

	Study group						Comparison test
	Tailored			Control			
	n	Median	IQR	n	Median	IQR	
IPAQ walk at discharge	12	635.2	1212.7	15	693.0	511.5	K-S Z=0.732, p=0.494, r=0.14
IPAQ walk at 1 month after discharge	10	594.0	726.0	12	544.5	730.1	K-S Z=0.272, p=0.995, r=0.06
IPAQ walk at 3 months after discharge	6	940.5	891.0	10	486.7	742.5	K-S Z=1.226, p=0.050, r=0.31
IPAQ moderate at discharge	12	240.0	1620.0	17	1200.0	2700.0	K-S Z=1.014, p=0.181, r=0.19
IPAQ moderate at 1 month after discharge	9	720.0	1070.0	12	0	1860.0	K-S Z=0.756, p=0.431, r=0.16
IPAQ moderate at 3 months after discharge	7	1440.0	2000.0	9	480.0	1080.0	K-S Z=0.976, p=0.209, r=0.24
IPAQ vigorous at discharge	12	840.0	3600.0	18	2400.0	2120.0	K-S Z=1.043, p=0.169, r=0.19
IPAQ vigorous at 1 month after discharge	10	1200.0	2340.0	11	480.0	1416.0	K-S Z=0.728, p=0.446, r=0.16
IPAQ vigorous at 3 months after discharge	6	2300.0	1824.0	11	0	1920.0	K-S Z=1.134, p=0.075, r=0.27

