

Guest Editorial

The WWW of the World Wide Web: Who, What, and Why?

John Powell, MSc, MRCPsych, MFPHM; Aileen Clarke, MD, MRCP, FFPHM

Health Services Research Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK

Corresponding Author:

John Powell, MSc, MRCPsych, MFPHM
Health Services Research Unit
London School of Hygiene and Tropical Medicine
Keppel Street
London WC1E 7HT
UK
Email: powell@lshtm.ac.uk

(*J Med Internet Res* 2002;4(1):e4) doi:[10.2196/jmir.4.1.e4](https://doi.org/10.2196/jmir.4.1.e4)

Current estimates suggest that 25 million people in the UK have access to the World Wide Web, and 14 million use it regularly [1]. Worldwide over 500 million people have logged on [2]. They have access to over 3 billion Web documents [3], and at least 2% of Web sites are health related [4]. Indeed, accessing health information is one of the commonest reasons for going on-line: surveys show that 50% to 75% of World Wide Web users have used it to look for health information [5,6,7], and those who do so access such information over 3 times a month [5]. In December 2001 the NHS Direct consumer health information Web site (www.nhsdirect.nhs.uk) dealt with 5.2 million hits from 171900 visitors [8].

Physicians are increasingly experiencing patients bringing Internet printouts to the consultation, although estimates of the frequency of this occurrence vary from 1-2% [9], to 58% [10], to over 70% [11]. The low prevalence of Internet-savvy patients of only 1-2% in the study by Potts and Wyatt [9], published in this issue of the *Journal of Medical Internet Research*, is surprising given the findings from consumer surveys on the frequency of accessing online health information that are cited above. Potts and Wyatt used a cross-sectional survey method asking respondents to retrospectively estimate the number of their patients who had used the Internet for health information in the past month. It is possible that recall bias may have led to an underestimate, but it is also likely that not all patients who consult the Internet reveal this to their doctor. The potential impact of the wide availability of online health information on the practitioner-patient relationship has been debated [12,13]. The Internet is a key influence in changing the balance of (knowledge) power between health care professionals and the public, empowering patients to become more involved in health care decision making and contributing to the deprofessionalization of medicine. Empirical research is beginning to investigate this impact [9,14].

Much of the limited evidence as to who the consumers of Internet health information are and what they are looking for comes from United States market-research surveys and Web-usage statistics, both quantifying the numbers of users and

types of information accessed. Women are more likely than men to seek health care information on-line, and the highest proportion of usage is in those between 30 and 64 years old [15]. Use of the Internet for health information declines with age [16,17]. Despite the much-discussed "digital divide" between the higher-income, more-educated "have-nets" and the lower-income, less-educated "have-nots," there is no evidence of differences in health-information seeking by income group once they have on-line access [18,19].

A 1999 Harris Poll of 2000 US adults found that mental health issues dominated the most popular on-line health topics, with depression, bipolar disorder, and anxiety problems accounting for 42% of the use of the Web to find health information [20]. Further work to investigate which health topics are most frequently accessed on-line will be valuable. Most on-line health seekers are looking for a specific answer to a specific health question, and start by submitting a topic to a general search engine [21]. Far fewer users go to health portals or direct to a specific health site, and in general, users do not simply browse for health-related information [22]. Most users research specific health issues that are currently affecting a friend, relative, or themselves, frequently in connection with a visit to their doctor [15]. Few use health sites to communicate with health services, purchase pharmaceuticals, or participate in health-related chat-room discussions [15]. However, the majority of US users report a desire for more on-line interaction with their doctors, including e-mail consultations and reminders [23].

The research in this area is notable for a relative lack of qualitative work exploring the reasons behind on-line information seeking and the attitudes and behavior of health users towards the World Wide Web. Sociological work has led to a better understanding of the process of help-seeking behavior, but this work now needs to be updated to take into account the use of the Internet by patients and caregivers.

Users report valuing the convenience, anonymity, and volume of online information [15]. It is likely that individuals will use the Web at different points in the trajectory of illness and health care. The California Healthcare Foundation has attempted to

categorize 3 types of user - the well; the newly diagnosed; and the chronically ill and their caregivers [24]. The well group carries out episodic searching for information relating to short-term medical conditions, pregnancy, and prevention issues. The newly diagnosed carries out very intensive searching for specific information, valuing the ease of access and broad range of information. The chronically ill and their caregivers carry out regular searching for information related to new treatments, nutrition advice, and alternative therapies. In addition, the latter 2 groups both value and use on-line communities and chat rooms. Several studies have shown the importance of the World Wide Web in providing social support, particularly to groups with chronic health problems such as diabetes patients [25, 26] or individuals with HIV [27].

It is likely that much of what is required from online information will be similar to that required from more-traditional routes:

clear, well-presented information, with advice on further sources. However, there may well also be particular advantages that can be gained from the interactivity, personalization, and creative ways of managing knowledge that the Internet provides. For example, preliminary work suggests that the Internet may be an effective means of delivering psychological therapies [28].

In an era of user involvement, consumer empowerment, and the wide dissemination of information on health and health services, it is important that we identify who the consumers of online health information are, what their information needs are, and understand why and how they seek information online. This will enable information to be provided in ways that will have benefits from the worldwide to the individual level, and will inform current debates over the quantity and quality of information provision and issues of privacy and access.

Acknowledgments

Dr Powell is funded by the NHS (South East) Research and Knowledge Management Programme, on a Training Fellowship in Health Services Research.

Conflicts of Interest

Dr Powell has worked with the consumer-health-information site NetDoctor.co.uk and the professional medical site www.Doctors.net.uk.

References

1. ; Nielsen/NetRatings Inc. Internet usage statistics for the month of December 2001 (United Kingdom). URL: <http://epm.netratings.com/uk/web/NRpublicreports.usagemonthly> [accessed 2002 Jan 24]
2. ; Nua Internet Surveys. How many online?. 2001 Aug. URL: http://www.nua.com/surveys/how_many_online/index.html [accessed 2002 Jan 24]
3. Google offers immediate access to 3 billion Web documents [press release]. Mountain View, Calif: Google Inc; 2001 Dec 11. URL: <http://www.google.com/press/pressrel/3billion.html> [accessed 2002 Feb 15]
4. Kiley R. Editorial - Internet Statistics. Health information on the Internet 1999;10:1-2 [FREE Full text]
5. Taylor H. The Harris Poll #19: Cyberchondriacs Update. 2001 Apr 18. URL: http://www.harrisinteractive.com/harris_poll/index.asp?PID=229 [accessed 2002 Feb 15]
6. Tatsumi H, Mitani H, Haruki Y, Ogushi Y. Internet medical usage in Japan: current situation and issues. J Med Internet Res 2001 Mar 17;3(1):e12 [FREE Full text] [Medline: 21578019]
7. Murero M, D'ancona G, Karamanoukian H. Use of the Internet by patients before and after cardiac surgery: telephone survey. J Med Internet Res 2001 Sep 30;3(3):e27 [FREE Full text] [Medline: 21578034] [doi: 10.2196/jmir.3.3.e27]
8. ; Kable's Government Computing. New year cheer for NHS Direct. 2002 Jan 4. URL: <http://www.kablenet.com/kd.nsf/printview/A33471B29784EDD480256B37003DC0EF?OpenDocument> [accessed 2002 Feb 15]
9. Potts HWW, Wyatt JC. Survey of doctors' experience of patients using the Internet. J Med Internet Res 2002 Mar 31;4(1):e5 [FREE Full text] [Medline: 21953333] [doi: 10.2196/jmir.4.1.e5]
10. Wilson SM. Impact of the internet on primary care staff in Glasgow. J Med Internet Res 1999 Nov 19;1(2):e7 [FREE Full text] [Medline: 21577981] [doi: 10.2196/jmir.1.2.e7]
11. Hjortdahl P, Nylenna M, Gerlow AO. Internett og lege-pasient-forholdet - fra "takk" til "hvorfor?". Tidsskr Norsk Laegeforening 1999;119:4339-4341.
12. Gerber BS, Eiser AR. The patient physician relationship in the Internet age: future prospects and the research agenda. J Med Internet Res 2001 Apr 9;3(2):e15 [FREE Full text] [Medline: 21578022] [doi: 10.2196/jmir.3.2.e15]
13. Hardey M. Doctor in the house: the Internet as a source of lay health knowledge and the challenge to expertise. Sociology of Health & Illness 1999;21(6):820-835. [doi: 10.1111/1467-9566.00185]
14. Nwosu C, Cox B. The impact of the Internet on doctor-patient relationships. Journal of Medical Informatics 2000;6(3):156-161.
15. Fox S, Rainie L; The Pew Internet & American Life Project. The online health care revolution: how the Web helps Americans take better care of themselves. 2000 Nov 26. URL: <http://www.pewinternet.org/reports/toc.asp?Report=26> [accessed 2002 Feb 15]

16. Smith-barbaro PA, Licciardone JC, Clarke HF, Coleridge ST. Factors associated with intended use of a Web site among family practice patients. *J Med Internet Res* 2001 May 17;3(2):e17 [FREE Full text] [Medline: [21578024](#)] [doi: [10.2196/jmir.3.2.e17](#)]
17. Licciardone JC, Smith-barbaro P, Coleridge ST. Use of the internet as a resource for consumer health information: results of the second osteopathic survey of health care in America (OSTEOSURV-II). *J Med Internet Res* 2001 Dec 26;3(4):e31 [FREE Full text] [Medline: [21635292](#)] [doi: [10.2196/jmir.3.4.e31](#)]
18. Brodie M, Flournoy RE, Altman DE, Blendon RJ, Benson JM, Rosenbaum MD. Health information, the Internet, and the digital divide. *Health Aff (Millwood)* 2000 Dec;19(6):255-265 [FREE Full text] [Medline: [21033799](#)] [doi: [10.1377/hlthaff.19.6.255](#)]
19. Pingree S, Hawkins RP, Gustafson DH, Boberg EW, Bricker E, Wise M, et al. Will the disadvantaged ride the information superhighway? Hopeful answers from a computer-based health crisis system. *J of Broadcasting and Electronic Media* 1996;40:331-353.
20. Taylor H. The Harris Poll #47: Explosive growth of "cyberchondriacs" continues. 1999 Aug 5. URL: http://www.harrisinteractive.com/harris_poll/index.asp?PID=117 [accessed 2002 Feb 15]
21. Lovich D, Silverstein MB, Lesser R. Vital signs: the impact of e-health on patients and physicians. Boston Consulting Group. 2001 Feb. URL: http://www.bcg.com/publications/files/Summary_Vital_Signs_5_01_report.pdf [accessed 2002 Feb 15]
22. Poensgen A, Larsson S. Patients, physicians and the Internet: myth, reality and implications. Boston Consulting Group. 2001 Jan. URL: http://www.bcg.com/publications/files/Patients_Physicians_and_the_Internet_Jan_01_summary.pdf [accessed 2002 Feb 15]
23. ; Cyber Dialogue. *Cybercitizen Health* 2001. New York: Cyber Dialogue; 2001.
24. Kahn J, Wayne JC. Health e-people: the online consumer experience. Oakland, California: California Healthcare Foundation and the Institute for the Future; 2000. URL: <http://admin.chcf.org/documents/ehealth/HealthEPeople.pdf> [accessed 2002 Feb 15]
25. Glasgow RE, Barrera M, McKay HG, Boles SM. Social support and health among participants in an Internet-based diabetes support program: a multi-dimensional investigation. *Diabetes* 1998;47:1263.
26. Zrebiec JF, Jacobson AM. What attracts patients with diabetes to an internet support group? A 21-month longitudinal website study. *Diabet Med* 2001 Feb;18(2):154-158. [Medline: [21150039](#)] [doi: [10.1046/j.1464-5491.2001.00443.x](#)]
27. Reeves PM. Coping in cyberspace: the impact of Internet use on the ability of HIV-positive individuals to deal with their illness. *J Health Commun* 2000;5 Suppl(5 Suppl):47-59. [Medline: [20394543](#)] [doi: [10.1080/10810730050019555](#)]
28. Christensen H, Griffiths KM, Korten A. Web-based cognitive behavior therapy: analysis of site usage and changes in depression and anxiety scores. *J Med Internet Res* 2002 Feb 15;4(1):e3 [FREE Full text] [Medline: [21953331](#)] [doi: [10.2196/jmir.4.1.e3](#)]

Edited by G. Eysenbach; submitted 04.02.02; peer-reviewed by G Eysenbach; comments to author 04.02.02; accepted 08.02.02; published 18.02.02

Please cite as:

Powell J, Clarke A

The WWW of the World Wide Web: Who, What, and Why?

J Med Internet Res 2002;4(1):e4

URL: <http://www.jmir.org/2002/1/e4/>

doi: [10.2196/jmir.4.1.e4](#)

PMID: [11956036](#)

© John Powell, Aileen Clarke. Originally published in the *Journal of Medical Internet Research* (<http://www.jmir.org>), 18.2.2002. Except where otherwise noted, articles published in the *Journal of Medical Internet Research* are distributed under the terms of the Creative Commons Attribution License (<http://www.creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited, including full bibliographic details and the URL (see "please cite as" above), and this statement is included.