

Original Paper

# Crystal Methamphetamine Use Predicts Incident STD Infection Among Men Who Have Sex With Men Recruited Online: A Nested Case-Control Study

Sabina Hirshfield<sup>1</sup>, PhD; Robert H Remien<sup>2</sup>, PhD; Imelda Walavalkar<sup>1</sup>, BA; Mary Ann Chiasson<sup>1</sup>, DrPH

<sup>1</sup>Medical and Health Research Association of New York City Inc, New York NY, USA

<sup>2</sup>HIV Center for Clinical and Behavioral Studies, New York State Psychiatric Institute and Columbia University, New York NY, USA

**Corresponding Author:**

Sabina Hirshfield, PhD

Medical and Health Research Association of New York City, Inc

40 Worth Street, Suite 720

New York NY 10013

USA

Phone: +1 212 285 0220 ext 115

Fax: +1 212 385 0565

Email: [shirshfield@mhra.org](mailto:shirshfield@mhra.org)

---

## Abstract

**Background:** Among men who have sex with men (MSM), the number of newly diagnosed human immunodeficiency virus (HIV) infections has increased by approximately 60% since 1999. Factors that may be contributing to this resurgence include a widely reported increase in bacterial sexually transmitted diseases (STDs) among HIV-positive and HIV-negative MSM, as well as unsafe sexual practices.

**Objective:** This research was undertaken to learn more about risk behaviors associated with an incident STD among MSM.

**Methods:** A nested case-control study was conducted, using data from a cross-sectional Internet survey of MSM (N=2643), which investigated risk behaviors during a 6-month period in 2001. Chi-square and logistic regression methods were used to estimate the likelihood of acquiring an incident STD versus no STD.

**Results:** Eighty-five percent of the respondents were white, 46% were under age 30, and 80% had met sex partners online; 7% were HIV-positive. Men with an incident STD were more likely than men without an STD to report drug use (crystal methamphetamine odds ratio 3.8; 95% confidence interval 2.1-6.7; cocaine OR 2.3; 95% CI 1.2-4.2; ecstasy OR 2.2; 95% CI 1.3-3.8; Viagra OR 2.1; 95% CI 1.2-3.7), alcohol before or during sex (OR 1.9; 95% CI 1.2-2.9), and high-risk sexual behavior (unprotected anal intercourse OR 5.0; 95% CI 2.8-8.9; multiple sex partners OR 5.9; 95% CI 2.5-13.8). In the multivariate analysis, significant independent predictors associated with an incident STD were crystal methamphetamine use (adjusted OR 2.0; 95% CI 1.1-3.8), unprotected anal intercourse (adjusted OR 3.4; 95% CI 1.9-6.3), and 6 or more sex partners during the study period (adjusted OR 3.3; 95% CI 1.4-7.8).

**Conclusion:** Identifying and treating MSM who have STDs, or who are at increased risk for acquiring STDs, is crucial in preventing the further spread of disease. In addition, there is a need to integrate HIV/STD and substance use prevention and education into Web-based and community-based venues.

(*J Med Internet Res* 2004;6(4):e41) doi:[10.2196/jmir.6.4.e41](https://doi.org/10.2196/jmir.6.4.e41)

---

## KEYWORDS

Internet; sexually transmitted diseases; methamphetamine; HIV

---

## Introduction

Among men who have sex with men (MSM), the number of newly diagnosed human immunodeficiency virus (HIV) infections has increased by approximately 60% since 1999 [1]. Several factors may be contributing to the increase in HIV


transmission. One is the widely reported increase in bacterial sexually transmitted diseases (STDs), namely syphilis and gonorrhea, among MSM [2,3]. Not only are STDs a marker for unsafe sexual behavior, but ulcerative and non-ulcerative STDs facilitate the transmission and acquisition of HIV [4,5] and increase HIV viral load and infectivity in persons with HIV

[6,7]. Moreover, studies have found a high proportion of HIV-positive MSM with incident STDs [8,9], suggesting continued unsafe sexual practices and exposure of others to HIV. Substance use has also been associated with sexual risk behaviors among MSM [10-12].

Crystal methamphetamine (crystal) use in MSM communities has been problematic in the Western US since the early 1990s [13,14], and has more recently spread to the Midwest [15], as well as the East Coast [16]. Crystal use is associated with "marathon sex" (prolonged sexual activity), receptive and insertive anal sex without a condom, the ability to have sustained arousal for multiple partners, and unsafe sex with HIV-serodiscordant partners or partners of unknown HIV serostatus [15,17,18]. Thus, substance use and its relationship to high-risk sexual behavior among MSM is of particular concern, as drugs may help men to avoid feelings of anxiety associated with same-sex behavior and their own awareness of HIV risk [17,19,20].

This research was undertaken to learn more about risk behaviors associated with an incident STD among MSM. We compared sexual and drug use behaviors between men with a self-reported incident bacterial or viral STD and men without an STD.

**Figure 1.** Survey banner ad



In order to minimize non-valid data, we incorporated reliability checks into the survey for age and certain risk behaviors. To reduce the likelihood of participants' completing multiple surveys, the study banner was rotated through the online chat rooms approximately every 20 minutes. Also, it was not technically possible for participants to bookmark the survey, and there were no monetary incentives to complete the survey.

The general interest, gay-oriented website agreed to host the banner in all of its US adult chat rooms. Individuals had to be registered with the website in order to enter chat rooms. The chat room banner provided the only link to the survey. No personally identifying information was collected. The survey did not use cookies and neither collected user IP addresses nor stored them with submitted data. Study participation was limited to those 18 and older, and all participants clicked on an online consent form before gaining access to the anonymous survey. The Medical and Health Research Association of New York City, Inc. (MHRA) institutional review board approved the study.

The survey included information on demographics (age group, race/ethnicity, education, income and residence), and assessment

## Methods

We conducted an anonymous, cross-sectional Internet study, inquiring about sexual and drug-using behaviors among MSM between June and December 2001, as part of a larger study of behavior change pre- and post-September 11, 2001. The banner ad (Figure 1) linking to the survey was posted in the online chat rooms of a general interest, gay-oriented website between June 3 and July 24, 2002. Overall, 2284 individuals clicked on the banner but exited the survey without answering any questions; 3697 clicked on the banner and answered the survey. A total of 2949 questionnaires were complete enough for statistical analysis (79% completion rate): 2934 were completed by men (18 of whom were exclusively heterosexual), 10 by women, and 5 by transgendered individuals. Analysis was limited to the 2643 men who reported sex with other men or who self-identified as gay or bisexual, excluding those who never had sex, those who were missing information on STDs, and those who had chronic viral STDs. To assess incident STDs, the questionnaire asked if the respondent had been diagnosed with any STDs during two consecutive 3-month periods between June and December 2001 and provided a checklist of the most common infections. For men reporting viral STDs, we included only viral STDs that were reported in the second 3-month period. Overall, 102 (4%) men reported being diagnosed with an incident bacterial or viral STD during the 6-month study period.

of risk behaviors, such as type of sexual contact (anal, oral, vaginal; with and without condoms) with main and non-main partners, knowledge of partners' HIV status, type of illicit drug use before or during sex, alcohol use before or during sex, how sex partners were met, and HIV testing. Links to STD prevention/treatment websites and mental health hotlines appeared at the end of the survey. Survey questions were adapted from questionnaires used by the investigators in previous studies.

Data analyses were conducted using SPSS 9.0 for Windows [21]. Bivariate categorical data were evaluated using chi-square and odds ratios. Statistically significant bivariate analyses were simultaneously assessed by multiple logistic regression models. To guard against Type I error, we set the *P*-value to .01 in the bivariate analyses, given that controls outnumbered cases almost 25 to 1.

Respondents were asked how many sex partners they had during two distinct 3-month periods. Respondents could only choose one response from a pull-down menu for each time period. Answer choices were none, 1, 2-5, 6-10, 11-20, 21-50, and 51 or higher. This variable was collapsed for the entire 6-month period; men who reported no partners or one were grouped into

the first category. Men who reported 2-5 partners were grouped into the second category, and men who reported 6 partners or more were grouped into the third. For this analysis, "multiple sex partners" refers to 2 or more partners during the study period. Regarding unprotected anal intercourse (UAI), respondents were asked about insertive and receptive sex without a condom. The UAI variable represents men who reported any unprotected receptive and/or insertive anal intercourse. Age was categorized in a pull-down menu: 18-24, 25-29, 30-39, 40-49, 50-59, 60 and older. For ease of analysis, age was collapsed into three categories.

## Results

Overall, the study sample was representative of the host website user population. The host website was able to provide several demographic variables on new registrants from the entire site for the month prior to sample recruitment. Most new registrants were male (87%), and the study sample was identical to the site population in age, and similar in educational attainment and by US region. Although two of the demographic variables were significantly different, the findings may be a reflection of the large samples sizes, which can detect small differences.

**Table 1.** Characteristics of study sample and host Internet website

	Study N (%)	Host Website N (%)	P *
<b>Age †</b>	n=2599	n=10 124	
18-24	730 (28)	3090 (29)	
25-34	809 (31)	3091 (29)	
35-44	635 (24)	2451 (23)	
45-54	331 (13)	1172 (11)	
55-64	80 (3)	320 (3)	
65 and older (reference)	14 (<1)	74 (<1)	0.168
<b>Education</b>	n=2633	n=6394	
High school or less (reference)	335 (13)	939 (15)	
Some college	1093 (41)	2578 (40)	
College degree or more	1205 (46)	2877 (45)	0.050
<b>US Regional Breakdown</b>	n=2874	n=8846	
Northeast	472 (16)	1658 (19)	
CT, ME, MA, NH, RI, VT, NJ, NY, PA			
Midwest	661 (23)	1998 (22)	
IN, IL, MI, OH, WI, IA, KS, MN, MO, NE, ND, SD			
South	1026 (36)	2910 (33)	
DE, DC, FL, GA, MD, NC, SC, VA, WV, AL, KY, MS, TN, AR, LA, OK, TX			
West (Reference)	715 (25)	2280 (26)	0.006
AZ, CO, ID, NM, MT, UT, NV, WY, AK, CA, HI, OR, WA			

\* Chi-square goodness-of-fit statistic used.

† Age brackets were recategorized in order to compare to the host website.

Participants resided in all 50 states, roughly in proportion to the population of each state. Less than 1% resided in Guam, Puerto Rico, and a few locations outside the United States. Approximately half (46%) of the study participants were younger than 30 and had at least a college degree (46%). Most were white (85%). Overall, 6% reported crystal use, 7% reported cocaine, 9% reported ecstasy, 9% reported Viagra, and about half (48%) reported drinking alcohol before or during sex. The aforementioned drugs were commonly used before or during sex (over 85% reported these drugs before or during sex). Most (80%) reported meeting new sexual partners online, and most (80%) engaged in sex with multiple partners. The majority

(81%) engaged in sex exclusively with men, and 7% were HIV-positive.

Respondents reported newly diagnosed bacterial or viral STDs (n=102), which included syphilis (n=9), genital herpes (n=4), genital warts/anal warts/HPV (n=16), gonorrhea (n=49), hepatitis B (n=2), chlamydia (n=29), and non-gonococcal urethritis (n=24). Sixteen men reported 2 STDs, and 7 reported 3 or more. Many respondents in the STD group made an effort to notify partners of potential exposure. Approximately 30% notified all partners, 26% told some partners but not all, 17% tried to notify their partners but could not locate them, and less than 5% had

the health department notify their sex partners; 21% told none of their partners.

Men with new STDs were more likely to be between 30 and 39 years of age than the controls (see Table 2). The bivariate and multivariate analyses of risk correlates for STDs were structured by drug use and behavioral risk categories as there were no demographic differences between cases and controls. Cases

were significantly more likely to report drug use before or during sex (crystal, cocaine, ecstasy, and Viagra), alcohol use before or during sex, and sexual risk behaviors (ie, UAI and multiple sex partners) than the controls (see Table 2). Gamma hydroxy butyrate (GHB), poppers (nitrite inhalants), ketamine, and marijuana use were excluded from the analyses, as their use was not statistically different between groups.

**Table 2.** Comparison of demographic and behavioral characteristics of men with incident STDs and controls

(N=2643)	STD	Controls	P*
Demographics	N (%)	N (%)	
<b>Age</b>	<b>(n=102)</b>	<b>(n=2541)</b>	
18-29	51 (50)	1167 (46)	.073
30-39	33 (32)	695 (27)	.050
40+ (reference group)	18 (18)	679 (27)	--
<b>Race/Ethnicity</b>			
White	81 (82)	2126 (85)	.316
Black	3 (3)	51 (2)	.894
Hispanic	5 (5)	140 (6)	.464
Other/mixed race (reference group)	10 (10)	186 (7)	--
<b>Education</b>			
High school or less (reference group)	10 (10)	325 (13)	--
Some college	44 (44)	1049 (41)	.384
College degree or more	47 (46)	1158 (46)	.434
<b>Income</b>			
Up to \$40 000	58 (64)	1375 (60)	.453
\$41 000 or more	33 (36)	924 (40)	
<b>Met Partners Online</b>			
Yes	87 (87)	1954 (80)	.080
No	13 (13)	492 (20)	
<b>HIV Status</b>			
Positive	11 (11)	180 (7)	.157
Negative or unknown	91 (89)	2361 (93)	

\* Age, race, and education used logistic regression to calculate the *P*-value. Income, meeting partners online, and HIV status used chi-square to calculate the *P*-value.

To test for multicollinearity, we ran a linear regression with “any STD” as the dependent variable and the drug and behavioral risk variables from the bivariate analyses as the independent variables. The variance inflation factor (VIF) value for each variable was below 1.5, indicating that multicollinearity was not present. We separated the drug and behavioral risk variables for the multivariate logistic analyses into 3 logistic regression models to assess risk correlates for acquiring an incident STD: model 1 comprised crystal, cocaine, ecstasy, Viagra, and alcohol before or during sex; model 2 comprised UAI and number of sex partners during the study; and model 3 (see Table 3) comprised the significant variables from models 1 and 2.

In model 1, only crystal and alcohol before/during sex were predictive of acquiring an incident STD (crystal, OR 2.7, 95% CI 1.2-6.0, *P*<.05; alcohol, OR 1.6, 95% CI 1.0-2.6, *P*<.05). In model 2, UAI and having 6 or more sex partners during the study were predictive of acquiring an incident STD (UAI OR 3.9, 95% CI 2.2-7.1, *P*<.001; 6 or more partners, OR 4.3, 95% CI 1.8-10.1, *P*=.001). In the final multivariate model (see Table 3), alcohol before/during sex lost significance and UAI, crystal use before or during sex, and having 6 or more sex partners were the strongest predictors of acquiring an incident STD.

In order to assess the potential for HIV transmission, we compared the HIV status of the participants to that of their partners. Among HIV-positive men with multiple sex partners who reported UAI (n=109), 47% reported UAI with

HIV-negative/unknown partners only, 43% reported UAI with positive and negative/unknown partners, and 10% reported UAI with positive partners only. Seven of the 8 respondents with an STD in this subgroup reported sex with serodiscordant partners.

**Table 3.** Bivariate and multivariate analyses: factors associated with incident STDs

	Incident STD #		Drug Use and Behavioral Risk		Crystal and Behavioral Risk	
	Yes	No	Bivariate Odds Ratio (95% CI)	<i>P</i>	Multivariate Adjusted Odds Ratio* (95% CI)	<i>P</i>
	N (%)	N (%)				
<b>Drug Use Before/During Sex §</b>	n=94	n=2411				
Crystal methamphetamine	16 (17)	124 (5)	3.8 (2.1-6.7)	<.001	2.0 (1.1-3.8)	.024
Cocaine	13 (14)	159 (7)	2.3 (1.2-4.2)	.007		
Ecstasy	17 (18)	219 (9)	2.2 (1.3-3.8)	.003		
Viagra	16 (17)	211 (9)	2.1 (1.2-3.7)	.006		
<b>Alcohol Use</b>	n=98	n=2499				
Alcohol before sex†	62 (63)	1192 (48)	1.9 (1.2-2.9)	.002	1.3 (0.8-2.1)	.207
<b>Behavioral Risk</b>	n=102	n=2538				
Unprotected anal intercourse‡	88 (86)	1409 (55)	5.0 (2.8-8.9)	<.001	3.4 (1.9-6.3)	<.001
<b>Sex partners</b>	n=100	n=2494				
0-1 (reference)	6 (6)	503 (20)	--			
2-5	31 (31)	1103 (44)	2.3 (0.9-5.7)	.056	1.6 (0.6-3.9)	.294
6-100+	63 (63)	888 (36)	5.9 (2.5-13.8)	<.001	3.3 (1.4-7.8)	.007

\* Adjusted odds ratio = the odds ratio estimated after adjusting for all other variables included in the parsimonious model.

‡ Note: In model 1 (data not shown), crystal use and alcohol before/during sex were associated with incident STDs. In model 2 (data not shown), UAI and having 6 or more sex partners were associated with incident STDs.

† Sometimes/most of the time

‡ Receptive and/or insertive UAI

§ Drug use variables are not mutually exclusive

## Discussion

In this case-control study of men recruited through the Internet, strong associations were found between unprotected anal intercourse, crystal use, and multiple sex partners and an incident STD. In the overall sample, 4% reported a diagnosis of an incident bacterial or viral STD during the 6-month study period. The great majority of HIV-positive men with multiple sex partners reported unprotected sex with HIV-negative or status unknown partners, which signifies the continued risk of spreading HIV and other STDs to non-infected individuals [8]. Other studies of HIV-positive men report a range of serodiscordant or potentially discordant sex, from 21% to 49% [18,22,23]. An average of 80% of our sample met sex partners online, and study findings indicate risk comparable to other Web-based studies on recent sexual risk behavior trends among MSM [24-26].

Men who begin having sex with men while on drugs may develop a pattern of using drugs during sexual experiences [27], and certain drugs such as nitrite inhalants (poppers) and crystal may be used specifically to enhance sexual experiences [27]. Impaired judgment due to drug use may lead to unprotected

sex, increasing the risk of HIV/STD transmission [13]. It has been hypothesized that substance use may help men avoid feelings of anxiety associated with same-sex behavior and concerns about HIV risk [20]. Reback's report [17] found that MSM used crystal to cope with negative internal messages about gay sexuality, and HIV-positive MSM reported using it to cope with the fear of transmitting HIV. Reback's report also found that most HIV-positive participants reported that they did not disclose their HIV status to casual sex partners as it was their partner's responsibility to use protection or to set behavioral limits.

Certain limitations of this study deserve mention. Our survey was posted on only one gay-oriented website. We do not know whether survey respondents would differ if the survey had been posted on multiple sites or on sites that specifically facilitate meeting sex partners. Minority MSM were underrepresented in the sample; our data suggest that white, non-Hispanic MSM were unintentionally oversampled, as those who have computer skills and access to participate in online sex surveys tend to be younger, wealthier, educated white males [28-30]. STD was self-report only and we did not ask for the site of infection. There may have been underreporting in this sample, as certain

STDs, like chlamydia and gonorrhea, are often asymptomatic and go undetected and unreported [31]. This may also be true for certain viral STDs such as genital herpes [32]. Finally, it is not possible to determine whether respondents who participated in this Internet-based survey are representative of MSM who use the Internet, of MSM in general, or of MSM with HIV, since the MSM population has never been enumerated. Despite these limitations, Internet research is an efficient and inexpensive way to reach large samples of high-risk groups.

Identifying and treating MSM who have STDs, or who are at increased risk of acquiring STDs, is crucial in preventing the further spread of disease. The Internet is a necessary and appropriate medium to reach sex-seeking populations for prevention and intervention efforts [33], especially when factoring in increasing numbers of people living with HIV who are resuming sexual activity as a result of improved treatment regimens [34]. Just as bathhouses and shooting galleries have been used to deliver STD prevention messages, Internet-based interventions should be considered for those seeking sex online [35]. Results of preliminary research on Internet HIV prevention for MSM are promising, suggesting that the Internet may be a

reliable resource for studying and targeting risk behaviors in MSM [36].

Studies conducted over the past 20 years have found associations between substance abuse treatment and a reduction in HIV risk behaviors [37]. Primary and secondary substance abuse treatment among MSM has been successful, as treatment can affect decisions about sexual behavior uninfluenced by drugs and alcohol [38]. However, treatment remains challenging, and it has been suggested that men need to abstain from drug use and learn skills to meet and initiate sex with men while sober [12]. The rise in crystal use among the MSM population may require a special focus on current substance abuse treatment approaches, such as addressing drugs in exchange for sex, and disclosure of HIV status [17]. In addition, there is a need to integrate HIV/STD and substance use prevention and education into Web-based and community-based venues. Study findings raise questions concerning the spread of disease and the multiple high-risk behaviors, specifically, how drug use is situated within the trajectory to unprotected sex, multiple partners, and ultimately HIV/STD transmission. Additional data are needed to better understand specific pathways between sexual and drug using practices and HIV/STD transmission among MSM.

## Acknowledgments

Data analysis and manuscript preparation were funded in part through CDC Contract Number 200-97-0621, Task 33 to RTI International, and Subcontract Number 10-46U-6900 from RTI to Medical and Health Research Association of New York City, Inc. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the US Government.

We would like to thank Mike Humberstone of Data Link, MHRA, Inc. for his technical and programming support.

## Conflicts of Interest

None declared.

## References

1. ; Centers for Disease Control and Prevention (CDC). Increases in HIV diagnoses--29 States, 1999-2002. *MMWR Morb Mortal Wkly Rep* 2003 Nov 28;52(47):1145-1148 [[FREE Full text](#)] [Medline: [14647015](#)]
2. ; Centers for Disease Control and Prevention. Primary and secondary syphilis among men who have sex with men - New York City, 2001. *Morb Mortal Wkly Rep* 2002;51(38):853-856 [[FREE Full text](#)] [Medline: [12363336](#)]
3. Rietmeijer CA, Patnaik JL, Judson FN, Douglas JM. Increases in gonorrhea and sexual risk behaviors among men who have sex with men: a 12-year trend analysis at the Denver Metro Health Clinic. *Sex Transm Dis* 2003 Jul;30(7):562-567. [Medline: [22722350](#)]
4. ; Centers for Disease Control and Prevention. HIV prevention through early detection and treatment of other sexually transmitted diseases - United States recommendations of the Advisory Committee for HIV and STD Prevention. *MMWR Recomm Rep* 1998 Jul 31;47(RR-12):1-24 [[FREE Full text](#)] [Medline: [98365274](#)]
5. Dickerson MC, Johnston J, Delea TE, White A, Andrews E. The causal role for genital ulcer disease as a risk factor for transmission of human immunodeficiency virus. An application of the Bradford Hill criteria. *Sex Transm Dis* 1996;23(5):429-440. [Medline: [97039525](#)]
6. Mertz KJ, Weiss JB, Webb RM, Levine WC, Lewis JS, Orle KA, et al. An investigation of genital ulcers in Jackson, Mississippi, with use of a multiplex polymerase chain reaction assay: high prevalence of chancroid and human immunodeficiency virus infection. *J Infect Dis* 1998 Oct;178(4):1060-1066. [Medline: [99022982](#)] [doi: [10.1086/314502](#)]
7. Mbopi-kéou FX, Grésengué G, Mayaud P, Weiss HA, Gopal R, Matta M, et al. Interactions between herpes simplex virus type 2 and human immunodeficiency virus type 1 infection in African women: opportunities for intervention. *J Infect Dis* 2000 Oct;182(4):1090-1096. [Medline: [20435900](#)] [doi: [10.1086/315836](#)]
8. Erbeling EJ, Chung SE, Kamb ML, Irwin KL, Rompalo AM. New sexually transmitted diseases in HIV-infected patients: markers for ongoing HIV transmission behavior. *J Acquir Immune Defic Syndr* 2003 Jun 1;33(2):247-252. [Medline: [22679045](#)]

9. Scheer S, Chu PL, Klausner JD, Katz MH, Schwarcz SK. Effect of highly active antiretroviral therapy on diagnoses of sexually transmitted diseases in people with AIDS. *Lancet* 2001 Feb 10;357(9254):432-435. [Medline: [21119874](#)] [doi: [10.1016/S0140-6736\(00\)04007-1](#)]
10. Colfax GN, Mansergh G, Guzman R, Vittinghoff E, Marks G, Rader M, et al. Drug use and sexual risk behavior among gay and bisexual men who attend circuit parties: a venue-based comparison. *J Acquir Immune Defic Syndr* 2001 Dec 1;28(4):373-379. [Medline: [21564241](#)]
11. Mansergh G, Colfax GN, Marks G, Rader M, Guzman R, Buchbinder S. The Circuit Party Men's Health Survey: findings and implications for gay and bisexual men. *Am J Public Health* 2001 Jun;91(6):953-958. [Medline: [21285318](#)]
12. Stall R, Purcell D. Intertwining epidemics: a review of research on substance use among men who have sex with men and its connection to the AIDS epidemic. *AIDS and Behavior* 2000;4(2):181-192. [doi: [10.1023/A:1009516608672](#)]
13. Halkitis PN, Parsons JT, Stirratt MJ. A double epidemic: crystal methamphetamine drug use in relation to HIV transmission among gay men. *J Homosex* 2001;41(2):17-35. [Medline: [21374611](#)] [doi: [10.1300/J082v41n02\\_02](#)]
14. Semple SJ, Patterson TL, Grant I. Binge use of methamphetamine among HIV-positive men who have sex with men: pilot data and HIV prevention implications. *AIDS Educ Prev* 2003 Apr;15(2):133-147. [Medline: [22624715](#)] [doi: [10.1521/aeap.15.3.133.23835](#)]
15. Semple SJ, Patterson TL, Grant I. Motivations associated with methamphetamine use among HIV+ men who have sex with men. *J Subst Abuse Treat* 2002 Apr;22(3):149-156. [Medline: [22035746](#)] [doi: [10.1016/S0740-5472\(02\)00223-4](#)]
16. Halkitis P. The crystal meth-HIV connection. HIV Forum NYC. New York; 2004. URL: <http://www.amfar.org/cgi-bin/iowa/news/feat/record.html?record=158> [accessed 2004 Nov 20]
17. Reback C, Ditman D. The social construction of a gay drug: methamphetamine use among gay and bisexual males in Los Angeles. Executive Summary. Los Angeles: City of Los Angeles, AIDS Coordinator; 1997.
18. Whittington WLH, Collis T, Dithmer-schreck D, Handsfield HH, Shalit P, Wood RW, et al. Sexually transmitted diseases and human immunodeficiency virus-discordant partnerships among men who have sex with men. *Clin Infect Dis* 2002 Oct 15;35(8):1010-1017. [Medline: [22241421](#)] [doi: [10.1086/342693](#)]
19. Mckirnan DJ, Ostrow DG, Hope B. Sex, drugs and escape: a psychological model of HIV-risk sexual behaviours. *AIDS Care* 1996 Dec;8(6):655-669. [Medline: [97146860](#)] [doi: [10.1080/09540129650125371](#)]
20. Mckirnan DJ, Vanable PA, Ostrow DG, Hope B. Expectancies of sexual "escape" and sexual risk among drug and alcohol-involved gay and bisexual men. *J Subst Abuse* 2001;13(1-2):137-154. [Medline: [21430446](#)] [doi: [10.1016/S0899-3289\(01\)00063-3](#)]
21. ; SPSS Inc. SPSS. Chicago, IL; 1999. URL: <http://www.spss.com/> [accessed 2004 Nov 20]
22. Chen SY, Gibson S, Weide D, McFarland W. Unprotected anal intercourse between potentially HIV-serodiscordant men who have sex with men, San Francisco. *J Acquir Immune Defic Syndr* 2003 Jun 1;33(2):166-170. [Medline: [22679032](#)]
23. Halkitis PN, Parsons JT. Intentional unsafe sex (barebacking) among HIV-positive gay men who seek sexual partners on the internet. *AIDS Care* 2003 Jun;15(3):367-378. [Medline: [22629792](#)] [doi: [10.1080/0954012031000105423](#)]
24. Kim AA, Kent C, McFarland W, Klausner JD. Cruising on the Internet highway. *J Acquir Immune Defic Syndr* 2001 Sep 1;28(1):89-93. [Medline: [21462991](#)]
25. McFarlane M, Bull SS, Rietmeijer CA; The Internet as a newly emerging risk environment for sexually transmitted diseases. *JAMA* 2000 Jul 26;284(4):443-446. [Medline: [20366357](#)] [doi: [10.1001/jama.284.4.443](#)]
26. Bull SS, McFarlane M, Rietmeijer C. HIV and sexually transmitted infection risk behaviors among men seeking sex with men on-line. *Am J Public Health* 2001 Jun;91(6):988-989. [Medline: [21285325](#)]
27. Lewis LA, Ross MW. A Select Body: The Gay Dance Party Subculture and the HIV-AIDS Pandemic. New York: Cassell; Mar 1, 1996.
28. Binik Y, Mah K, Kiesler S. Ethical issues in conducting sex research on the Internet. *J Sex Res* 1999;36(1):82-90.
29. Lenhart A, Horrigan J, Rainie L, Allen K, Boyce A, Madden M, et al. Ever-Shifting Internet Population: A New Look at Internet Access and the Digital Divide. Washington, DC: Diane Pub Co; Oct 1, 2003. URL: [http://www.pewinternet.org/pdfs/PIP\\_Shifting\\_Net\\_Pop\\_Report.pdf](http://www.pewinternet.org/pdfs/PIP_Shifting_Net_Pop_Report.pdf)
30. Toomey KE, Rothenberg RB. Sex and cyberspace-virtual networks leading to high-risk sex. *JAMA* 2000 Jul 26;284(4):485-487. [Medline: [20366365](#)] [doi: [10.1001/jama.284.4.485](#)]
31. Turner CF, Rogers SM, Miller HG, Miller WC, Gribble JN, Chromy JR, et al. Untreated gonococcal and chlamydial infection in a probability sample of adults. *JAMA* 2002 Feb 13;287(6):726-733. [Medline: [21841467](#)] [doi: [10.1001/jama.287.6.726](#)]
32. Langenberg AG, Corey L, Ashley RL, Leong WP, Straus SE. A prospective study of new infections with herpes simplex virus type 1 and type 2. Chiron HSV Vaccine Study Group. *N Engl J Med* 1999 Nov 4;341(19):1432-1438 [FREE Full text] [Medline: [20001875](#)] [doi: [10.1056/NEJM199911043411904](#)]
33. Ross MW, Tikkanen R, Månsson SA. Differences between Internet samples and conventional samples of men who have sex with men: implications for research and HIV interventions. *Soc Sci Med* 2000 Sep;51(5):749-758. [Medline: [20426946](#)] [doi: [10.1016/S0277-9536\(99\)00493-1](#)]
34. Rietmeijer CA, Bull SS, McFarlane M. Sex and the internet. *AIDS* 2001 Jul 27;15(11):1433-1434. [Medline: [21395680](#)] [doi: [10.1097/00002030-200107270-00013](#)]

35. Bull SS, McFarlane M, King D. Barriers to STD/HIV prevention on the Internet. *Health Educ Res* 2001 Dec;16(6):661-670. [Medline: [21639061](#)] [doi: [10.1093/her/16.6.661](#)]
36. Rhodes S, Glorioso J, Hergenrather K. Use of Internet chat rooms for HIV prevention among men who have sex with men in the United States: an evaluation of a community-based initiative. 2002 Presented at: XIV International Conference on AIDS; Jul 7-12; Barcelona, Spain.
37. Metzger DS, Navaline H. Human immunodeficiency virus prevention and the potential of drug abuse treatment. *Clin Infect Dis* 2003;37(Suppl 5):S451-S456. [Medline: [14648463](#)]
38. Shoptaw S, Frosch D. Substance abuse treatment as HIV prevention for men who have sex with men. *AIDS Behav* 2000;4(2):193-203. [doi: [10.1023/A:1009568725510](#)]

## Abbreviations

**HIV:** Human immunodeficiency virus

**MSM:** Men who have sex with men

**STD:** Sexually transmitted disease

**UAI:** Unprotected anal intercourse

*Edited by G. Eysenbach; submitted 17.03.04; peer-reviewed by M McFarlane, G Rebchook; comments to author 25.05.04; revised version received 09.07.04; accepted 19.08.04; published 29.11.04*

*Please cite as:*

*Hirshfield S, Remien RH, Walavalkar I, Chiasson MA*

*Crystal Methamphetamine Use Predicts Incident STD Infection Among Men Who Have Sex With Men Recruited Online: A Nested Case-Control Study*

*J Med Internet Res* 2004;6(4):e41

URL: <http://www.jmir.org/2004/4/e41/>

doi: [10.2196/jmir.6.4.e41](#)

PMID: [15631965](#)

© Sabina Hirshfield, Robert H Remien, Imelda Walavalkar, Mary Ann Chiasson. Originally published in the *Journal of Medical Internet Research* (<http://www.jmir.org>), 29.11.2004. Except where otherwise noted, articles published in the *Journal of Medical Internet Research* are distributed under the terms of the Creative Commons Attribution License (<http://www.creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited, including full bibliographic details and the URL (see "please cite as" above), and this statement is included.