

Original Paper

A Web-Based Nutrition Program Reduces Health Care Costs in Employees With Cardiac Risk Factors: Before and After Cost Analysis

Naomi Sacks¹, MA; Howard Cabral¹, PhD; Lewis E Kazis¹, ScD; Kelli M Jarrett¹, MPH; Delia Vetter²; Russell Richmond³, MD; Thomas J Moore⁴, MD

¹Boston University School of Public Health, Boston, MA, USA

²EMC Corporation, Boston, MA, USA

³Verisk Health, Inc, Boston, MA, USA

⁴Boston University School of Medicine, Boston, MA, USA

Corresponding Author:

Naomi Sacks, MA

Boston University School of Public Health

Department of Health Policy and Management

715 Albany Street

Boston, MA 02118

USA

Phone: +1 617 638 4640

Fax: +1 617 638 5374

Email: nsacks@bu.edu

Abstract

Background: Rising health insurance premiums represent a rapidly increasing burden on employer-sponsors of health insurance and their employees. Some employers have become proactive in managing health care costs by providing tools to encourage employees to directly manage their health and prevent disease. One example of such a tool is DASH for Health, an Internet-based nutrition and exercise behavior modification program. This program was offered as a free, opt-in benefit to US-based employees of the EMC Corporation.

Objective: The aim was to determine whether an employer-sponsored, Internet-based diet and exercise program has an effect on health care costs.

Methods: There were 15,237 total employees and spouses who were included in our analyses, of whom 1967 enrolled in the DASH for Health program (DASH participants). Using a retrospective, quasi-experimental design, study year health care costs among DASH participants and non-participants were compared, controlling for baseline year costs, risk, and demographic variables. The relationship between how often a subject visited the DASH website and health care costs also was examined. These relationships were examined among all study subjects and among a subgroup of 735 subjects with cardiovascular conditions (diabetes, hypertension, hyperlipidemia). Multiple linear regression analysis examined the relationship of program use to health care costs, comparing study year costs among DASH participants and non-participants and then examining the effects of increased website use on health care costs. Analyses were repeated among the cardiovascular condition subgroups.

Results: Overall, program use was not associated with changes in health care costs. However, among the cardiovascular risk study subjects, health care costs were US\$827 lower, on average, during the study year ($P = .05$; $t_{729} = 1.95$). Among 1028 program users, increased website use was significantly associated with lower health care costs among those who visited the website at least nine times during the study year (US\$14 decrease per visit; $P = .04$; $t_{1022} = 2.05$), with annual savings highest among 80 program users with targeted conditions (US\$55 decrease per visit; $P < .001$; $t_{74} = 2.71$).

Conclusions: An employer-sponsored, Internet-based diet and exercise program shows promise as a low-cost benefit that contributes to lower health care costs among persons at higher risk for above-average health care costs and utilization.

(*J Med Internet Res* 2009;11(4):e43) doi:[10.2196/jmir.1263](https://doi.org/10.2196/jmir.1263)

KEYWORDS

Employer health costs; disease management; health promotion; wellness programs; costs and cost analysis

Introduction

Health insurance premiums have risen faster than inflation for the past 10 years, placing an increasing burden on employer-sponsors of health insurance and their employees [1,2]. Some employers have become proactive in managing health costs, providing tools that encourage employees to directly manage their health and prevent disease [3]. Examples include smoking cessation and stress management programs, gym and health club memberships, and formal disease management programs, many of which are popular with employees and improve employee satisfaction. But there is very little evidence that any of these initiatives actually reduce health care costs [4].

Recent reviews of employer health promotion programs show some success in improving employee health and productivity, but show mixed results as to whether or not these programs have an impact on health care costs [5]. Employers commonly offer nutrition education programs [6,7], but there is little evidence that such programs alter eating behaviors or change health care costs. Yet rapidly rising rates of overweight and obesity can contribute to a number of high-cost chronic diseases (eg, diabetes, hypertension, heart disease), increasing the likelihood that health care costs for these already highly prevalent and expensive conditions will increase dramatically in the future.

To address the issue of poor nutrition and the diseases associated with it, we designed an Internet-based nutrition and exercise behavior modification program called DASH for Health. Our program was based on the National Heart Lung and Blood Institute (NHLBI) DASH diet, which was originally developed

to lower blood pressure and which has been demonstrated in randomized controlled trials to lower blood pressure and cholesterol levels and heighten insulin sensitivity [8-12]. We developed the Web-based DASH for Health program in collaboration with EMC Corporation, a Massachusetts-based global information infrastructure company. The program was offered as a free employee benefit to US-based EMC employees and their family members, who could opt-in to the program and were free to use the program however intensely they chose. Employees and their spouses were eligible to enroll in the program at the beginning of the calendar year in which it was offered. During the first year this program was offered at EMC Corporation (the same year that the health costs reported in this paper were collected), enrollees in the program were found to have significantly lost weight, lowered their blood pressure, and improved their healthy eating habits [11]. Examples of articles provided on the DASH for Health website are shown in [Figure 1](#) and [Figure 2](#).

To determine whether an Internet-based behavior modification program like DASH for Health has any effect on health care costs, we analyzed the costs for EMC employees and their spouses during the 12 months preceding the initial launch of the DASH for Health program and during the 12 months following the launch. The baseline year was the 12 months immediately preceding the initial launch of the program, and the study year was the 12 months immediately following the launch. We compared health care costs of those who participated in the DASH for Health program with those of nonparticipants. We analyzed costs for all study subjects and then performed a more focused analysis on employees and spouses with medical conditions (hypertension, hyperlipidemia, and/or diabetes) targeted by the DASH program.

Figure 1. Example of article provided on the DASH for Health website



DASH for Health™



EMC²
where information lives

Your Personal Online Program
To Improve Nutrition, Exercise And Health



HOME
YOUR STUFF
ARCHIVES
DASH STORE
TIP OF THE WEEK

HEALTH CALCULATORS

FAQ'S

SEND US A QUESTION

SEND US A RECIPE

ABOUT THE PROGRAM

LINKS TO DASH

ABOUT US

RECIPES

HEALTH LINK

LOG OUT

SEARCH

ENTER YOUR FOOD

Date:

Today's Weight:

Today's Blood Pressure: /

SUBMIT

CREAM OR MILK IN YOUR COFFEE?

What you add to your cup of coffee can make a big difference in the calories and fat you consume on a daily, monthly, and even yearly basis. It all adds up! National chains are starting to recognize that, too, and are offering lower-fat options for their customers. Starbucks, for example, now offers 2% milk, a much lighter version that still satisfies those customers that want a creamy cup.

Read on to learn more about how the way you dress your java and why it counts:

Think of it in terms of butter. Did you know that when you add cream, light cream, or even half and half to your coffee, you're effectively adding butter? That's right, the fat in cream is the same as the fat in butter. These coffee "creamers" have much more fat than even whole milk, 2% milk, and milk with less fat. But, you say, how much different can it make?? A LOT.

If you add cream or half-and-half to your coffee, most people add around 4 ounces to each "large" cup (about 16 ounces; equivalent to a Grande at Starbucks or a Medium at Dunkin Donuts). Check the table below to see how much BUTTER and how many calories you are adding EACH MONTH if you drink just one large coffee with milk or cream per day.



Light Cream in Your Coffee adds 7.6 Sticks of Butter Every Month

| 4 oz. per day serving | Sticks of butter per month | Calories per month |
|-----------------------|----------------------------|--------------------|
| Light Cream | 7.6 sticks | 7,100 |
| Half and Half | 4.6 sticks | 4,700 |
| Whole Milk | 1.3 sticks | 2,400 |
| 2% milk | 0.4 sticks | 1,830 |

Figure 2. Example of article provided on the DASH for Health website

The screenshot shows the DASH for Health website interface. At the top, there is a logo for 'DASH for Health' and 'EMC² Your Personal Online Program To Improve Nutrition, Exercise And Health'. Below the logo is a navigation menu with links for HOME, YOUR STUFF, ARCHIVES, DASH STORE, and TIP OF THE WEEK. On the left side, there is a vertical menu with links for HEALTH CALCULATORS, FAQ'S, SEND US A QUESTION, SEND US A RECIPE, ABOUT THE PROGRAM, LINKS TO DASH, ABOUT US, RECIPES, HEALTH LINK, and LOG OUT. Below this menu is a search bar and a form to 'ENTER YOUR FOOD' with fields for Date, Today's Weight, and Today's Blood Pressure, and a SUBMIT button. The main content area features an article titled 'EAT DASH, STAY SMART' with a sub-header 'The DASH eating plan grabbed the health headlines again last week when a team of researchers presenting data at the International Conference on Alzheimer's Disease in Switzerland announced their finding that the DASH Diet slowed the rate of mental decline.' The article text describes a study of 4000 people over 65 years old, published in the Cache County study of Memory Health and Aging. To the right of the article is an image of a human head profile with several colorful gears (blue, yellow, green, red) inside, symbolizing cognitive function. At the bottom of the article, there are links for CONTACT US, PRIVACY POLICY, and DISCLAIMER, and a footer with contact information for EMC Corporation.

Methods

Study Setting

The DASH for Health website was developed by a team at Boston University School of Medicine (BUSM). The website provides tools for enrollees to record and follow their body weight, blood pressure, eating habits, and exercise habits as well as providing a variety of healthy recipes based on the DASH diet. These tools include easy-to-use entry fields for entering weight, blood pressure, 24-hour food recall, and exercise habits. Although enrollees were encouraged to use these data “progress reports” as a means of encouraging their adherence to the program, there were no requirements or set expectations of how often enrollees would enter their own data. The website also provides two new articles each week on improving nutritional habits (based on the DASH diet) or healthy exercise. Everyone who enrolls in the program gets a reminder email each time a new article is posted on the website. The website is hosted at SignalZ Corporation (Montpelier, VT, USA). Self-reported data on demographics (age, gender, and employee/dependent status) and website visits during each year were collected by SignalZ and transferred to the research team

at BUSM. Because enrollees had to enter the site using their username and password, standard website monitoring software was used to track how many times a particular enrollee visited the site. We define a “visit” as a unique instance of an enrollee logging on to the site.

When the program was about to launch, EMC Corporation informed employees that DASH for Health would be available at no cost to them and their adult household members. EMC had no other role in the program. EMC management had no access to the identities of the subjects or to their medical data. Once the program began, the employer took no role in encouraging website usage. The announcing emails from EMC leadership clearly stated that the employer would not know any enrollee’s individual data or the identities of people who did and did not enroll.

EMC’s clinical data warehouse, Ingenix, provided demographic and medical and pharmacy cost data for employees and their adult dependents. These data included date of birth, gender, and employee status. D2Hawkeye, a medical analytics firm specializing in data warehousing and health claims analysis, provided data cleansing services and summarized study subjects’

demographic information, condition specific diagnoses, and total health care costs. D2Hawkeye calculated a risk adjustment factor for each study subject, using a proprietary methodology that combines diagnostic, procedural, and pharmacy information in health claims data with pattern recognition and times series analysis. The risk index provides a single representation of an individual's disease burden and has been shown to correctly distinguish high-cost and low-cost persons 77% of the time. Additionally, the likelihood that an individual will be in a high-cost group increases as the risk index value increases. We used an extension of the risk index, the adjusted risk index (ARI), which incorporates assessments of gaps in care (medical or pharmaceutical) for various conditions [13].

We used individual subject-level linkages to merge data on DASH for Health participants' website visit frequency with health care cost data and other variables. A subset of study subjects was identified with evidence of diabetes, hypertension, or hyperlipidemia, conditions that could be expected to benefit from the improved dietary behaviors that are encouraged by the DASH for Health program. This subset is referred to as the cardiovascular (CV) risk group. Data were managed and analyzed using SAS, version 9.1.3 (SAS Institute Inc, Cary, NC, USA). Approval for this study was obtained from the Boston University Institutional Review Board.

Study subjects with missing data for key fields such as age, employee status, or gender were excluded, as were DASH participants with no matching entries in the health care cost files. These enrollees were either not covered by the EMC insurance plan or were not affiliated with EMC at the end of the study period. The study was limited to study subjects with health care costs in both study years to ensure that those with bad claims data reported as negative or zero amounts were excluded from the analysis and to increase the likelihood that the analysis was restricted to those with primary insurance coverage through EMC. Information on job classification was not included since baseline year health care costs and risk assessment are proximal for our analysis.

The total number of EMC employees and their spouses was 29,675, of which approximately 75% ($n = 22,354$) were reported to have received health coverage through EMC and had health care costs in the baseline or study year. Sixty-eight percent ($n = 15,237$) of these subjects had costs in both years. Additionally, 3797 EMC employees or spouses enrolled in the DASH for Health program and used the DASH program website at least once in the study year (DASH participants). Seventy-three percent ($n = 2756$) of these DASH participants were found in the health claims summary file, and 71% ($n = 1967$) of these 2756 had health care costs in both years. The final study sample ($N = 15,237$, DASH = 1967, non-DASH = 13,270) also reflected the restrictions described above.

Analysis

In univariate analyses to describe our sample, we generated means and standard deviations for continuous variables and counts with percentages for categorical variables. We used bivariate analyses to examine the relationships among demographic, annual health care cost, and website usage data for 15,237 study subjects over the 2-year baseline and study

year period. In these analyses, we employed chi-square tests for analyses of categorical data and two-sample *t* tests for comparisons of continuous by categorical variables. Multiple linear regression analysis examined the relationship of DASH program enrollment with study year health care costs, controlling for other salient factors. To control for potential bias of self-selection of the DASH participants, we used a logistic regression model to construct a propensity score that represented the likelihood of participation in the DASH program. Studies that use observational data, as ours does, are subject to bias because the "treatment" and "control groups" are not randomly assigned and may differ in ways that affect the outcome of interest. A propensity score, which represents the conditional probability of receiving a given treatment, given a vector of measured covariates, is frequently used in such studies to adjust for differences in the observed characteristics between the treatment and the control groups. Our use of logistic regression is one of several recommended approaches for determining the propensity score [14].

All available study variables were included in the propensity score model (baseline year risk index and ARI, baseline year costs, age, gender, and employee status). We then used a linear regression model to compare study year costs among DASH and non-DASH participants, controlling for age, gender, ARI, baseline year costs, and likelihood of DASH participation (propensity score). In this model, study year total health care costs was the dependent (outcome) variable; independent (predictor) variables were DASH participation, age, gender, employee status, baseline year ARI, and baseline year total health care costs. Next, because stronger effects of an intervention are often observed in the most compliant or frequent participants, we examined whether more frequent use of the website than is typical is more strongly associated with reduced costs among DASH participants than across the full range of website use overall. For this, we used a similar linear regression analysis to evaluate the relationship of website usage intensity at or above the median to health care costs in the study year. Study year total health care costs was the dependent variable; independent variables were number of website visits, age, gender, employee status, baseline year ARI, and baseline year total health care costs. We repeated these analyses using the 735 DASH CV risk group and non-DASH CV risk group study subjects who showed evidence in both study years of diabetes, hyperlipidemia, or hypertension.

To address the skewed distribution of the cost data, baseline and study year costs in all analyses were top-coded at US\$25,000, which represents the 99th percentile of annual health care costs in our study population. Website usage was top-coded at 75 website visits to address skewness. The analyses were repeated using trimming to remove high-cost outliers, using the entire study population and using top-coding at other thresholds. As top-coding is recommended for reducing the effects of high-cost outliers on model results while retaining useable observations, we report results of those analyses here. Our choice of US\$25,000, or the 99th percentile, is also consistent with others [15-17].

As part of model development, we added interaction terms as covariates and conducted additional analyses where we limited

study subjects to employees, males, and male employees, reflecting the larger number of employees and males in the study group. To account for nonlinear effects of DASH participation and website use intensity, we grouped study year costs into quartiles; we also grouped website visits and study year costs into quartiles. There were no differences suggesting that the interaction terms, study population restrictions or groupings should be included in the final model as none of these modifications affected the results. We repeated the analyses we report here using square root and log transformations for all cost variables because of their good performance with heteroscedastic health care cost data [18]. Our results based on the transformed data were similar to those observed using the

untransformed data. Thus, to allow for ease of interpretation, we present results here for the untransformed data.

Results

All Study Subjects

Demographic, ARI, and baseline medical and pharmacy cost information were gathered for all study subjects (N = 15,237) and for the subgroup of subjects with CV risk conditions (N = 735). These measures were also examined for DASH (N = 1967) and non-DASH (N = 13,270) participants overall and in the CV risk group (DASH: N = 134; non-DASH: N = 601). These results are shown in Table 1.

Table 1. Baseline characteristics of DASH and non-DASH participants

| | All Study Subjects | | | CV Risk Subgroup ^a | | |
|-----------------------------------|-----------------------|--------------------------------------|--|-------------------------------|--------------------------------|---|
| | Total (N = 15,237) | DASH Partici- pants (N = 1967) | Non-DASH Partici- pants (N = 13,270) | Total (N = 735) | DASH Participants (N = 134) | Non-DASH Partici- pants (N = 601) |
| Cost baseline year (US\$) | | | | | | |
| mean (SD) | 2684 (7164) | 2181 (4351) | 2758 (7489) | 5663 (10,089) | 4239 (6335) | 5980 (10,727) |
| P25/P50/P75 ^b | 327/934/2612 | 345/933/2224 | 324/935/2563 | 1401/2772/5783 | 1002/2028/4527 | 1490/2849/6020 |
| Cost DASH year (US\$) | | | | | | |
| mean (SD) | 2814 (7835) | 2413 (4315) | 2879 (8228) | 5929 (13,611) | 3425 (3667) | 6487 (14,897) |
| P25/P50/P75 | 358/1006/2621 | 442/1145/2700 | 347/981/2607 | 1394/2681/5755 | 1146/2318/4222 | 1467/2848/6152 |
| Baseline year ARI | | | | | | |
| mean (SD) | 3.70 (6.80) | 3.52 (6.04) | 3.72 (6.91) | 12.14 (13.70) | 10.10 (9.73) | 12.59 (14.41) |
| P25/P50/P75 | 1/1/3 | 1/1/3 | 1/1/3 | 3/7/16 | 2/6/14 | 3/8/17 |
| Age (years) | | | | | | |
| mean (SD) | 40.2 (9.2) | 40.7 (9.1) | 40.1 (9.2) | 47.5 (8.54) | 46.1 (8.28) | 47.8 (8.57) |
| P25/P50/P75 | 33/40/46 | 34/41/47 | 33/40/46 | 42/49/54 | 41/47/52 | 42/49/54 |
| Gender, % (no.) | | | | | | |
| Male | 46 (7041) | 56 (1116) | 45 (5925) | 65 (476) | 73 (98) | 63 (378) |
| Female | 54 (8196) | 44 (851) | 55 (7345) | 35 (259) | 27 (36) | 37 (223) |
| Enrollment status, % (no.) | | | | | | |
| Employee | 55 (8384) | 84 (1659) | 51 (6725) | 64 (469) | 85 (114) | 59 (355) |
| Spouse | 45 (6853) | 16 (308) | 45 (6853) | 36 (266) | 15 (20) | 41 (246) |
| Website visits | | | | | | |
| mean (SD) | N/A ^c | 12.0 (17.0) | N/A | N/A | 16.9 (26.3) | N/A |
| P25/P50/P75 | N/A | 3/9/12 | N/A | N/A | 3/9/17 | N/A |

^a CV risk group subjects show evidence of hyperlipidemia, hypertension, and/or diabetes in both years.

^b P25/P50/P75 equals 25th, 50th, and 75th percentiles.

^cNot available.

Among the 15,237 study subjects, 55% (n = 8384) were employees, while 45% (n = 6853) were spouses. The overall study sample was 46% male (n = 7041), with an average age of 40.2 years. Average total baseline year health care costs were US\$2684. A slightly higher proportion of the DASH participants was male, compared to the non-DASH participants (56% vs

45%). The average age of the DASH participants was 40.7 years, slightly higher than the non-DASH participants (40.1 years). DASH participants were mostly employees (84%; n = 1659). DASH participants had a mean ARI of 3.52, which was slightly lower than the mean ARI for the non-DASH participants (3.72). Average total costs among DASH participants in the baseline

year were lower than among the non-DASH participants (US\$2181 vs US\$2758).

Table 1 also shows demographic, ARI, and baseline year health care costs for the DASH and non-DASH CV risk groups. This subgroup was 65% male (n = 476) and 64% employees (n =

469), with an average age of 47.5 years, older than the overall study sample. The mean baseline year ARI of 12.14 (DASH: 10.10, non-DASH: 12.59) was also higher than in the general study sample, as were mean total baseline health care costs (overall: US\$5663; DASH participants: US\$4239, non-DASH: US\$5980).

Table 2. Predictors of costs in DASH year: DASH vs non-DASH (overall and CV risk group)^a

| | All Study Subjects (N = 15,237) | | CV Risk Group (n = 735) | |
|----------------------------------|---|---------------------|---|---------------------|
| | Difference in Mean Study Year Cost ^b (SE) | P (t ^c) | Difference in Mean Study Year Cost ^b (SE) | P (t ^c) |
| DASH use vs non-use ^d | \$85.14 (\$90.83) | .35 (0.94) | -\$826.95 (\$424.81) | .05 (1.95) |
| Age | \$8.18 (\$3.31) | .01 (2.47) | \$50.40 (\$25.85) | .05 (1.95) |
| Male vs female | -\$453.05 (\$156.98) | .004 (2.89) | \$88.90 (\$645.85) | .89 (0.14) |
| Employee vs non-employee | -\$136.45 (\$370.54) | .72 (0.37) | -\$2862.85 (\$1564.49) | .07 (1.83) |
| Baseline year ARI ^e | \$123.46 (\$5.92) | < .001 (20.85) | \$133.69 (\$17.61) | < .001 (7.59) |
| Baseline year cost ^f | \$0.29 (\$0.01) | < .001 (29.00) | \$0.34 (\$0.04) | < .001 (8.50) |

^a Baseline and study year costs top-coded at US\$25,000; study year website visits top-coded at 75; probability of DASH participation included as model covariate (not shown).

^b For age, ARI, baseline year cost: difference in mean study year costs per unit difference; unit is one year (age), one integer (ARI), one US dollar (baseline year cost).

^c Degrees of freedom = n - 6.

^d DASH participants' health care costs were, on average, US\$85 higher than those of nonparticipants, although this result was not statistically significant.

^e Higher baseline year ARI increases were associated with higher study year costs. On average, study year costs increased US\$123 with each additional unit increase in the baseline year ARI. A unit refers to an integer; as an example, an ARI of 10 is one unit greater than an ARI of 9.

^f Higher baseline year health care costs were associated with higher study year costs. On average, study year costs were US\$0.29 higher for each additional dollar in baseline year cost.

The results of the linear regression analysis of study year costs for DASH vs non-DASH in the full sample are shown in Table 2. Among all study subjects, DASH participation was associated with increased health care costs, although this result was not statistically significant (difference in mean costs, DASH vs non-DASH = US\$85.14; $P = .35$). Model covariates associated with significantly higher study year costs were older age ($P = .01$), being female ($P = .004$), higher baseline ARI ($P < .001$), and higher baseline year costs ($P < .001$).

Results of the linear regression analysis of study year costs for DASH and non-DASH CV risk group study subjects are also shown in Table 2. DASH CV risk group members' study year health care costs were US\$827 lower, on average, than those of the non-DASH CV risk group members ($P = .05$). Higher baseline year ARI was significantly associated with higher study year costs, with each additional unit of risk associated with an increase, on average, of US\$134 ($P < .001$). Each additional dollar in baseline year costs was associated with an additional US\$0.34, on average, in study year costs ($P < .001$).

Table 3. Cost in DASH year as a function of intensity of website use, adjusting for covariates^a

| | All DASH Participants (n = 1967) | | CV Risk Group (n = 134) | |
|---|---|---------------------|---|---------------------|
| | Difference in Mean Study Year Cost ^b (SE) | P (t ^c) | Difference in Mean Study Year Cost ^b (SE) | P (t ^c) |
| Change in costs per website visit ^d | -\$6.66 (\$5.34) | .21 (1.25) | -\$28.45 (\$14.61) | .054 (1.95) |
| Age ^e | \$34.07(\$8.05) | < .001 (4.23) | \$17.50 (\$31.57) | .58 (0.55) |
| Male vs female | -\$458.43 (\$161.07) | .005 (2.85) | -\$1121.62 (\$686.75) | .11 (1.63) |
| Employee vs non-employee | \$66.82 (\$215.01) | .76 (0.31) | -\$39.67 (\$861.64) | .96 (0.05) |
| Baseline year ARI ^f | \$60.24 (\$16.74) | < .001 (3.60) | \$54.48 (\$36.69) | .14 (1.48) |
| Baseline year cost ^g | \$0.30 (\$0.03) | < .001 (10.00) | \$0.34 (\$0.07) | < .001 (4.86) |
| | DASH Participants Website Use at or Above Median ^h (n = 1028) | | CV Risk Group Website Use at or Above Median ^h (n = 80) | |
| | Difference in Mean Study Year Cost ^b (SE) | P (t ^d) | Difference in Mean Study Year Cost ^b (SE) | P (t ^d) |
| Change in costs per website visit | -\$14.26 (\$6.97) | .04 (2.05) | -\$54.61 (\$20.16) | .01 (2.71) |
| Age | \$49.22 (\$11.87) | < .001 (4.15) | \$18.69 (\$47.47) | .70 (0.39) |
| Male vs female | -\$444.33 (\$230.67) | .05 (1.93) | -\$1855.55 (\$940.40) | .05 (1.97) |
| Employee vs non-employee | \$141.08 (\$301.12) | .64 (0.47) | \$757.00 (\$1286.59) | .56 (0.59) |
| Baseline year ARI | \$109.73 (\$26.49) | < .001 (4.14) | \$98.59 (\$55.97) | .08 (1.76) |
| Baseline year cost | \$0.24 (\$0.04) | < .001 (6.00) | \$0.25 (\$0.12) | .04 (2.08) |

^a Baseline and study year costs top-coded at US\$25,000; study year website visits top-coded at 75.

^b For number of website visits, age, ARI, baseline year costs: difference in mean study year costs per unit difference; unit is one year (age), one integer (ARI), one US dollar (baseline year cost).

^c Degrees of freedom = n - 6.

^d Among all DASH participants, each additional website visit was associated, on average, with a US\$6.66 decrease in study year health care cost. This result was not statistically significant. Among CV risk group DASH participants, each additional website visit was associated with a US\$28 decrease in study year cost; this result was not statistically significant at the $P < .05$ level. Among DASH participants who visited the website at least the median number of times during the study year (nine visits), each additional visit was associated with a US\$14 study year cost decrease. Among CV risk group DASH participants who visited the website at least the median number of times for the CV risk group (also nine visits), each additional visit was associated with a US\$55 decrease in study year cost.

^e Among all DASH participants, each additional year of age was associated with US\$34, on average, higher study year health care cost. Among DASH participants who visited the website nine or more times during the study year, each additional year of age was associated with US\$49 higher study year health care cost. Among CV risk group DASH participants, the relationship of age to study year health care costs was not statistically significant at the $P < .05$ level.

^f Among all DASH participants, each additional increment in baseline year ARI was associated with US\$60 higher study year health care cost.

^g Each additional dollar in baseline year costs was associated with increased study year health care costs as follows: US\$0.30 among all DASH participants; US\$0.34 among CV risk group DASH participants; US\$0.24 among DASH participants who visited the website at least nine times; US\$0.25 among CV risk group DASH participants who visited the website at least nine times.

^h Median website usage for all DASH and CV risk group DASH participants: nine visits.

Results of the analyses of intensity of website usage and study year costs are shown in Table 3. Among 1967 DASH participants, each additional website visit was associated with almost a US\$7 decrease in study year health care costs, but this result was not statistically significant ($P = .21$). Among

participants whose website usage was at or above the nine-visit study year median (n = 1028), each additional website visit was associated with a US\$14 decrease in health care costs on average ($P = .04$).


Figure 3. CV risk group change in unadjusted total costs from baseline to DASH study year in DASH participants vs nonparticipants

Figure 3 compares the change in crude study year costs, relative to baseline year costs, among DASH participants versus nonparticipants in the CV risk group. Among DASH CV risk group participants, study year health care costs were US\$814 lower than baseline year costs. Among DASH nonparticipants, study year costs were US\$507 higher than baseline year costs. These results are not adjusted to control for age, gender, employee status, or baseline ARI.

Results of the analysis of website use intensity in the DASH CV risk group are also shown in Table 3. Among the 134 DASH CV risk group members, each additional website visit was associated with a decrease of approximately US\$28 in health care costs ($P = .05$). Among those who used the website at least the median number of nine times during the year ($n = 80$), each additional visit was associated with a US\$55 decrease in study year costs ($P = .01$).

Discussion

We did not find DASH participation to be associated with lower health care costs in the overall study sample. With an average age of 40 years, an average ARI of 3.7, and minimal health services utilization, this study sample is relatively young and healthy. The significance of higher baseline year risk and costs, female gender, and increasing age as predictors of study year costs is consistent with a relatively young, commercially insured population. Improvements in diet and exercise habits would not be expected to address the highest expenditures in this group—pregnancy and childbirth, depression, and back pain.

We found suggestive evidence that DASH program use was associated with decreased health care spending among study subjects with CV risk. Within the DASH CV risk group, participants' study year health care costs were US\$827 lower than for the DASH nonparticipants. Although this result only achieved a significance level of $P = .05$, the reduction in mean costs from DASH participation (US\$827), which was nearly twice the standard deviation (US\$425), indicates that the effect size is quite large; the model R-square, 39%, is also noteworthy. Improvements in diet and exercise would be expected to address health care expenditures among study subjects with hyperlipidemia, hypertension, or diabetes. This subgroup is older (average age 47 years) and less healthy, with a baseline year ARI more than three times that of the general study sample.

We also found evidence suggestive of a dose-response relationship. Among DASH participants who visited the website at least nine times during the study year, each additional visit was associated with lower study year costs overall ($P = .04$) and in the DASH CV risk group ($P = .01$). Evidence of this dose-response was strongest among the DASH participants in the CV risk group, where each additional website visit was associated with a US\$55 decrease in study year health care costs ($P = .01$).

The DASH CV risk group participants' baseline ARI (10.10) suggests that they may be less sick than the non-DASH CV risk group (baseline ARI = 12.59). However, the DASH CV risk group, with conditions targeted by the DASH program, was the only subgroup that showed a decrease in health spending. Health care costs among DASH enrollees decreased 24% between the baseline (US\$4239) and study years (US\$3425) in the DASH

CV risk group. By contrast, health spending increased US\$507 (8%) in the non-DASH CV risk group and approximately US\$130 per person (4.8%) across the general study sample.

We also found evidence of selective enrollment in the DASH program by study subjects with hypertension, hyperlipidemia, and diabetes. Despite being offered as a benefit to all employees and their dependents, participation in DASH for Health was proportionally higher (18%) among individuals with CV risk conditions (ie, those who might benefit most from nutrition improvement) than it was among individuals without those conditions, of whom 13% signed up for the DASH program ($P < .001$). For employers interested in offering benefit programs equally to all employees rather than targeting a selected subset of the employee population, this finding provides evidence that the DASH for Health program benefits the subset of the employee population whose health risks are of concern and whose health status is targeted by the program.

Our results expand on the already-published reports that Internet-based programs can have positive effects on clinical parameters such as weight and blood pressure [11,19-22] and contribute to our understanding of the effects of nutrition, weight management, and exercise programs on health care costs in targeted populations [23-26], including workplace populations [3,27-32]. Several recent studies indicate that workplace-sponsored, Web-based programs can lead to these improvements in clinical parameters [11,33-36]. Our focus on the effects of an employer-sponsored, Web-based diet and exercise program on health care costs also expands our understanding of the effects of employer benefits that encourage employees to better manage their health status and contain health care costs. Hsu et al [37] observed significant increases in nonpharmaceutical health care expenses over a short time period among persons with chronic conditions who reduced their prescription medication use in response to increased cost sharing in benefit design. By contrast, regular use of the DASH for Health program may encourage health behavior changes that result in cost savings among persons with chronic conditions in a similarly short time frame. The benefits of DASH participation among healthier, younger enrollees may be evident over a longer time period than the year evaluated for this study. Employers, particularly those who are self-insured, may be interested in both short- and long-term employee costs and health status and choose to invest in health status improvements that will show benefits over the long term [38-40].

Limitations

A number of study limitations should be noted. Our analysis used observational data and is vulnerable to selection effects, which represent the largest threat to validity in observational studies. Our use of a control group and a pre-test, coupled with our use of a propensity score, are standard mechanisms for addressing selection bias in quasi-experimental study designs. However, our model would not control for the possibility that study subjects who were motivated to manage their CV risk conditions were more likely to enroll in DASH for Health and would visit the website more often. However, it is unlikely that our finding that even moderate use of the DASH for Health program is associated with lower health care costs is attributable

to motivation and not to participation in the program. This finding is consistent with clinically oriented studies of the DASH diet program indicating that the DASH diet is associated with improved blood pressure, lower cholesterol levels, and increased insulin sensitivity. Such clinical improvements would be expected to be associated with lower health care costs.

Several other limitations should be noted. First, the study sample was limited to persons with health care costs in both years. The proportion of study subjects with costs in only one year (costs in baseline year only: 90%; costs in study year only: 79%) is consistent with the national Medical Expenditure and Panel Survey (MEPS) data indicating that 11% of commercially insured persons do not use health services in a given year. D2Hawkeye received and cleaned the source claims data and reported that the proportion of study subjects with zero costs was higher than expected. Because the data could not be sent back to the data warehouse for review, we chose to restrict the study to persons with costs in both years. In analyses with all study subjects, including those with no costs in either or both years, costs for all DASH participants were significantly lower than for non-DASH participants, and the cost savings were higher among DASH participants with CV risk conditions. (Results available on request.) Restricting study subjects to those with costs understates the effects of the DASH program on costs. Second, our analysis compares study subjects who visited the DASH website at least once with those who did not visit it at all. However, subjects who visited the website once during the study year would not be expected to benefit from the nutrition and exercise education that the DASH for Health website program offers. Our finding of a dose-response effect would be strengthened by the inclusion of subjects who signed up for the program but never visited the website. From the perspective of benefit design, this effect would be more useful for guiding planning or coaching efforts. Third, our propensity score is based only on available data and therefore may not completely address potential selection bias. It is possible that DASH participants with CV risk conditions would have had decreased health care costs without participation in DASH. However, the clinically observable effects of the DASH diet on CV risk conditions make this result less likely. Fourth, our results are based on all health care costs for all conditions. Costs for conditions targeted by the DASH diet cannot be distinguished from costs for medical conditions that are not likely to benefit from improved diet, such as pregnancy and childbirth and back pain. However, the dose-response effect among participants who use the DASH website more often and the increasing significance of additional website usage in the DASH CV risk group suggest that participation in the DASH program is related to decreases in health care costs among persons at higher risk for health care expenses. We performed additional analyses of the effects of DASH participation on study year costs for subjects with different baseline year risk levels. Based on the regression model, at higher levels of baseline year risk, DASH participants had significantly lower study year costs than nonparticipants. (Results available on request.) These results are of particular interest given that higher levels of baseline year risk are predictive of higher costs [13]. Finally, study year costs include health care costs incurred during the initial enrollment period. The effects of DASH for

Health program participation would not be expected to be evident during the initial launch of the program or during the first few months of the year. However, this limitation suggests that our study results understate the effects of DASH for Health program participation since they include this initial time frame.

Our focus on short-term cost savings does not examine whether these savings are offset by increases in other costs [40]. Although the DASH for Health program is relatively inexpensive, further study to evaluate its cost-effectiveness as a corporate benefit would be beneficial. Finally, the study sample was limited to employees at one technology firm and their dependents. Our study results may not be generalizable to persons without employer-based insurance or to adults who are

less comfortable using the Internet for information about diet, exercise, and health management. Research into the effectiveness of DASH for Health in other populations is warranted.

Conclusions

Use of an Internet-based program that targets changes in diet and exercise to reduce weight, cholesterol, and blood pressure shows evidence of reducing short-term health care costs among persons at high risk for health care expenditures with conditions targeted by the diet. Offering access to a website with diet and exercise information appears to have promise as a low-cost, employer-sponsored benefit that contributes to lower health care costs among persons at higher risk for above-average health costs and utilization.

Acknowledgments

We wish to thank Diana Lehman for her assistance in preparing this manuscript; Jack Mollen, Lauri Tenney, and Leslie Berger for their help in introducing the DASH for Health program at EMC Corporation; and Binod Bhattarai for claims data analysis guidance at D2 Hawkeye, Inc. The development of the DASH for Health™ program was supported by donations from several companies that produce or promote healthy food products (Cabot Cheese, California Table Grape Commission, ConAgra Foods, Dannon, Diamond Nuts of California, Florida Department of Citrus, Garelick Farms, General Mills, International Banana Association, Minute Maid, National Dairy Council, The Peanut Institute, Sunkist, and SunMaid) as well as donations from EMC Corporation and Stop and Shop supermarkets.

Conflicts of Interest

Dr. Moore is Chairman of E-Havior Change, LLC, which owns the copyright to the DASH for Health™ program.

References

1. ; Kaiser Family Foundation and Health Research & Educational Trust. Employer Health Benefits: 2008 Annual Survey. Menlo Park, CA: The Henry J. Kaiser Family Foundation; 2008. URL: <http://ehbs.kff.org/pdf/7790.pdf> [accessed 2009-05-05] [[WebCite Cache ID 5gXrWjDI3](#)]
2. Kaiser Family Foundation. 2009. Trends in Health Care Costs and Spending. URL: http://www.kff.org/insurance/upload/7692_02.pdf [accessed 2009-05-05] [[WebCite Cache ID 5gXrkt8a1](#)]
3. Okie S. The employer as health coach. *N Engl J Med* 2007 Oct 11;357(15):1465-1469. [Medline: [17928592](#)] [doi: [10.1056/NEJMp078152](#)]
4. Mattke S, Seid M, Ma S. Evidence for the effect of disease management: is \$1 billion a year a good investment? *Am J Manag Care* 2007 Dec;13(12):670-676 [[FREE Full text](#)] [Medline: [18069910](#)]
5. Goetzel RZ, Ozminkowski RJ. The health and cost benefits of work site health-promotion programs. *Annu Rev Public Health* 2008;29(1):303-323. [Medline: [18173386](#)] [doi: [10.1146/annurev.publhealth.29.020907.090930](#)]
6. Buller DB, Morrill C, Taren D, Aickin M, Sennott-Miller L, Buller MK, et al. Randomized trial testing the effect of peer education at increasing fruit and vegetable intake. *J Natl Cancer Inst* 1999 Sep 1;91(17):1491-1500 [[FREE Full text](#)] [Medline: [10469751](#)] [doi: [10.1093/jnci/91.17.1491](#)]
7. Byers T, Mullis R, Anderson J, Dusenbury L, Gorsky R, Kimber C, et al. The costs and effects of a nutritional education program following work-site cholesterol screening. *Am J Public Health* 1995 May;85(5):650-655 [[FREE Full text](#)] [Medline: [7733424](#)] [doi: [10.2105/AJPH.85.5.650](#)]
8. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med* 1997 Apr 17;336(16):1117-1124 [[FREE Full text](#)] [Medline: [9099655](#)] [doi: [10.1056/NEJM199704173361601](#)]
9. Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, et al; DASH-Sodium Collaborative Research Group. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med* 2001 Jan 4;344(1):3-10 [[FREE Full text](#)] [Medline: [11136953](#)]
10. Lien LF, Brown AJ, Ard JD, Loria C, Erlinger TP, Feldstein AC, et al. Effects of PREMIER lifestyle modifications on participants with and without the metabolic syndrome. *Hypertension* 2007 Oct;50(4):609-616 [[FREE Full text](#)] [Medline: [17698724](#)] [doi: [10.1161/HYPERTENSIONAHA.107.089458](#)]
11. Moore TJ, Alsabeeh N, Apovian CM, Murphy MC, Coffman GA, Cullum-Dugan D, et al. Weight, blood pressure, and dietary benefits after 12 months of a Web-based Nutrition Education Program (DASH for health): longitudinal observational study. *J Med Internet Res* 2008;10(4):e52 [[FREE Full text](#)] [Medline: [19073541](#)] [doi: [10.2196/jmir.1114](#)]

12. Levitan EB, Wolk A, Mittleman MA. Consistency with the DASH diet and incidence of heart failure. *Arch Intern Med* 2009 May 11;169(9):851-857. [Medline: [19433696](#)] [doi: [10.1001/archinternmed.2009.56](#)]
13. Gautam S, Singh S. 2Hawkeye, Inc.. 2008. Predicting Overall and Impactable Future Costs with the D2Hawkeye Risk Modelling System. URL: http://www.d2hawkeye.com/Web%20files/D2_Risk_Models_White_Paper.pdf [accessed 2009-05-05] [WebCite Cache ID 5gXsHXxAd]
14. D'Agostino RB. Propensity score methods for bias reduction in the comparison of a treatment to a non-randomized control group. *Stat Med* 1998 Oct 15;17(19):2265-2281. [Medline: [9802183](#)] [doi: [10.1002/\(SICI\)1097-0258\(19981015\)17:19<2265::AID-SIM918>3.0.CO;2-B](#)]
15. Zhao Y, Ash AS, Haughton J, McMillan B. Identifying Future High Cost Cases Through Predictive Modeling. *Dis Manage Health Outcomes* 2003;11(6):389-397. [doi: [10.2165/00115677-200311060-00005](#)]
16. Shwartz M, Ash A, Pekoz E. Risk adjustment and risk-adjusted provider profiles. *Int J Health Care Technol Manage* 2006;7(1/2):15-42. [doi: [10.1504/IJHTM.2006.007871](#)]
17. Yaisawarng S, Burgess JF. Performance-based budgeting in the public sector: an illustration from the VA health care system. *Health Econ* 2006 Mar;15(3):295-310. [Medline: [16331724](#)] [doi: [10.1002/hec.1060](#)]
18. Montez-Rath M, Christiansen CL, Ettner SL, Loveland S, Rosen AK. Performance of statistical models to predict mental health and substance abuse cost. *BMC Med Res Methodol* 2006;6(1):53 [FREE Full text] [Medline: [17067394](#)] [doi: [10.1186/1471-2288-6-53](#)]
19. Micco N, Gold B, Buzzell P, Leonard H, Pintauro S, Harvey-Berino J. Minimal in-person support as an adjunct to internet obesity treatment. *Ann Behav Med* 2007 Feb;33(1):49-56. [Medline: [17291170](#)] [doi: [10.1207/s15324796abm3301_6](#)]
20. Tate DF, Jackvony EH, Wing RR. Effects of Internet behavioral counseling on weight loss in adults at risk for type 2 diabetes: a randomized trial. *JAMA* 2003 Apr 9;289(14):1833-1836 [FREE Full text] [Medline: [12684363](#)] [doi: [10.1001/jama.289.14.1833](#)]
21. Tate DF, Jackvony EH, Wing RR. A randomized trial comparing human e-mail counseling, computer-automated tailored counseling, and no counseling in an Internet weight loss program. *Arch Intern Med* 2006;166(15):1620-1625 [FREE Full text] [Medline: [16908795](#)] [doi: [10.1001/archinte.166.15.1620](#)]
22. Williamson DA, Martin PD, White MA, Newton R, Walden H, York-Crowe E, et al. Efficacy of an internet-based behavioral weight loss program for overweight adolescent African-American girls. *Eat Weight Disord* 2005 Sep;10(3):193-203. [Medline: [16277142](#)]
23. Musich SA, Adams L, Edington DW. Effectiveness of health promotion programs in moderating medical costs in the USA. *Health Promot Int* 2000;15(1):5-15. [doi: [10.1093/heapro/15.1.5](#)]
24. Ozminkowski RJ, Goetzel RZ, Wang F, Gibson TB, Shechter D, Musich S, et al. The savings gained from participation in health promotion programs for Medicare beneficiaries. *J Occup Environ Med* 2006 Nov;48(11):1125-1132. [Medline: [17099448](#)] [doi: [10.1097/01.jom.0000240709.01860.8a](#)]
25. Ozminkowski RJ, Ling D, Goetzel RZ, Bruno JA, Rutter KR, Isaac F, et al. Long-term impact of Johnson & Johnson's Health & Wellness Program on health care utilization and expenditures. *J Occup Environ Med* 2002 Jan;44(1):21-29. [Medline: [11802462](#)] [doi: [10.1097/00043764-200201000-00005](#)]
26. Pelletier KR. A review and analysis of the clinical and cost-effectiveness studies of comprehensive health promotion and disease management programs at the worksite: update VI 2000-2004. *J Occup Environ Med* 2005 Oct;47(10):1051-1058. [Medline: [16217246](#)] [doi: [10.1097/01.jom.0000174303.85442.bf](#)]
27. Herman CW, Musich S, Lu C, Sill S, Young JM, Edington DW. Effectiveness of an incentive-based online physical activity intervention on employee health status. *J Occup Environ Med* 2006 Sep;48(9):889-895. [Medline: [16966955](#)] [doi: [10.1097/01.jom.0000232526.27103.71](#)]
28. McCarty CA, Scheuer D. Lessons learned from employee fitness programs at the Marshfield Clinic. *WMJ* 2005 Jul;104(5):61-65. [Medline: [16138518](#)]
29. Stein AD, Shakour SK, Zuidema RA. Financial incentives, participation in employer-sponsored health promotion, and changes in employee health and productivity: HealthPlus Health Quotient Program. *J Occup Environ Med* 2000 Dec;42(12):1148-1155. [Medline: [11125677](#)] [doi: [10.1097/00043764-200012000-00005](#)]
30. Voit S. Work-site health and fitness programs: Impact on the employee and employer. *Work* 2001;16(3):273-286. [Medline: [12441457](#)]
31. Young JM. Promoting health at the workplace: challenges of prevention, productivity, and program implementation. *N C Med J* 2006;67(6):417-424. [Medline: [17393702](#)]
32. Aldana SG. Financial impact of health promotion programs: a comprehensive review of the literature. *Am J Health Promot* 2001;15(5):296-320. [Medline: [11502012](#)]
33. Cook RF, Billings DW, Hersch RK, Back AS, Hendrickson A. A field test of a web-based workplace health promotion program to improve dietary practices, reduce stress, and increase physical activity: randomized controlled trial. *J Med Internet Res* 2007;9(2):e17 [FREE Full text] [Medline: [17581811](#)] [doi: [10.2196/jmir.9.2.e17](#)]
34. Norman GJ, Zabinski MF, Adams MA, Rosenberg DE, Yaroch AL, Atienza AA. A review of eHealth interventions for physical activity and dietary behavior change. *Am J Prev Med* 2007 Oct;33(4):336-345 [FREE Full text] [Medline: [17888860](#)] [doi: [10.1016/j.amepre.2007.05.007](#)]

35. Sternfeld B, Block C, Quesenberry CP, Block TJ, Husson G, Norris JC, et al. Improving diet and physical activity with ALIVE: a worksite randomized trial. *Am J Prev Med* 2009 Jun;36(6):475-483. [Medline: [19460655](#)] [doi: [10.1016/j.amepre.2009.01.036](#)]
36. Block G, Sternfeld B, Block CH, Block TJ, Norris J, Hopkins D, et al. Development of Alive! (A Lifestyle Intervention Via Email), and its effect on health-related quality of life, presenteeism, and other behavioral outcomes: randomized controlled trial. *J Med Internet Res* 2008;10(4):e43 [FREE Full text] [Medline: [19019818](#)] [doi: [10.2196/jmir.1112](#)]
37. Hsu J, Price M, Huang J, Brand R, Fung V, Hui R, et al. Unintended consequences of caps on Medicare drug benefits. *N Engl J Med* 2006 Jun 1;354(22):2349-2359 [FREE Full text] [Medline: [16738271](#)] [doi: [10.1056/NEJMsa054436](#)]
38. ; Kaiser Family Foundation and the National Governors Association Center for Best Practices. Elements of State Health Reform: Prevention and Wellness as Covered Benefits. Menlo Park, CA: Kaiser Family Foundation; 2007. URL: [url:http://www.kaisernetwork.org/health_cast/uploaded_files/060707_kff_nga_transcript.pdf](http://www.kaisernetwork.org/health_cast/uploaded_files/060707_kff_nga_transcript.pdf) [accessed 2009-05-05] [WebCite Cache ID 5gXsTEBrm]
39. Newhouse JP. Consumer-directed health plans and the RAND Health Insurance Experiment. *Health Aff (Millwood)* 2004;23(6):107-113. [Medline: [15584103](#)] [doi: [10.1377/hlthaff.23.6.107](#)]
40. Nicholson S, Pauly MV, Polsky D, Baase CM, Billotti GM, Ozminkowski RJ, et al. How to present the business case for healthcare quality to employers. *Appl Health Econ Health Policy* 2005;4(4):209-218. [Medline: [16466272](#)] [doi: [10.2165/00148365-200504040-00003](#)]

Abbreviations

ARI: adjusted risk index

BUSM: Boston University School of Medicine

CV: cardiovascular

DASH: Dietary Approaches to Stop Hypertension

Edited by G Eysenbach; submitted 15.05.09; peer-reviewed by G Block; comments to author 07.06.09; revised version received 21.08.09; accepted 27.08.09; published 23.10.09

Please cite as:

Sacks N, Cabral H, Kazis LE, Jarrett KM, Vetter D, Richmond R, Moore TJ

A Web-Based Nutrition Program Reduces Health Care Costs in Employees With Cardiac Risk Factors: Before and After Cost Analysis
J Med Internet Res 2009;11(4):e43

URL: <http://www.jmir.org/2009/4/e43/>

doi: [10.2196/jmir.1263](#)

PMID: [19861297](#)

© Naomi Sacks, Howard Cabral, Lewis E Kazis, Kelli M Jarrett, Delia Vetter, Russell Richmond, Thomas J Moore. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 23.10.2009. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.