
Review

Mapping mHealth Research: A Decade of Evolution

Maddalena Fiordelli, PhD; Nicola Diviani, PhD; Peter J Schulz, PhD

Institute of Communication and Health, Faculty of Communication Sciences, University of Lugano, Lugano, Switzerland

Corresponding Author:

Maddalena Fiordelli, PhD

Institute of Communication and Health

Faculty of Communication Sciences

University of Lugano

Via G. Buffi, 13

Lugano, 6900

Switzerland

Phone: 41 586664757

Fax: 41 586664647

Email: maddalena.fiordelli@usi.ch

Abstract

Background: For the last decade, mHealth has constantly expanded as a part of eHealth. Mobile applications for health have the potential to target heterogeneous audiences and address specific needs in different situations, with diverse outcomes, and to complement highly developed health care technologies. The market is rapidly evolving, making countless new mobile technologies potentially available to the health care system; however, systematic research on the impact of these technologies on health outcomes remains scarce.

Objective: To provide a comprehensive view of the field of mHealth research to date and to understand whether and how the new generation of smartphones has triggered research, since their introduction 5 years ago. Specifically, we focused on studies aiming to evaluate the impact of mobile phones on health, and we sought to identify the main areas of health care delivery where mobile technologies can have an impact.

Methods: A systematic literature review was conducted on the impact of mobile phones and smartphones in health care. Abstracts and articles were categorized using typologies that were partly adapted from existing literature and partly created inductively from publications included in the review.

Results: The final sample consisted of 117 articles published between 2002 and 2012. The majority of them were published in the second half of our observation period, with a clear upsurge between 2007 and 2008, when the number of articles almost doubled. The articles were published in 77 different journals, mostly from the field of medicine or technology and medicine. Although the range of health conditions addressed was very wide, a clear focus on chronic conditions was noted. The research methodology of these studies was mostly clinical trials and pilot studies, but new designs were introduced in the second half of our observation period. The size of the samples drawn to test mobile health applications also increased over time. The majority of the studies tested basic mobile phone features (eg, text messaging), while only a few assessed the impact of smartphone apps. Regarding the investigated outcomes, we observed a shift from assessment of the technology itself to assessment of its impact. The outcome measures used in the studies were mostly clinical, including both self-reported and objective measures.

Conclusions: Research interest in mHealth is growing, together with an increasing complexity in research designs and aim specifications, as well as a diversification of the impact areas. However, new opportunities offered by new mobile technologies do not seem to have been explored thus far. Mapping the evolution of the field allows a better understanding of its strengths and weaknesses and can inform future developments.

(*J Med Internet Res* 2013;15(5):e95) doi: [10.2196/jmir.2430](https://doi.org/10.2196/jmir.2430)

KEYWORDS

mHealth; systematic review; health outcomes

Introduction

In the last decade, mobile health (mHealth), the branch of eHealth broadly defined as “the use of mobile computing and communication technologies in health care and public health” [1], has been constantly expanding. Mobile applications for health can target heterogeneous audiences such as doctors, nurses, patients, or even healthy people [1]. Different features of mobile phones may address specific needs in different situations. Available literature suggests that the use of mobile phones serves a wide variety of purposes [2], such as smoking cessation, weight loss, diet and physical activity, treatment adherence, and disease management. The biggest advantages of using mobile devices, and in particular mobile phones, for health are that these devices are personal, intelligent, connected, and always with people [3,4]. Therefore, they can serve patients both in everyday life and during hospitalization or rehabilitation, as well as health care providers during emergency or routine visits. Current evidence suggests that the use of mobile technology can improve diagnosis and compliance with treatment guidelines, as well as patient information, and can increase administrative efficiency [5]. In particular, short message service (SMS) text messaging reminders have been shown to be a simple and efficient option for health services to use in order to improve service delivery, resulting in health benefits for the patients who receive them [6]. Mobile phone technologies have also been shown to be effective in smoking cessation, weight loss, physical activity, diabetes management, STD prevention and treatment, and hypertension [7].

The mobile phone market is constantly evolving. The first digital mobile phones appeared in the early 1990s, and since then, mobile technology has continued to be refined thanks to the development of new features and better networks. Current smartphones have been defined as “mobile telephones with computer features that may enable them to interact with computerized systems, send e-mails, and access the web” [8]. Over a third of US mobile phone users own a smartphone [3,9], and it is estimated that 67.6% of adults worldwide own a mobile phone [2,10], making it the most equitable communication technology [1]. It has been argued that mobile phones could be a solution to overcome the traditional digital divide derived from the introduction of the Internet because they provide new opportunities to reach underserved and previously unreachable parts of the population worldwide, especially in developing countries [2].

Mobile technology, with its diffusion and characteristics, holds a great potential for health care applications. However the use of mobile phones in health care delivery has not been fully explored, and the diverse outcomes of mHealth have barely been documented. Although some literature reviews cover one part or the other of the field [6,11,12], an overall picture is still missing, possibly due to the field’s constant evolution. A recent methodological review sought to map the domain of mobile phone health interventions [13], but it relied on describing the design of the interventions, with a clear focus on technology, rather than the outcomes. As the authors stated, their motivation lay in the fact that “effectiveness reviews can be best done at

the level of a particular pathology”, while they wanted to draw a more comprehensive taxonomy of the field.

The main objective of this paper, as stated in the title, is to map the field, but without omitting the outcome measures. This means that our intention is to investigate how the impact of mobile phones on health has been assessed in peer-reviewed scientific literature. In particular, we are interested in understanding the evolution over the past decade, how the interventions have been developed, the main health care delivery areas where the impact of mobile technologies has been assessed, the methodology and features used, and finally, the type of outcome measures and general impact of the intervention.

The second objective of this review is to understand, after the 5 years since the introduction of the new generation of smartphones (eg, the iPhone in 2007), whether and how these devices have triggered research. The appeal of these new devices resides in the fact that they include several computer-like built-in features (eg, the GPS or the accelerometer) allowing the monitoring of a whole series of behaviors. Additionally, new mobile operating systems allow users to customize their devices according to their needs, by downloading apps available for free or for a low price from a central store. Klasnja and Pratt named this kind of feature “native application” [13], which is a typical complex and sophisticated application that can be implemented on major smartphone platforms (iOS, Android, Symbian, BlackBerry, webOS, and Windows Phone). In 2012, smartphone users spent US \$8 billion for paid apps in the top 5 app platforms, and the European mobile app market size reached €1.68 billion [14]. Therefore, iPhones and similar devices are potentially very interesting for application in health care—they already integrate most of the features that researchers previously had to add to traditional mobile phones in order to use them for health-related purposes and monitoring [15,16].

Methods

The objective of this study was to provide a comprehensive picture of how the impact of mHealth was assessed in the scientific literature in its first decade of existence. For this purpose, a systematic literature review was conducted in which relevant studies were categorized in a two-step process. The first step included the review of the titles and abstracts of all publications that were identified as potentially relevant, with the goal of assessing whether they might meet the inclusion criteria for the systematic review. Selected abstracts were categorized at this stage using general typologies partly adapted from existing literature [1,2,12] and partly created inductively from a subsample of the publications. Categories referred to the type of methodology used, the impact area (ie, remote monitoring, data gathering, communication, self-management, training/education, improve adherence, health promotion), and the type of study. In a second step, all the publications not excluded during the abstract and title review stage underwent a full-text review. All publications that met all eligibility criteria (see below) made up the final sample.

Search Strategy

In February 2012, five electronic databases (CINAHL, Communication and Mass Media Complete, PubMed, PsycINFO, and Web of Science) were systematically searched. The choice of databases was deemed to reflect the multidisciplinary nature of the field. Among the most used medical databases, we decided to include PubMed only, since it comprises MEDLINE, while Embase was excluded because it has a stronger drug coverage, which was not relevant for the purposes of our research. A list of keywords was created around the two domains of “health” and “mobile technology”. A search string was constructed using both the conjunction “AND” and the disjunction “OR” logical operators ([health OR medicine OR medical OR telemedicine OR health care OR “mHealth” OR “mobile health” OR “m - health” OR “mobile - health”] AND [“mobile phone” OR “cell phone” OR “cellphone” OR “cell - phone” OR “smartphone” OR “iPhone” OR “blackberry” OR “android”]). The search was based on metadata, ie, title, abstract, and keywords. Reference lists of selected studies were also checked for other potentially relevant studies.

Selection Criteria

Eligibility criteria for inclusion were as follows: records had to be written in English and discuss/acknowledge the role of mobile technology as a tool for promoting, managing, or monitoring health. This could include interventions, cross-sectional studies, literature reviews, conceptual papers, etc. All articles dealing with health effects of mobile phones (eg, effects of non-ionizing radiation on health or effect of mobile phone use on adolescents) were excluded. Records had to be officially published, either online or in print in a peer-reviewed publication (ie, journal articles, book chapters, and published proceedings papers). This means that poster presentations, (extended) abstracts, and encyclopedia entries were excluded. No time restriction was given; all publication dates were eligible for inclusion. Also, there was no restriction on the field of studies, ie, records that could be classified as social sciences, humanities, medicine, and others were all included.

The exclusion criteria that accounted for the biggest number of excluded articles included the following: the study provided descriptive summaries of mHealth programs but failed to provide an evaluation of the program; study provided a short description of multiple mHealth programs without providing specific details on an mHealth intervention; and the study focused on mHealth application design. The title and abstract review allowed us to exclude system design articles and to better identify all the studies that involved people in the testing of the intervention. A full-text article review was therefore conducted only on studies evaluating and assessing mHealth applications. The categories for full-text review were the following: continent where the study took place, condition addressed, type of technology, features used, basis for the intervention development, study design, sample size, aim of the evaluation, outcome measures, and overall impact assessed.

Results

The flowchart in [Figure 1](#) summarizes the different steps of the literature search and review process. A first search identified

4039 articles. After checking for duplicates, 747 articles met the predefined inclusion criteria. Initially, articles were categorized by type of study: quantitative, qualitative, mixed methods, review, and system design. Since articles in the last typology described the development of a mobile technology but did not include any actual testing, they were excluded from further analysis together with reviews, reducing the final sample to 352 articles. More than half of the 352 studies (56%) included at least some testing of a mobile phone application via proper interventions or in small samples. Most of the studies analyzed (86%) applied a quantitative methodology and were designed to address simultaneously one or more impact areas. An upsurge was noted, starting from 2008, when the articles doubled in comparison with the previous year, and this upward trend reached its maximum in 2011 (36% of the total in a 10-year time period). The search of scientific databases without a time limit yielded an article distribution on the topic over 10 years, from 2002 to 2012.

The final sample for the full-text review included 117 [17-133] articles out of the 352 described above. After title and abstract review, an additional 157 articles were excluded because of no actual testing, while another 78 were excluded during full-text revision for different reasons (eg, no patients involved, mobile device other than phone, study duplicates). Looking at this past decade ([Figure 2](#)), we again observed an upsurge in the field: from 1 article in 2002 to 30 articles in 2011. The largest upsurge again came between 2007 and 2008, when the articles almost doubled, similar to what was already noted during the abstract review phase.

In order to better reflect the objectives of our review and to mirror the development over time, all the results are presented by splitting our observation period in two halves (2002-2007 and 2008-2012). The first period includes 23 articles, while the second period includes 94. The 117 articles in the final sample were published in 77 different journals, which can be grouped in four disciplinary fields: technology, medicine, social sciences, and the intersection between technology and medicine. During the first half of the observed decade, most of the articles on mHealth were published in medical journals (52%) and in journals focused on medicine and technology (44%). The remaining 4% of articles were published in journals focused only on technology. In the second half, the share of articles published in medical journals grew from 52% to 60%, while coverage of the topic by technology journals did not change (4%). At the same time, a decrease in the share of articles published by journals dealing with medicine and technology was observed (from 44% to 35%). In the second period, we found one article from a new disciplinary field, the social sciences.

The geographical areas focusing on this type of research were mainly Europe (34%) and North America (33%). However, if we look at results split by time periods, Europe’s interest seems to decrease from the first period (52%) to the second (30%), and the same happens in Asia (from 17% to 10%). A different picture can be found on all the other continents, where the number of studies in the field increased. This is the case in North America (from 17% to 37%), Australia (from 13% to 15%),

and especially Africa (from 0% to 6%) and South America (from 0% to 2%).

Specific health conditions addressed in the studies ranged from diabetes to mental health, from obesity to well-being and postoperative care. Figure 3 shows the number of articles for every health condition for which mHealth applications were studied. As shown in the graph, diabetes has received a great deal of attention. Moreover, after grouping the conditions into larger classes, it becomes clear that the focus of mHealth research is chronic conditions (74 studies), followed by prevention/well-being (22 studies), and acute conditions (21 studies).

In reviewing the background of the studies, we found that a description of the development of the intervention, and especially of how this was grounded, occurred more often in the second period (84%) than in the first (65%). During the last 5 years, only 1 study was uniquely theory-based (1%), while the majority was evidence-based (73%) or based both on theory and evidence (10%); 15 studies (16%) provided a more general description that was based on neither theory nor previous evidence.

From a methodological point of view, the majority of articles were clinical trials (50%), followed by pilot studies (44%). However, both of these study designs diminished over the last years of our observation period as new types of research designs were introduced, namely observational studies (2% of all articles in the second part of the observation period), case studies (2%), case series (2%), and cross-sectional studies (2%). The samples used to test mobile health applications were mostly small (less than 50 people) in both the first (61%) and the second half (49%) of our observation. Interestingly in the second half, the number of medium-sized samples increased (from 17% to 33%). Larger samples were used in 21 (19%) articles; however, they were more frequent in the first half of the observation (22%) than in the second half (18%).

Moving from research methodology to the actual target of investigation, ie, mobile phones, our classification highlighted a more rigorous and diversified description of the technology used in interventions. In recent years, new kinds of mobile phones have been used, such as smartphones (8%) and ad hoc phones (3%), which are devices developed specifically by the researchers to manage a specific condition. Unfortunately, the kind of mobile phone used was not even specified most of the time (71% of the overall sample).

We identified seven main categories of mobile features used in the studies, and an article could fall in one or more of these (ie, the categories were not mutually exclusive). Half of the studies (49%) applied text messaging, and 32% applied some features developed ad hoc for a specific condition. Add-ons (eg, a glucometer to measure blood sugar or a pedometer for physical activity) were used in 12% of the cases together with ad hoc features. Other features such as voice (10%), video (6%), and

multimedia messaging service (MMS) (3%) were used less frequently. Native applications for smartphones were applied in 7 studies (6%) out of the 8 using smartphones. However, none of them applied already existing and publicly available apps.

The impact areas to which interventions were directed were coded into seven categories, again not mutually exclusive. The majority of articles addressed health promotion (38%) and self-management (33%), but also communication (22%), remote monitoring (21%), data gathering (21%), improvement of adherence (20%), and training/education (13%). The focus on most of these areas increased over time, eg, on self-management (from 30% to 33%) and communication (from 17% to 23%). Only health promotion (from 29% to 27%) and training/education (from 13% to 10%) had a slight decrease.

Regarding the aims of the interventions, both the evaluation of the technology itself (35%) and of its impact on health outcomes (43%) dominated in the first 5 years. In the second half, however, interest clearly moved toward evaluating the impact of mobile technology on health outcomes (73%). While the majority of the studies investigated only the impact of the mobile application on health outcomes (51%), some also assessed both the technology and its impact on health outcomes (22%).

Another point of interest was the outcome measure used to assess the impact of mobile phones. In the majority of cases, the outcome measures were a combination of both self-reported and objective data (44% of the overall sample). If we look at the evolution over time, self-reported measures increased (from 9% to 20%), whereas objective measures decreased slightly (from 39% to 36%), and this was also the case for the combination of self-reported and objective measures (from 52% to 43%).

Our examination of the type of data collected showed that clinical measures were often the only outcomes observed (30%), and this phenomenon increased over time (from 22% to 31%). 14% of the articles were focused only on user assessment of the technology, even if this decreased during the observation period (from 17% to 13%). Psychosocial measures were the outcome in 9% of the studies, and this increased slightly over time (from 9% to 10%). The remaining articles (53%) considered outcomes deriving from all possible combinations of these main three. The most frequent combination was clinical measures together with user assessment of technology (17%).

An overall positive impact of the intervention was described by a total of 69 studies (60%). In the first period, the impact of the interventions was mainly either mixed (43%) or positive (57%). In the second part of the observation, the number of interventions with a positive impact slightly increased (60%), while the number of those with mixed impact decreased (33%). In this second period, interventions with negative (6%) or no impact (1%) were reported as well.

Figure 1. Summary of literature search and review process.




Figure 2. Distribution of articles over time.





Figure 3. Conditions addressed by mHealth applications in mHealth research.



Discussion

This systematic literature review has found encouraging trends pointing toward the development of mHealth as an autonomous field of study, which is of interest to different disciplines. In the case of social sciences for instance, this is shown by the publication of a special issue of the *Journal of Health Communication* dedicated entirely to mHealth in May 2012.

We took this into account but did not include it in the current review because of the date of publication.

The increase in the number of articles over the past decade indicates an increasing interest in peer-reviewed scientific literature on the topic. In particular, the number of articles almost doubled from 2007 to 2008. All continents engaged in research on mHealth during the last 5 years. Many different conditions were addressed by the studies analyzed, with an evident focus on chronic conditions. More recently, the

development of the interventions became more accurately described and grounded on past evidence or theory. In the last decade, the first in the history of mHealth, new methodologies were applied to the field, and while the majority of the studies were pilot studies or clinical trials, some new approaches were introduced. At the same time, the samples used in such studies grew, thus pointing toward a more reliable assessment. Newer and more advanced technologies were tested; however, the potential of smartphones does not seem to have been fully exploited yet. Indeed, half of the included studies applied very basic features of mobile phones, such as text messaging, which corroborates evidence already established in the field [7]. One third of the studies used features from add-ons, and only a few other examples used video, images, or MMS. Mobile technology interventions were directed to different impact areas, and most frequently to health promotion and self-management. Interestingly, research moved from an evaluation of the technology itself to an assessment of its impact on health outcomes, and an increase in the number of interventions with an overall positive impact was observed. These observations point toward a field that is becoming more structured, coherent, and solid. Among the outcome measures used, both self-reported and objective measures could be found, and in some cases were combined. In the last 5 years, the use of self-reported measures increased. Clinical measures only were observed as outcomes in almost one third of the studies, and an increasing trend was evident. Consistent with the observed trend towards an assessment of the impact of technology, we noticed an increase in psychosocial and clinical measures, sometimes in combination with users' assessment of the technology, and a decrease of assessments of the technology alone.

Recommendations

Although these findings are encouraging and can be seen as indicators of a promising field, they highlight certain gaps that future research should address. So far, the focus of mHealth interventions has been on chronic conditions, similar to eHealth [134]. However, it would be advisable to explore the impact of mobile health for acute conditions as well. Because of their wireless cellular communication capability, mobile phones allow users to have continuous, interactive communication from any location. In our view, this characteristic of mobile phones makes them an ideal tool to address in real-time the specific needs of patients experiencing acute conditions.

Another recommendation is to address methodological issues such as whether clinical trials are the most suitable design to use at every stage of research in this field. The studies we analyzed showed a diversification toward the second half of the decade, which seems to reflect more positive outcomes and a stronger evidence-based development of the interventions. Consistent with the conclusions of other authors [3,5,7,135], we believe the field will greatly profit from a diversification in research methodologies and that the multidisciplinary approach offered by different areas could be a fertile ground for the development of the field, both from a theoretical and methodological perspective.

Our last recommendation is for research to fully exploit the potential of technologies, especially of smartphones. We

expected in our review to find more results detailing applications of new built-in features, which are the "specialty" of smartphones. However we found that only a few interventions aimed at assessing the impact of native applications for smartphones had been reported so far in the literature. Moreover, in all the cases, the apps were not available to the public but had been created ad hoc for research purposes.

This last recommendation brings up a new topic of discussion. Currently, there are more than 15,000 health-related apps (free and paid) on app stores, but we were not able to find any study assessing any of them. So what is publicly available has not been evaluated, and what has been evaluated is not publicly available. At least three possible scenarios could explain the lack of scholarly interest in studying the effects of interventions based on publicly available apps. First, no one, so far, has conducted such studies. If this is true, it provides great opportunities for investigation in this area, since some basic features of mobile health such as text messaging have already proved to be effective. A second scenario could be that it is still too early for results of such assessments to have been published. However, this seems to be less plausible, because iPhones and similar devices have already been on the market for 5 years. Finally, it could be that studies evaluating the impact of native applications have indeed been published, but not in the peer-reviewed scientific literature. A search of the gray literature, such as more consumer-oriented magazines, websites, or blogs (eg, iMedicalApps) could then yield some results. If this last, and more plausible, explanation is true, then we need to document the exploitation of smartphones in a way that is easily accessible to the scientific community. If only a minority of good quality applications and solutions seek clinical mHealth research to prove effectiveness, we should question why this is happening. It is thus essential to understand why scientific literature is not keeping up with advancements of the field, to find where any discussion is taking place, and to find the evidence of apps' effectiveness, which seems to be missing [3].

Limitations

A limitation of our study, which is common to most systematic literature reviews, resides in the fact that the last articles analyzed were published in February 2012. In this constantly evolving field, this could make a difference, especially because we are well aware that the publication process is often lengthy and time-consuming, and therefore we could have missed some studies on smartphones.

A second possible limitation lies in the choice of databases. For our systematic search, we focused on more medical and social sciences-oriented databases. However a preliminary search conducted on a more technology-oriented database (ACM) resulted in a long list of peer-reviewed articles mainly focused on system design, which we would have excluded from the final sample since our research interest was on the impact of mobile technology on health outcomes.

Conclusion

With this systematic literature review, we sought to map a field that is becoming more and more visible in the literature. This

review is an essential first step to the understanding of strengths and limitations in mHealth research.

Some questions remain about the lack of information on the newest technological opportunities and on the best methodologies to assess mHealth's impact on health outcomes.

Is scientific literature the appropriate place to find studies on the effectiveness of mobile applications? Are we exploiting all the potential of smartphones? Why is the scientific research not keeping up with the market evolution? Further research will help to answer these questions.

Authors' Contributions

All authors took part in the design of the search strategy. MF and ND conducted the search, extracted the data, and analyzed the content. MF drafted the manuscript and together with ND contributed to the writing of the final manuscript. MF revised the manuscript with the input and advice from all authors.

Conflicts of Interest

None declared.

References

1. Free C, Phillips G, Felix L, Galli L, Patel V, Edwards P. The effectiveness of M-health technologies for improving health and health services: a systematic review protocol. *BMC Res Notes* 2010;3:250 [FREE Full text] [doi: [10.1186/1756-0500-3-250](https://doi.org/10.1186/1756-0500-3-250)] [Medline: [20925916](https://pubmed.ncbi.nlm.nih.gov/20925916/)]
2. Riley WT, Rivera DE, Atienza AA, Nilsen W, Allison SM, Mermelstein R. Health behavior models in the age of mobile interventions: are our theories up to the task? *Transl Behav Med* 2011 Mar 1;1(1):53-71 [FREE Full text] [doi: [10.1007/s13142-011-0021-7](https://doi.org/10.1007/s13142-011-0021-7)] [Medline: [21796270](https://pubmed.ncbi.nlm.nih.gov/21796270/)]
3. Whittaker R. Issues in mHealth: findings from key informant interviews. *J Med Internet Res* 2012;14(5):e129 [FREE Full text] [doi: [10.2196/jmir.1989](https://doi.org/10.2196/jmir.1989)] [Medline: [23032424](https://pubmed.ncbi.nlm.nih.gov/23032424/)]
4. Fogg BJ, Adler R. *Texting 4 Health: A Simple, Powerful Way to Change Lives* (Stanford University) (Stanford University). Stanford, CA: Captology Media; 2009.
5. Sherry JM, Ratzan SC. Measurement and evaluation outcomes for mHealth communication: don't we have an app for that? *J Health Commun* 2012 May;17 Suppl 1:1-3. [doi: [10.1080/10810730.2012.670563](https://doi.org/10.1080/10810730.2012.670563)] [Medline: [22548591](https://pubmed.ncbi.nlm.nih.gov/22548591/)]
6. Guy R, Hocking J, Wand H, Stott S, Ali H, Kaldor J. How effective are short message service reminders at increasing clinic attendance? A meta-analysis and systematic review. *Health Serv Res* 2012 Apr;47(2):614-632. [doi: [10.1111/j.1475-6773.2011.01342.x](https://doi.org/10.1111/j.1475-6773.2011.01342.x)] [Medline: [22091980](https://pubmed.ncbi.nlm.nih.gov/22091980/)]
7. Evans WD, Abroms LC, Poropatich R, Nielsen PE, Wallace JL. Mobile health evaluation methods: the Text4baby case study. *J Health Commun* 2012;17 Suppl 1:22-29. [doi: [10.1080/10810730.2011.649157](https://doi.org/10.1080/10810730.2011.649157)] [Medline: [22548595](https://pubmed.ncbi.nlm.nih.gov/22548595/)]
8. Collins English Online Dictionary. 2013. "Smartphone" URL: <http://www.collinsdictionary.com/dictionary/english/smartphone?showCookiePolicy=true> [accessed 2013-04-10] [WebCite Cache ID 6Fm8JS74o]
9. Smith A. Americans and their cell phones. Washington, DC: Pew Internet & American Life Project; 2011 Aug 15. URL: <http://pewinternet.org/Reports/2011/Cell-Phones.aspx> [accessed 2013-04-10] [WebCite Cache ID 6Fm0QIEp8]
10. International Communications Union. 2012. URL: http://www.itu.int/ITU-D/ict/publications/material/LDB_ICT_2012.pdf [accessed 2013-04-10] [WebCite Cache ID 6Fm0SamWW]
11. Chomutare T, Fernandez-Luque L, Arsand E, Hartvigsen G. Features of mobile diabetes applications: review of the literature and analysis of current applications compared against evidence-based guidelines. *J Med Internet Res* 2011;13(3):e65 [FREE Full text] [doi: [10.2196/jmir.1874](https://doi.org/10.2196/jmir.1874)] [Medline: [21979293](https://pubmed.ncbi.nlm.nih.gov/21979293/)]
12. Krishna S, Boren SA, Balas EA. Healthcare via cell phones: a systematic review. *Telemed J E Health* 2009 Apr;15(3):231-240. [doi: [10.1089/tmj.2008.0099](https://doi.org/10.1089/tmj.2008.0099)] [Medline: [19382860](https://pubmed.ncbi.nlm.nih.gov/19382860/)]
13. Klasnja P, Pratt W. Healthcare in the pocket: mapping the space of mobile-phone health interventions. *J Biomed Inform* 2012 Feb;45(1):184-198 [FREE Full text] [doi: [10.1016/j.jbi.2011.08.017](https://doi.org/10.1016/j.jbi.2011.08.017)] [Medline: [21925288](https://pubmed.ncbi.nlm.nih.gov/21925288/)]
14. Research2guidance. 2013. The research mobile specialists URL: <http://www.research2guidance.com/> [accessed 2013-04-07] [WebCite Cache ID 6Fhsvb7a4]
15. Boulos MN, Wheeler S, Tavares C, Jones R. How smartphones are changing the face of mobile and participatory healthcare: an overview, with example from eCAALYX. *Biomed Eng Online* 2011;10:24 [FREE Full text] [doi: [10.1186/1475-925X-10-24](https://doi.org/10.1186/1475-925X-10-24)] [Medline: [21466669](https://pubmed.ncbi.nlm.nih.gov/21466669/)]
16. Terry M. Medical Apps for Smartphones. *Telemed J E Health* 2010 Feb;16(1):17-22. [doi: [10.1089/tmj.2010.9999](https://doi.org/10.1089/tmj.2010.9999)] [Medline: [20070172](https://pubmed.ncbi.nlm.nih.gov/20070172/)]
17. Aguilera A, Muñoz RF. Text messaging as an adjunct to CBT in low-income populations: A usability and feasibility pilot study. *Professional Psychology: Research and Practice* 2011 2011;42(6):472-478. [doi: [10.1037/a0025499](https://doi.org/10.1037/a0025499)]
18. Anhøj J, Møldrup C. Feasibility of collecting diary data from asthma patients through mobile phones and SMS (short message service): response rate analysis and focus group evaluation from a pilot study. *J Med Internet Res* 2004 Dec 2;6(4):e42 [FREE Full text] [doi: [10.2196/jmir.6.4.e42](https://doi.org/10.2196/jmir.6.4.e42)] [Medline: [15631966](https://pubmed.ncbi.nlm.nih.gov/15631966/)]

19. Arsand E, Tataru N, Østengen G, Hartvigsen G. Mobile phone-based self-management tools for type 2 diabetes: the few touch application. *J Diabetes Sci Technol* 2010 Mar;4(2):328-336 [FREE Full text] [Medline: 20307393]
20. Beebe LH, Smith K, Bennett C, Bentley K, Walters AB, Hancock B, et al. Keeping in touch. Cell phone use in people with schizophrenia disorders. *J Psychosoc Nurs Ment Health Serv* 2010 Apr;48(4):32-37. [doi: [10.3928/02793695-20100304-99](https://doi.org/10.3928/02793695-20100304-99)] [Medline: 20349885]
21. Blasco A, Carmona M, Fernández-Lozano I, Salvador CH, Pascual M, Sagredo PG, et al. Evaluation of a telemedicine service for the secondary prevention of coronary artery disease. *J Cardiopulm Rehabil Prev* 2012;32(1):25-31. [doi: [10.1097/HCR.0b013e3182343aa7](https://doi.org/10.1097/HCR.0b013e3182343aa7)] [Medline: 22113368]
22. Bopp JM, Miklowitz DJ, Goodwin GM, Stevens W, Rendell JM, Geddes JR. The longitudinal course of bipolar disorder as revealed through weekly text messaging: a feasibility study. *Bipolar Disord* 2010 May;12(3):327-334 [FREE Full text] [doi: [10.1111/j.1399-5618.2010.00807.x](https://doi.org/10.1111/j.1399-5618.2010.00807.x)] [Medline: 20565440]
23. Bramley D, Riddell T, Whittaker R, Corbett T, Lin RB, Wills M, et al. Smoking cessation using mobile phone text messaging is as effective in Maori as non-Maori. *N Z Med J* 2005 Jun 3;118(1216):U1494. [Medline: 15937529]
24. Brendryen H, Drozd F, Kraft P. A digital smoking cessation program delivered through internet and cell phone without nicotine replacement (happy ending): randomized controlled trial. *J Med Internet Res* 2008 Jan;10(5):e51 [FREE Full text] [doi: [10.2196/jmir.1005](https://doi.org/10.2196/jmir.1005)] [Medline: 19087949]
25. Burns MN, Begale M, Duffecy J, Gergle D, Karr CJ, Giangrande E, et al. Harnessing context sensing to develop a mobile intervention for depression. *J Med Internet Res* 2011 Jan;13(3):e55 [FREE Full text] [doi: [10.2196/jmir.1838](https://doi.org/10.2196/jmir.1838)] [Medline: 21840837]
26. Carroll AE, DiMeglio LA, Stein S, Marrero DG. Using a cell phone-based glucose monitoring system for adolescent diabetes management. *Diabetes Educ* 2011;37(1):59-66. [doi: [10.1177/0145721710387163](https://doi.org/10.1177/0145721710387163)] [Medline: 21106908]
27. Charpentier G, Benhamou PY, Dardari D, Clergeot A, Franc S, Schaepelynck-Belicar P, TeleDiab Study Group. The Diabeo software enabling individualized insulin dose adjustments combined with telemedicine support improves HbA1c in poorly controlled type 1 diabetic patients: a 6-month, randomized, open-label, parallel-group, multicenter trial (TeleDiab 1 Study). *Diabetes Care* 2011 Mar;34(3):533-539 [FREE Full text] [doi: [10.2337/dc.10-1259](https://doi.org/10.2337/dc.10-1259)] [Medline: 21266648]
28. Chi BH, Stringer JS. Mobile phones to improve HIV treatment adherence. *Lancet* 2010 Nov 27;376(9755):1807-1808. [doi: [10.1016/S0140-6736\(10\)62046-6](https://doi.org/10.1016/S0140-6736(10)62046-6)] [Medline: 21071073]
29. Cho JH, Lee HC, Lim DJ, Kwon HS, Yoon KH. Mobile communication using a mobile phone with a glucometer for glucose control in Type 2 patients with diabetes: as effective as an Internet-based glucose monitoring system. *J Telemed Telecare* 2009;15(2):77-82. [doi: [10.1258/jtt.2008.080412](https://doi.org/10.1258/jtt.2008.080412)] [Medline: 19246607]
30. Cocosila M, Archer N, Haynes RB, Yuan Y. Can wireless text messaging improve adherence to preventive activities? Results of a randomised controlled trial. *Int J Med Inform* 2009 Apr;78(4):230-238. [doi: [10.1016/j.ijmedinf.2008.07.011](https://doi.org/10.1016/j.ijmedinf.2008.07.011)] [Medline: 18778967]
31. Cocosila M, Archer N, Yuan Y. Would people pay for text messaging health reminders? *Telemed J E Health* 2008 Dec;14(10):1091-1095. [doi: [10.1089/tmj.2008.0047](https://doi.org/10.1089/tmj.2008.0047)] [Medline: 19119832]
32. Cummings E, Hauser J, Cameron-Tucker H, Fitzpatrick P, Jessup M, Walters EH, et al. Enhancing self-efficacy for self-management in people with cystic fibrosis. *Stud Health Technol Inform* 2011;169:33-37. [Medline: 21893709]
33. da Costa TM, Salomão PL, Martha AS, Pisa IT, Sigulem D. The impact of short message service text messages sent as appointment reminders to patients' cell phones at outpatient clinics in São Paulo, Brazil. *Int J Med Inform* 2010 Jan;79(1):65-70. [doi: [10.1016/j.ijmedinf.2009.09.001](https://doi.org/10.1016/j.ijmedinf.2009.09.001)] [Medline: 19783204]
34. de Oliveira R, Cherubini M, Oliver N. MoviPill? Improving Medication Compliance for Elders Using a Mobile Persuasive Social Game. In: Proceedings of the 12th ACM International Conference on Ubiquitous Computing. 2010 Presented at: 12th ACM International Conference on Ubiquitous Computing; September 26-29, 2010; Copenhagen p. 251-260.
35. Depp CA, Mausbach B, Granholm E, Cardenas V, Ben-Zeev D, Patterson TL, et al. Mobile interventions for severe mental illness: design and preliminary data from three approaches. *J Nerv Ment Dis* 2010 Oct;198(10):715-721 [FREE Full text] [doi: [10.1097/NMD.0b013e3181f49ea3](https://doi.org/10.1097/NMD.0b013e3181f49ea3)] [Medline: 20921861]
36. Dunton GF, Liao Y, Intille SS, Spruijt-Metz D, Pentz M. Investigating children's physical activity and sedentary behavior using ecological momentary assessment with mobile phones. *Obesity (Silver Spring)* 2011 Jun;19(6):1205-1212. [doi: [10.1038/oby.2010.302](https://doi.org/10.1038/oby.2010.302)] [Medline: 21164502]
37. Faridi Z, Liberti L, Shuval K, Northrup V, Ali A, Katz DL. Evaluating the impact of mobile telephone technology on type 2 diabetic patients' self-management: the NICHE pilot study. *J Eval Clin Pract* 2008 Jun;14(3):465-469. [doi: [10.1111/j.1365-2753.2007.00881.x](https://doi.org/10.1111/j.1365-2753.2007.00881.x)] [Medline: 18373577]
38. Ferrer-Roca O, Cárdenas A, Diaz-Cardama A, Pulido P. Mobile phone text messaging in the management of diabetes. *J Telemed Telecare* 2004;10(5):282-285. [Medline: 15494086]
39. Finkelstein J, Wood J, Cha E. Introducing a BlackBerry eLearning Platform for Interactive Hypertension Education. In: Proceedings of the 2nd International Conference on Mobile, Hybrid, and On-Line Learning. 2010 Presented at: 2nd International Conference on Mobile, Hybrid, and On-Line Learning; Feb. 10-16, 2010; Washington, DC p. 77-81.
40. Fjeldsoe BS, Miller YD, Marshall AL. MobileMums: a randomized controlled trial of an SMS-based physical activity intervention. *Ann Behav Med* 2010 May;39(2):101-111. [doi: [10.1007/s12160-010-9170-z](https://doi.org/10.1007/s12160-010-9170-z)] [Medline: 20174902]

41. Free C, Whittaker R, Knight R, Abramsky T, Rodgers A, Roberts IG. Txt2stop: a pilot randomised controlled trial of mobile phone-based smoking cessation support. *Tob Control* 2009 Apr;18(2):88-91. [doi: [10.1136/tc.2008.026146](https://doi.org/10.1136/tc.2008.026146)] [Medline: [19318534](https://pubmed.ncbi.nlm.nih.gov/19318534/)]
42. Free C, Knight R, Robertson S, Whittaker R, Edwards P, Zhou W, et al. Smoking cessation support delivered via mobile phone text messaging (txt2stop): a single-blind, randomised trial. *Lancet* 2011 Jul 2;378(9785):49-55 [FREE Full text] [doi: [10.1016/S0140-6736\(11\)60701-0](https://doi.org/10.1016/S0140-6736(11)60701-0)] [Medline: [21722952](https://pubmed.ncbi.nlm.nih.gov/21722952/)]
43. Fukuoka Y, Kamitani E, Dracup K, Jong SS. New insights into compliance with a mobile phone diary and pedometer use in sedentary women. *J Phys Act Health* 2011 Mar;8(3):398-403. [Medline: [21487139](https://pubmed.ncbi.nlm.nih.gov/21487139/)]
44. Fukuoka Y, Vittinghoff E, Jong SS, Haskell W. Innovation to motivation--pilot study of a mobile phone intervention to increase physical activity among sedentary women. *Prev Med* 2010;51(3-4):287-289 [FREE Full text] [doi: [10.1016/j.ypmed.2010.06.006](https://doi.org/10.1016/j.ypmed.2010.06.006)] [Medline: [20600263](https://pubmed.ncbi.nlm.nih.gov/20600263/)]
45. Gerber BS, Stolley MR, Thompson AL, Sharp LK, Fitzgibbon ML. Mobile phone text messaging to promote healthy behaviors and weight loss maintenance: a feasibility study. *Health Informatics J* 2009 Mar;15(1):17-25 [FREE Full text] [doi: [10.1177/1460458208099865](https://doi.org/10.1177/1460458208099865)] [Medline: [19218309](https://pubmed.ncbi.nlm.nih.gov/19218309/)]
46. Gold J, Aitken CK, Dixon HG, Lim MS, Gouillou M, Spelman T, et al. A randomised controlled trial using mobile advertising to promote safer sex and sun safety to young people. *Health Educ Res* 2011 Oct;26(5):782-794 [FREE Full text] [doi: [10.1093/her/cyr020](https://doi.org/10.1093/her/cyr020)] [Medline: [21447750](https://pubmed.ncbi.nlm.nih.gov/21447750/)]
47. Gold J, Lim MS, Hocking JS, Keogh LA, Spelman T, Hellard ME. Determining the impact of text messaging for sexual health promotion to young people. *Sex Transm Dis* 2011 Apr;38(4):247-252. [doi: [10.1097/OLQ.0b013e3181f68d7b](https://doi.org/10.1097/OLQ.0b013e3181f68d7b)] [Medline: [20966830](https://pubmed.ncbi.nlm.nih.gov/20966830/)]
48. Granholm E, Ben-Zeev D, Link PC, Bradshaw KR, Holden JL. Mobile Assessment and Treatment for Schizophrenia (MATS): a pilot trial of an interactive text-messaging intervention for medication adherence, socialization, and auditory hallucinations. *Schizophr Bull* 2012 May;38(3):414-425. [doi: [10.1093/schbul/sbr155](https://doi.org/10.1093/schbul/sbr155)] [Medline: [22080492](https://pubmed.ncbi.nlm.nih.gov/22080492/)]
49. Grassi A, Preziosa A, Villani D, Riva G. A Relaxing Journey: the Use of Mobile Phones for Well-Being Improvement. *Annual Review of CyberTherapy and Telemedicine* 2007;5:123-131.
50. Haapala I, Barengo NC, Biggs S, Surakka L, Manninen P. Weight loss by mobile phone: a 1-year effectiveness study. *Public Health Nutr* 2009 Dec;12(12):2382-2391. [doi: [10.1017/S1368980009005230](https://doi.org/10.1017/S1368980009005230)] [Medline: [19323865](https://pubmed.ncbi.nlm.nih.gov/19323865/)]
51. Hanauer DA, Wentzell K, Laffel N, Laffel LM. Computerized Automated Reminder Diabetes System (CARDS): e-mail and SMS cell phone text messaging reminders to support diabetes management. *Diabetes Technol Ther* 2009 Feb;11(2):99-106. [doi: [10.1089/dia.2008.0022](https://doi.org/10.1089/dia.2008.0022)] [Medline: [19848576](https://pubmed.ncbi.nlm.nih.gov/19848576/)]
52. Hoffman JA, Cunningham JR, Suleh AJ, Sundsmo A, Dekker D, Vago F, et al. Mobile direct observation treatment for tuberculosis patients: a technical feasibility pilot using mobile phones in Nairobi, Kenya. *Am J Prev Med* 2010 Jul;39(1):78-80. [doi: [10.1016/j.amepre.2010.02.018](https://doi.org/10.1016/j.amepre.2010.02.018)] [Medline: [20537846](https://pubmed.ncbi.nlm.nih.gov/20537846/)]
53. Hurling R, Catt M, Boni MD, Fairley BW, Hurst T, Murray P, et al. Using internet and mobile phone technology to deliver an automated physical activity program: randomized controlled trial. *J Med Internet Res* 2007;9(2):e7 [FREE Full text] [doi: [10.2196/jmir.9.2.e7](https://doi.org/10.2196/jmir.9.2.e7)] [Medline: [17478409](https://pubmed.ncbi.nlm.nih.gov/17478409/)]
54. Istepanian RS, Zitouni K, Harry D, Moutosammy N, Sungoor A, Tang B, et al. Evaluation of a mobile phone telemonitoring system for glycaemic control in patients with diabetes. *J Telemed Telecare* 2009;15(3):125-128. [doi: [10.1258/jtt.2009.003006](https://doi.org/10.1258/jtt.2009.003006)] [Medline: [19364893](https://pubmed.ncbi.nlm.nih.gov/19364893/)]
55. Joo NS, Kim BT. Mobile phone short message service messaging for behaviour modification in a community-based weight control programme in Korea. *J Telemed Telecare* 2007;13(8):416-420. [doi: [10.1258/135763307783064331](https://doi.org/10.1258/135763307783064331)] [Medline: [18078554](https://pubmed.ncbi.nlm.nih.gov/18078554/)]
56. Juzang I, Fortune T, Black S, Wright E, Bull S. A pilot programme using mobile phones for HIV prevention. *J Telemed Telecare* 2011;17(3):150-153. [doi: [10.1258/jtt.2010.091107](https://doi.org/10.1258/jtt.2010.091107)] [Medline: [21270049](https://pubmed.ncbi.nlm.nih.gov/21270049/)]
57. Kalichman SC, Kalichman MO, Cherry C, Swetzes C, Amaral CM, White D, et al. Brief behavioral self-regulation counseling for HIV treatment adherence delivered by cell phone: an initial test of concept trial. *AIDS Patient Care STDS* 2011 May;25(5):303-310 [FREE Full text] [doi: [10.1089/apc.2010.0367](https://doi.org/10.1089/apc.2010.0367)] [Medline: [21457056](https://pubmed.ncbi.nlm.nih.gov/21457056/)]
58. Katz DL, Nordwall B. Novel interactive cell-phone technology for health enhancement. *J Diabetes Sci Technol* 2008 Jan;2(1):147-153 [FREE Full text] [Medline: [19885191](https://pubmed.ncbi.nlm.nih.gov/19885191/)]
59. Katz KS, Rodan M, Milligan R, Tan S, Courtney L, Gantz M, et al. Efficacy of a randomized cell phone-based counseling intervention in postponing subsequent pregnancy among teen mothers. *Matern Child Health J* 2011 Dec;15 Suppl 1:S42-S53. [doi: [10.1007/s10995-011-0860-3](https://doi.org/10.1007/s10995-011-0860-3)] [Medline: [21809218](https://pubmed.ncbi.nlm.nih.gov/21809218/)]
60. Kearney N, McCann L, Norrie J, Taylor L, Gray P, McGee-Lennon M, et al. Evaluation of a mobile phone-based, advanced symptom management system (ASyMS) in the management of chemotherapy-related toxicity. *Support Care Cancer* 2009 Apr;17(4):437-444. [doi: [10.1007/s00520-008-0515-0](https://doi.org/10.1007/s00520-008-0515-0)] [Medline: [18953579](https://pubmed.ncbi.nlm.nih.gov/18953579/)]
61. Khokhar A. Short text messages (SMS) as a reminder system for making working women from Delhi Breast Aware. *Asian Pac J Cancer Prev* 2009;10(2):319-322 [FREE Full text] [Medline: [19537904](https://pubmed.ncbi.nlm.nih.gov/19537904/)]

62. Koller S, Hofmann-Wellenhof R, Hayn D, Weger W, Kastner P, Schreier G, et al. Teledermatological monitoring of psoriasis patients on biologic therapy. *Acta Derm Venereol* 2011 Oct;91(6):680-685 [FREE Full text] [doi: [10.2340/00015555-1148](https://doi.org/10.2340/00015555-1148)] [Medline: [21879250](https://pubmed.ncbi.nlm.nih.gov/21879250/)]
63. Kollmann A, Riedl M, Kastner P, Schreier G, Ludvik B. Feasibility of a mobile phone-based data service for functional insulin treatment of type 1 diabetes mellitus patients. *J Med Internet Res* 2007 Jan;9(5):e36 [FREE Full text] [doi: [10.2196/jmir.9.5.e36](https://doi.org/10.2196/jmir.9.5.e36)] [Medline: [18166525](https://pubmed.ncbi.nlm.nih.gov/18166525/)]
64. Koshy E, Car J, Majeed A. Effectiveness of mobile-phone short message service (SMS) reminders for ophthalmology outpatient appointments: observational study. *BMC Ophthalmol* 2008 Jan;8:9 [FREE Full text] [doi: [10.1186/1471-2415-8-9](https://doi.org/10.1186/1471-2415-8-9)] [Medline: [18513438](https://pubmed.ncbi.nlm.nih.gov/18513438/)]
65. Kristjánssdóttir Ó, Fors EA, Eide E, Finset A, van Dulmen S, Wigers SH, et al. Written online situational feedback via mobile phone to support self-management of chronic widespread pain: a usability study of a Web-based intervention. *BMC Musculoskelet Disord* 2011 Jan;12:51 [FREE Full text] [doi: [10.1186/1471-2474-12-51](https://doi.org/10.1186/1471-2474-12-51)] [Medline: [21352516](https://pubmed.ncbi.nlm.nih.gov/21352516/)]
66. Kunawararak P, Pongpanich S, Chantawong S, Pokaew P, Traisathit P, Srithanaviboonchai K, et al. Tuberculosis treatment with mobile-phone medication reminders in northern Thailand. *Southeast Asian J Trop Med Public Health* 2011 Nov;42(6):1444-1451. [Medline: [22299414](https://pubmed.ncbi.nlm.nih.gov/22299414/)]
67. Kunutsor S, Walley J, Katabira E, Muchuro S, Balidawa H, Namagala E, et al. Using mobile phones to improve clinic attendance amongst an antiretroviral treatment cohort in rural Uganda: a cross-sectional and prospective study. *AIDS Behav* 2010 Dec;14(6):1347-1352. [doi: [10.1007/s10461-010-9780-2](https://doi.org/10.1007/s10461-010-9780-2)] [Medline: [20700644](https://pubmed.ncbi.nlm.nih.gov/20700644/)]
68. Lakkis NA, Atfeh AM, El-Zein YR, Mahmassani DM, Hamadeh GN. The effect of two types of sms-texts on the uptake of screening mammogram: a randomized controlled trial. *Prev Med* 2011 Oct;53(4-5):325-327. [doi: [10.1016/j.ypmed.2011.08.013](https://doi.org/10.1016/j.ypmed.2011.08.013)] [Medline: [21871480](https://pubmed.ncbi.nlm.nih.gov/21871480/)]
69. Lee HJ, Lee SH, Ha KS, Jang HC, Chung WY, Kim JY, et al. Ubiquitous healthcare service using Zigbee and mobile phone for elderly patients. *Int J Med Inform* 2009 Mar;78(3):193-198. [doi: [10.1016/j.ijmedinf.2008.07.005](https://doi.org/10.1016/j.ijmedinf.2008.07.005)] [Medline: [18760959](https://pubmed.ncbi.nlm.nih.gov/18760959/)]
70. Leong KC, Chen WS, Leong KW, Mastura I, Mimi O, Sheikh MA, et al. The use of text messaging to improve attendance in primary care: a randomized controlled trial. *Fam Pract* 2006 Dec;23(6):699-705 [FREE Full text] [doi: [10.1093/fampra/cml044](https://doi.org/10.1093/fampra/cml044)] [Medline: [16916871](https://pubmed.ncbi.nlm.nih.gov/16916871/)]
71. Lester RT, Ritvo P, Mills EJ, Kariri A, Karanja S, Chung MH, et al. Effects of a mobile phone short message service on antiretroviral treatment adherence in Kenya (WelTel Kenya1): a randomised trial. *Lancet* 2010 Nov 27;376(9755):1838-1845. [doi: [10.1016/S0140-6736\(10\)61997-6](https://doi.org/10.1016/S0140-6736(10)61997-6)] [Medline: [21071074](https://pubmed.ncbi.nlm.nih.gov/21071074/)]
72. Lim EJ, Haar J, Morgan J. Can text messaging results reduce time to treatment of Chlamydia trachomatis? *Sex Transm Infect* 2008 Dec;84(7):563-564. [doi: [10.1136/sti.2008.032441](https://doi.org/10.1136/sti.2008.032441)] [Medline: [18723584](https://pubmed.ncbi.nlm.nih.gov/18723584/)]
73. Lim MS, Hocking JS, Aitken CK, Fairley CK, Jordan L, Lewis JA, et al. Impact of text and email messaging on the sexual health of young people: a randomised controlled trial. *J Epidemiol Community Health* 2012 Jan;66(1):69-74. [doi: [10.1136/jech.2009.100396](https://doi.org/10.1136/jech.2009.100396)] [Medline: [21415232](https://pubmed.ncbi.nlm.nih.gov/21415232/)]
74. Lim MS, Sacks-Davis R, Aitken CK, Hocking JS, Hellard ME. Randomised controlled trial of paper, online and SMS diaries for collecting sexual behaviour information from young people. *J Epidemiol Community Health* 2010 Oct;64(10):885-889. [doi: [10.1136/jech.2008.085316](https://doi.org/10.1136/jech.2008.085316)] [Medline: [19767322](https://pubmed.ncbi.nlm.nih.gov/19767322/)]
75. Lim S, Kang SM, Shin H, Lee HJ, Won Yoon J, Yu SH, et al. Improved glycemic control without hypoglycemia in elderly diabetic patients using the ubiquitous healthcare service, a new medical information system. *Diabetes Care* 2011 Feb;34(2):308-313 [FREE Full text] [doi: [10.2337/dc10-1447](https://doi.org/10.2337/dc10-1447)] [Medline: [21270188](https://pubmed.ncbi.nlm.nih.gov/21270188/)]
76. Logan AG, McIsaac WJ, Tisler A, Irvine MJ, Saunders A, Dunai A, et al. Mobile phone-based remote patient monitoring system for management of hypertension in diabetic patients. *Am J Hypertens* 2007 Sep;20(9):942-948. [doi: [10.1016/j.amjhyper.2007.03.020](https://doi.org/10.1016/j.amjhyper.2007.03.020)] [Medline: [17765133](https://pubmed.ncbi.nlm.nih.gov/17765133/)]
77. Macedo LG, Maher CG, Latimer J, McAuley JH. Feasibility of using short message service to collect pain outcomes in a low back pain clinical trial. *Spine* 2012 Jun 1;37(13):1151-1155. [doi: [10.1097/BRS.0b013e3182422df0](https://doi.org/10.1097/BRS.0b013e3182422df0)] [Medline: [22146289](https://pubmed.ncbi.nlm.nih.gov/22146289/)]
78. Maguire R, Miller M, McCann LA, Lesley T, Kearney N, Meurig S. Managing chemotherapy symptoms via mobile phones. *Nursing Times* 2008;104(22):28-29.
79. Mao Y, Zhang Y, Zhai S. Mobile phone text messaging for pharmaceutical care in a hospital in China. *J Telemed Telecare* 2008 Jan;14(8):410-414. [doi: [10.1258/jtt.2008.080406](https://doi.org/10.1258/jtt.2008.080406)] [Medline: [19047450](https://pubmed.ncbi.nlm.nih.gov/19047450/)]
80. Martínez-Ramos C, Cerdán MT, López RS. Mobile phone-based telemedicine system for the home follow-up of patients undergoing ambulatory surgery. *Telemed J E Health* 2009;15(6):531-537. [doi: [10.1089/tmj.2009.0003](https://doi.org/10.1089/tmj.2009.0003)] [Medline: [19566396](https://pubmed.ncbi.nlm.nih.gov/19566396/)]
81. Mattila E, Lappalainen R, Pärkkä J, Salminen J, Korhonen I. Use of a mobile phone diary for observing weight management and related behaviours. *J Telemed Telecare* 2010;16(5):260-264. [doi: [10.1258/jtt.2009.091103](https://doi.org/10.1258/jtt.2009.091103)] [Medline: [20483880](https://pubmed.ncbi.nlm.nih.gov/20483880/)]
82. McCall K, Keen J, Farrer K, Maguire R, McCann L, Johnston B, et al. Perceptions of the use of a remote monitoring system in patients receiving palliative care at home. *Int J Palliat Nurs* 2008 Sep;14(9):426-431. [Medline: [19060793](https://pubmed.ncbi.nlm.nih.gov/19060793/)]
83. McCann L, Maguire R, Miller M, Kearney N. Patients' perceptions and experiences of using a mobile phone-based advanced symptom management system (ASyMS) to monitor and manage chemotherapy related toxicity. *Eur J Cancer Care (Engl)* 2009 Mar;18(2):156-164. [doi: [10.1111/j.1365-2354.2008.00938.x](https://doi.org/10.1111/j.1365-2354.2008.00938.x)] [Medline: [19267731](https://pubmed.ncbi.nlm.nih.gov/19267731/)]

84. Meltzer EO, Kelley N, Hovell MF. Randomized, cross-over evaluation of mobile phone vs paper diary in subjects with mild to moderate persistent asthma. *Open Respir Med J* 2008 Jan;2:72-79 [FREE Full text] [doi: [10.2174/1874306400802010072](https://doi.org/10.2174/1874306400802010072)] [Medline: [19412327](https://pubmed.ncbi.nlm.nih.gov/19412327/)]
85. Milne RG. Reducing non-attendance at specialist clinics: an evaluation of the effectiveness and cost of patient-focused booking and SMS reminders at a Scottish health board. *Int J Consum Stud* 2010 Aug;34(5):570-580. [doi: [10.1111/j.1470-6431.2010.00903.x](https://doi.org/10.1111/j.1470-6431.2010.00903.x)]
86. Morak J, Schindler K, Goerzer E, Kastner P, Toplak H, Ludvik B, et al. A pilot study of mobile phone-based therapy for obese patients. *J Telemed Telecare* 2008;14(3):147-149. [doi: [10.1258/jtt.2008.003015](https://doi.org/10.1258/jtt.2008.003015)] [Medline: [18430285](https://pubmed.ncbi.nlm.nih.gov/18430285/)]
87. Morris ME, Kathawala Q, Leen TK, Gorenstein EE, Guilak F, Labhard M, et al. Mobile therapy: case study evaluations of a cell phone application for emotional self-awareness. *J Med Internet Res* 2010 Jan;12(2):e10 [FREE Full text] [doi: [10.2196/jmir.1371](https://doi.org/10.2196/jmir.1371)] [Medline: [20439251](https://pubmed.ncbi.nlm.nih.gov/20439251/)]
88. Mosso JL, Gorini A, De La Cerda G, Obrador T, Almazan A, Mosso D, et al. Virtual reality on mobile phones to reduce anxiety in outpatient surgery. *Stud Health Technol Inform* 2009;142:195-200. [Medline: [19377147](https://pubmed.ncbi.nlm.nih.gov/19377147/)]
89. Nakajima R, Nakamura K, Takano T, Seino K, Inose T. Improvements in health by consultations using mobile videophones among participants in a community health promotion programme. *J Telemed Telecare* 2007;13(8):411-415. [doi: [10.1258/135763307783064403](https://doi.org/10.1258/135763307783064403)] [Medline: [18078553](https://pubmed.ncbi.nlm.nih.gov/18078553/)]
90. Nguyen HQ, Gill DP, Wolpin S, Steele BG, Benditt JO. Pilot study of a cell phone-based exercise persistence intervention post-rehabilitation for COPD. *Int J Chron Obstruct Pulmon Dis* 2009 Jan;4:301-313 [FREE Full text] [Medline: [19750190](https://pubmed.ncbi.nlm.nih.gov/19750190/)]
91. Noh JH, Cho YJ, Nam HW, Kim JH, Kim DJ, Yoo HS, et al. Web-based comprehensive information system for self-management of diabetes mellitus. *Diabetes Technol Ther* 2010 May;12(5):333-337. [doi: [10.1089/dia.2009.0122](https://doi.org/10.1089/dia.2009.0122)] [Medline: [20388042](https://pubmed.ncbi.nlm.nih.gov/20388042/)]
92. Pallavicini F, Algeri D, Repetto C, Gorini A, Riva G. Biofeedback, Virtual Reality and Mobile Phones in the Treatment of Generalized Anxiety Disorder (GAD): a Phase-2 Controlled Clinical Trial. *Journal of CyberTherapy & Rehabilitation* 2009;2(4):315-327 [FREE Full text]
93. Patrick K, Raab F, Adams MA, Dillon L, Zabinski M, Rock CL, et al. A text message-based intervention for weight loss: randomized controlled trial. *J Med Internet Res* 2009 Jan;11(1):e1 [FREE Full text] [doi: [10.2196/jmir.1100](https://doi.org/10.2196/jmir.1100)] [Medline: [19141433](https://pubmed.ncbi.nlm.nih.gov/19141433/)]
94. Pérez F, Montón E, Nodal MJ, Viñoles J, Guillen S, Traver V. Evaluation of a mobile health system for supporting postoperative patients following day surgery. *J Telemed Telecare* 2006;12 Suppl 1:41-43. [doi: [10.1258/13576330677978506](https://doi.org/10.1258/13576330677978506)] [Medline: [16884576](https://pubmed.ncbi.nlm.nih.gov/16884576/)]
95. Pinsker M, Schindler K, Hayn D, Kastner P, Riedl M, Ludvik B. Experiences Using Mobile Phones as Patient-Terminal for Telemedical Home Care and Therapy Monitoring of Patients Suffering from Chronic Diseases. *Lecture Notes in Computer Science* 2008;5105:1305-1312. [doi: [10.1007/978-3-540-70540-6_195](https://doi.org/10.1007/978-3-540-70540-6_195)]
96. Pop-Eleches C, Thirumurthy H, Habyarimana JP, Zivin JG, Goldstein MP, de Walque D, et al. Mobile phone technologies improve adherence to antiretroviral treatment in a resource-limited setting: a randomized controlled trial of text message reminders. *AIDS* 2011 Mar 27;25(6):825-834. [doi: [10.1097/QAD.0b013e32834380c1](https://doi.org/10.1097/QAD.0b013e32834380c1)] [Medline: [21252632](https://pubmed.ncbi.nlm.nih.gov/21252632/)]
97. Prabhakaran L, Chee WY, Chua KC, Abisheganaden J, Wong WM. The use of text messaging to improve asthma control: a pilot study using the mobile phone short messaging service (SMS). *J Telemed Telecare* 2010;16(5):286-290. [doi: [10.1258/jtt.2010.090809](https://doi.org/10.1258/jtt.2010.090809)] [Medline: [20576744](https://pubmed.ncbi.nlm.nih.gov/20576744/)]
98. Puccio JA, Belzer M, Olson J, Martinez M, Salata C, Tucker D, et al. The use of cell phone reminder calls for assisting HIV-infected adolescents and young adults to adhere to highly active antiretroviral therapy: a pilot study. *AIDS Patient Care STDS* 2006 Jun;20(6):438-444. [doi: [10.1089/apc.2006.20.438](https://doi.org/10.1089/apc.2006.20.438)] [Medline: [16789857](https://pubmed.ncbi.nlm.nih.gov/16789857/)]
99. Quinn CC, Clough SS, Minor JM, Lender D, Okafor MC, Gruber-Baldini A. WellDoc mobile diabetes management randomized controlled trial: change in clinical and behavioral outcomes and patient and physician satisfaction. *Diabetes Technol Ther* 2008 Jun;10(3):160-168. [doi: [10.1089/dia.2008.0283](https://doi.org/10.1089/dia.2008.0283)] [Medline: [18473689](https://pubmed.ncbi.nlm.nih.gov/18473689/)]
100. Quinn CC, Gruber-Baldini AL, Shardell M, Weed K, Clough SS, Peeples M, et al. Mobile diabetes intervention study: testing a personalized treatment/behavioral communication intervention for blood glucose control. *Contemp Clin Trials* 2009 Jul;30(4):334-346. [doi: [10.1016/j.cct.2009.02.004](https://doi.org/10.1016/j.cct.2009.02.004)] [Medline: [19250979](https://pubmed.ncbi.nlm.nih.gov/19250979/)]
101. Quinn CC, Shardell MD, Terrin ML, Barr EA, Ballew SH, Gruber-Baldini AL. Cluster-randomized trial of a mobile phone personalized behavioral intervention for blood glucose control. *Diabetes Care* 2011 Sep;34(9):1934-1942 [FREE Full text] [doi: [10.2337/dc11-0366](https://doi.org/10.2337/dc11-0366)] [Medline: [21788632](https://pubmed.ncbi.nlm.nih.gov/21788632/)]
102. Rami B, Popow C, Horn W, Waldhoer T, Schober E. Telemedical support to improve glycemic control in adolescents with type 1 diabetes mellitus. *Eur J Pediatr* 2006 Oct;165(10):701-705. [doi: [10.1007/s00431-006-0156-6](https://doi.org/10.1007/s00431-006-0156-6)] [Medline: [16670859](https://pubmed.ncbi.nlm.nih.gov/16670859/)]
103. Reid SC, Kauer SD, Dudgeon P, Sanci LA, Shrier LA, Patton GC. A mobile phone program to track young people's experiences of mood, stress and coping. Development and testing of the mobiletype program. *Soc Psychiatry Psychiatr Epidemiol* 2009 Jun;44(6):501-507. [doi: [10.1007/s00127-008-0455-5](https://doi.org/10.1007/s00127-008-0455-5)] [Medline: [19011723](https://pubmed.ncbi.nlm.nih.gov/19011723/)]
104. Reid SC, Kauer SD, Hearps SJ, Crooke AH, Khor AS, Sanci LA, et al. A mobile phone application for the assessment and management of youth mental health problems in primary care: a randomised controlled trial. *BMC Fam Pract* 2011 Jan;12:131 [FREE Full text] [doi: [10.1186/1471-2296-12-131](https://doi.org/10.1186/1471-2296-12-131)] [Medline: [22123031](https://pubmed.ncbi.nlm.nih.gov/22123031/)]

105. Riley W, Obermayer J, Jean-Mary J. Internet and mobile phone text messaging intervention for college smokers. *J Am Coll Health* 2008;57(2):245-248. [doi: [10.3200/JACH.57.2.245-248](https://doi.org/10.3200/JACH.57.2.245-248)] [Medline: [18809542](https://pubmed.ncbi.nlm.nih.gov/18809542/)]
106. Rizvi SL, Dimeff LA, Skutch J, Carroll D, Linehan MM. A pilot study of the DBT coach: an interactive mobile phone application for individuals with borderline personality disorder and substance use disorder. *Behav Ther* 2011 Dec;42(4):589-600. [doi: [10.1016/j.beth.2011.01.003](https://doi.org/10.1016/j.beth.2011.01.003)] [Medline: [22035988](https://pubmed.ncbi.nlm.nih.gov/22035988/)]
107. Rodgers A, Corbett T, Bramley D, Riddell T, Wills M, Lin RB, et al. Do u smoke after txt? Results of a randomised trial of smoking cessation using mobile phone text messaging. *Tob Control* 2005 Aug;14(4):255-261 [FREE Full text] [doi: [10.1136/tc.2005.011577](https://doi.org/10.1136/tc.2005.011577)] [Medline: [16046689](https://pubmed.ncbi.nlm.nih.gov/16046689/)]
108. Rollo ME, Ash S, Lyons-Wall P, Russell A. Trial of a mobile phone method for recording dietary intake in adults with type 2 diabetes: evaluation and implications for future applications. *J Telemed Telecare* 2011;17(6):318-323. [doi: [10.1258/jtt.2011.100906](https://doi.org/10.1258/jtt.2011.100906)] [Medline: [21844173](https://pubmed.ncbi.nlm.nih.gov/21844173/)]
109. Scherr D, Zweiker R, Kollmann A, Kastner P, Schreier G, Fruhwald FM. Mobile phone-based surveillance of cardiac patients at home. *J Telemed Telecare* 2006;12(5):255-261. [doi: [10.1258/135763306777889046](https://doi.org/10.1258/135763306777889046)] [Medline: [16848939](https://pubmed.ncbi.nlm.nih.gov/16848939/)]
110. Scherr D, Kastner P, Kollmann A, Hallas A, Auer J, Krappinger H, MOBITEL Investigators. Effect of home-based telemonitoring using mobile phone technology on the outcome of heart failure patients after an episode of acute decompensation: randomized controlled trial. *J Med Internet Res* 2009;11(3):e34 [FREE Full text] [doi: [10.2196/jmir.1252](https://doi.org/10.2196/jmir.1252)] [Medline: [19687005](https://pubmed.ncbi.nlm.nih.gov/19687005/)]
111. Schiel R, Kaps A, Bieber G, Kramer G, Seebach H, Hoffmeyer A. Identification of determinants for weight reduction in overweight and obese children and adolescents. *J Telemed Telecare* 2010;16(7):368-373. [doi: [10.1258/jtt.2010.091005](https://doi.org/10.1258/jtt.2010.091005)] [Medline: [20679405](https://pubmed.ncbi.nlm.nih.gov/20679405/)]
112. Seid M, D'Amico EJ, Varni JW, Munafo JK, Britto MT, Kercksmar CM, et al. The in vivo adherence intervention for at risk adolescents with asthma: report of a randomized pilot trial. *J Pediatr Psychol* 2012 May;37(4):390-403. [doi: [10.1093/jpepsy/jsr107](https://doi.org/10.1093/jpepsy/jsr107)] [Medline: [22167121](https://pubmed.ncbi.nlm.nih.gov/22167121/)]
113. Shapiro JR, Bauer S, Hamer RM, Kordy H, Ward D, Bulik CM. Use of text messaging for monitoring sugar-sweetened beverages, physical activity, and screen time in children: a pilot study. *J Nutr Educ Behav* 2008;40(6):385-391 [FREE Full text] [doi: [10.1016/j.jneb.2007.09.014](https://doi.org/10.1016/j.jneb.2007.09.014)] [Medline: [18984496](https://pubmed.ncbi.nlm.nih.gov/18984496/)]
114. Sharma R, Hebbal M, Ankola AV, Murugabupathy V. Mobile-phone text messaging (SMS) for providing oral health education to mothers of preschool children in Belgaum City. *J Telemed Telecare* 2011;17(8):432-436. [doi: [10.1258/jtt.2011.110416](https://doi.org/10.1258/jtt.2011.110416)] [Medline: [22025742](https://pubmed.ncbi.nlm.nih.gov/22025742/)]
115. Spaniel F, Vohlídká P, Hrdlicka J, Kožený J, Novák T, Motlová L, et al. ITAREPS: information technology aided relapse prevention programme in schizophrenia. *Schizophr Res* 2008 Jan;98(1-3):312-317. [doi: [10.1016/j.schres.2007.09.005](https://doi.org/10.1016/j.schres.2007.09.005)] [Medline: [17920245](https://pubmed.ncbi.nlm.nih.gov/17920245/)]
116. Stapleton S, Adams M, Atterton L. A mobile phone as a memory aid for individuals with traumatic brain injury: a preliminary investigation. *Brain Inj* 2007 Apr;21(4):401-411. [doi: [10.1080/02699050701252030](https://doi.org/10.1080/02699050701252030)] [Medline: [17487638](https://pubmed.ncbi.nlm.nih.gov/17487638/)]
117. Sternfeld B, Jiang SF, Picchi T, Chasan-Taber L, Ainsworth B, Quesenberry CP. Evaluation of a cell phone-based physical activity diary. *Med Sci Sports Exerc* 2012 Mar;44(3):487-495. [doi: [10.1249/MSS.0b013e3182325f45](https://doi.org/10.1249/MSS.0b013e3182325f45)] [Medline: [21857369](https://pubmed.ncbi.nlm.nih.gov/21857369/)]
118. Strandbygaard U, Thomsen SF, Backer V. A daily SMS reminder increases adherence to asthma treatment: a three-month follow-up study. *Respir Med* 2010 Feb;104(2):166-171. [doi: [10.1016/j.rmed.2009.10.003](https://doi.org/10.1016/j.rmed.2009.10.003)] [Medline: [19854632](https://pubmed.ncbi.nlm.nih.gov/19854632/)]
119. Stuckey M, Fulkerson R, Read E, Russell-Minda E, Munoz C, Kleinstiver P, et al. Remote monitoring technologies for the prevention of metabolic syndrome: the Diabetes and Technology for Increased Activity (DaTA) study. *J Diabetes Sci Technol* 2011 Jul;5(4):936-944 [FREE Full text] [Medline: [21880237](https://pubmed.ncbi.nlm.nih.gov/21880237/)]
120. Tasker AP, Gibson L, Franklin V, Gregor P, Greene S. What is the frequency of symptomatic mild hypoglycemia in type 1 diabetes in the young?: assessment by novel mobile phone technology and computer-based interviewing. *Pediatr Diabetes* 2007 Feb;8(1):15-20. [doi: [10.1111/j.1399-5448.2006.00220.x](https://doi.org/10.1111/j.1399-5448.2006.00220.x)] [Medline: [17341287](https://pubmed.ncbi.nlm.nih.gov/17341287/)]
121. Treasure J, Macare C, Mentxaka IO, Harrison A. The use of a vodcast to support eating and reduce anxiety in people with eating disorder: A case series. *Eur Eat Disord Rev* 2010;18(6):515-521. [doi: [10.1002/erv.1034](https://doi.org/10.1002/erv.1034)] [Medline: [20669153](https://pubmed.ncbi.nlm.nih.gov/20669153/)]
122. Tsai C, Lee G, Raab F, Norman G, Sohn T, Griswold W, et al. Usability and Feasibility of PmEB: A Mobile Phone Application for Monitoring Real Time Caloric Balance. In: Proceedings of Pervasive Health Conference and Workshops, 2006. 2006 Presented at: Pervasive Health Conference and Workshops; Nov. 29-Dec. 1, 2006; Innsbruck URL: <http://dx.plos.org/10.1371/journal.pone.0044740>
123. Turner J, Larsen M, Tarassenko L, Neil A, Farmer A. Implementation of telehealth support for patients with type 2 diabetes using insulin treatment: an exploratory study. *Inform Prim Care* 2009;17(1):47-53. [Medline: [19490773](https://pubmed.ncbi.nlm.nih.gov/19490773/)]
124. Vidrine DJ, Arduino RC, Gritz ER. Impact of a cell phone intervention on mediating mechanisms of smoking cessation in individuals living with HIV/AIDS. *Nicotine Tob Res* 2006 Dec;8 Suppl 1:S103-S108. [Medline: [17491177](https://pubmed.ncbi.nlm.nih.gov/17491177/)]
125. Vidrine DJ, Marks RM, Arduino RC, Gritz ER. Efficacy of cell phone-delivered smoking cessation counseling for persons living with HIV/AIDS: 3-month outcomes. *Nicotine Tob Res* 2012 Jan;14(1):106-110 [FREE Full text] [doi: [10.1093/ntr/ntr121](https://doi.org/10.1093/ntr/ntr121)] [Medline: [21669958](https://pubmed.ncbi.nlm.nih.gov/21669958/)]
126. Vilella A, Bayas JM, Diaz MT, Guinovart C, Diez C, Simó D, et al. The role of mobile phones in improving vaccination rates in travelers. *Prev Med* 2004 Apr;38(4):503-509. [doi: [10.1016/j.ypmed.2003.12.005](https://doi.org/10.1016/j.ypmed.2003.12.005)] [Medline: [15020186](https://pubmed.ncbi.nlm.nih.gov/15020186/)]

127. Vogel PA, Launes G, Moen EM, Solem S, Hansen B, Håland AT, et al. Videoconference- and cell phone-based cognitive-behavioral therapy of obsessive-compulsive disorder: a case series. *J Anxiety Disord* 2012 Jan;26(1):158-164. [doi: [10.1016/j.janxdis.2011.10.009](https://doi.org/10.1016/j.janxdis.2011.10.009)] [Medline: [22119331](https://pubmed.ncbi.nlm.nih.gov/22119331/)]
128. Wang DH, Kogashiwa M, Ohta S, Kira S. Validity and reliability of a dietary assessment method: the application of a digital camera with a mobile phone card attachment. *J Nutr Sci Vitaminol (Tokyo)* 2002 Dec;48(6):498-504. [Medline: [12775117](https://pubmed.ncbi.nlm.nih.gov/12775117/)]
129. Wangberg SC, Arsand E, Andersson N. Diabetes education via mobile text messaging. *J Telemed Telecare* 2006;12 Suppl 1:55-56. [doi: [10.1258/13576330677978515](https://doi.org/10.1258/13576330677978515)] [Medline: [16884582](https://pubmed.ncbi.nlm.nih.gov/16884582/)]
130. Whittaker R, Dorey E, Bramley D, Bullen C, Denny S, Elley CR, et al. A theory-based video messaging mobile phone intervention for smoking cessation: randomized controlled trial. *J Med Internet Res* 2011 Jan;13(1):e10 [FREE Full text] [doi: [10.2196/jmir.1553](https://doi.org/10.2196/jmir.1553)] [Medline: [21371991](https://pubmed.ncbi.nlm.nih.gov/21371991/)]
131. Whittaker R, Maddison R, McRobbie H, Bullen C, Denny S, Dorey E, et al. A multimedia mobile phone-based youth smoking cessation intervention: findings from content development and piloting studies. *J Med Internet Res* 2008 Jan;10(5):e49 [FREE Full text] [doi: [10.2196/jmir.1007](https://doi.org/10.2196/jmir.1007)] [Medline: [19033148](https://pubmed.ncbi.nlm.nih.gov/19033148/)]
132. Woolford SJ, Clark SJ, Strecher VJ, Resnicow K. Tailored mobile phone text messages as an adjunct to obesity treatment for adolescents. *J Telemed Telecare* 2010;16(8):458-461 [FREE Full text] [doi: [10.1258/jtt.2010.100207](https://doi.org/10.1258/jtt.2010.100207)] [Medline: [20959393](https://pubmed.ncbi.nlm.nih.gov/20959393/)]
133. Zanner R, Wilhelm D, Feussner H, Schneider G. Evaluation of M-AID, a first aid application for mobile phones. *Resuscitation* 2007 Sep;74(3):487-494. [doi: [10.1016/j.resuscitation.2007.02.004](https://doi.org/10.1016/j.resuscitation.2007.02.004)] [Medline: [17452068](https://pubmed.ncbi.nlm.nih.gov/17452068/)]
134. Ahern DK, Kreslake JM, Phalen JM. What is eHealth (6): perspectives on the evolution of eHealth research. *J Med Internet Res* 2006;8(1):e4 [FREE Full text] [doi: [10.2196/jmir.8.1.e4](https://doi.org/10.2196/jmir.8.1.e4)] [Medline: [16585029](https://pubmed.ncbi.nlm.nih.gov/16585029/)]
135. Nilsen W, Kumar S, Shar A, Varoquiers C, Wiley T, Riley WT, et al. Advancing the science of mHealth. *J Health Commun* 2012 May;17 Suppl 1:5-10. [doi: [10.1080/10810730.2012.677394](https://doi.org/10.1080/10810730.2012.677394)] [Medline: [22548593](https://pubmed.ncbi.nlm.nih.gov/22548593/)]

Abbreviations

ACM: Association for Computing Machinery

CINAHL: Cumulative Index to Nursing and Allied Health Literature

MMS: multimedia messaging service

SMS: short message service

Edited by G Eysenbach; submitted 09.11.12; peer-reviewed by Z Hajar, R Whittaker; comments to author 22.01.13; revised version received 05.03.13; accepted 28.03.13; published 21.05.13

Please cite as:

Fiordelli M, Diviani N, Schulz PJ

Mapping mHealth Research: A Decade of Evolution

J Med Internet Res 2013;15(5):e95

URL: <http://www.jmir.org/2013/5/e95/>

doi: [10.2196/jmir.2430](https://doi.org/10.2196/jmir.2430)

PMID: [23697600](https://pubmed.ncbi.nlm.nih.gov/23697600/)

©Maddalena Fiordelli, Nicola Diviani, Peter J. Schulz. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 21.05.2013. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.