

Original Paper

Secure Asynchronous Communication Between Smokers and Tobacco Treatment Specialists: Secondary Analysis of a Web-Assisted Tobacco Intervention in the QUIT-PRIMO and National Dental PBRN Networks

Rajani Shankar Sadasivam^{1*}, PhD; Ariana Kamberi^{1*}, MBA; Kathryn DeLaughter^{2*}, MA; Barrett Phillips^{3*}, MPH; Jessica H Williams^{4*}, MPH, PhD; Sarah L Cutrona^{1*}, MD; Midge N Ray^{4*}, PhD; Gregg H Gilbert^{4*}, DDS, MBA; Thomas K Houston^{1*}, MD, MPH; QUITPRIMO^{4*}; National Dental PBRN Collaborative Group^{4*}

¹University of Massachusetts Medical School, Worcester, MA, United States

²Edith Nourse Rogers Memorial Veterans Hospital, Bedford, MA, United States

³Veterans Affairs Central Western Massachusetts Healthcare System, Leeds, MA, United States

⁴University of Alabama at Birmingham, Birmingham, AL, United States

* all authors contributed equally

Corresponding Author:

Rajani Shankar Sadasivam, PhD

University of Massachusetts Medical School

365 Plantation Street

Worcester, MA, 01605

United States

Phone: 1 508 856 8923

Email: rajani.sadasivam@umassmed.edu

Abstract

Background: Within a web-assisted tobacco intervention, we provided a function for smokers to asynchronously communicate with a trained tobacco treatment specialist (TTS). Previous studies have not attempted to isolate the effect of asynchronous counseling on smoking cessation.

Objective: This study aimed to conduct a semiquantitative analysis of TTS-smoker communication and evaluate its association with smoking cessation.

Methods: We conducted a secondary analysis of data on secure asynchronous communication between trained TTSs and a cohort of smokers during a 6-month period. Smokers were able to select their preferred TTS and message them using a secure web-based form. To evaluate whether the TTS used evidence-based practices, we coded messages using the Motivational Interviewing Self-Evaluation Checklist and Smoking Cessation Counseling (SCC) Scale. We assessed the content of messages initiated by the smokers by creating topical content codes. At 6 months, we assessed the association between smoking cessation and the amount of TTS use and created a multivariable model adjusting for demographic characteristics and smoking characteristics at baseline.

Results: Of the 725 smokers offered asynchronous counseling support, 33.8% (245/725) messaged the TTS at least once. A total of 1082 messages (TTSs: 565; smokers 517) were exchanged between the smokers and TTSs. The majority of motivational interviewing codes were those that supported client strengths (280/517, 54.1%) and promoted engagement (280/517, 54.1%). SCC code analysis showed that the TTS provided assistance to smokers if they were willing to quit (247/517, 47.8%) and helped smokers prepare to quit (206/517, 39.8%) and anticipate barriers (197/517, 38.1%). The majority of smokers' messages discussed motivations to quit (234/565, 41.4%) and current and past treatments (talking about their previous use of nicotine replacement therapy and medications; 201/565, 35.6%). The majority of TTS messages used behavioral strategies (233/517, 45.1%), offered advice on treatments (189/517, 36.5%), and highlighted motivations to quit (171/517, 33.1%). There was no association between the amount of TTS use and cessation. In the multivariable model, after adjusting for gender, age, race, education, readiness at baseline, number of cigarettes smoked per day at baseline, and the selected TTS, smokers messaging the TTS one or two times

had a smoking cessation odds ratio (OR) of 0.8 (95% CI 0.4-1.4), and those that messaged the TTS more than two times had a smoking cessation OR of 1.0 (95% CI 0.4-2.3).

Conclusions: Our study demonstrated the feasibility of using asynchronous counseling to deliver evidence-based counseling. Low participant engagement or a lack of power could be potential explanations for the nonassociation with smoking cessation. Future trials should explore approaches to increase participant engagement and test asynchronous counseling in combination with other approaches for improving the rates of smoking cessation.

(*J Med Internet Res* 2020;22(5):e13289) doi: [10.2196/13289](https://doi.org/10.2196/13289)

KEYWORDS

distance counseling; tobacco cessation; internet-based intervention; smoking cessation

Introduction

Background

Smoking is the number one preventable cause of death [1]. Nearly 1 in 5 adults in the United States, or 42.1 million people, currently smoke cigarettes [2]. Quitting reduces the risk of smoking-related diseases, including cardiovascular diseases [3]. Current guidelines recommend that smokers use behavioral counseling to aid their quitting efforts [4]. However, most smokers do not use counseling in their quit attempts. For instance, despite the wide availability of Quitlines, a telephone-based tobacco counseling that has been established in all 50 states, it has been reported that only 1% to 2% of smokers use Quitline services [5,6]. More innovations are needed to increase access to counseling resources [1].

Providing behavioral counseling by trained tobacco treatment specialists (TTSs) over the internet may help overcome barriers to access. Internet access is almost ubiquitous in the United States, especially with the increasing use of mobile phones [7-9]. Users have also become comfortable communicating over the web (eg, emails, text messages, and on social networks). Web-based counseling is usually delivered synchronously through live chat, where the counselor and smoker are both online and engage via video, audio, or messaging. Synchronous counseling has the benefit of the smokers being able to directly interact with the counselors, and for the counselors to be able to tailor the counseling to the smoker. Previous papers have evaluated synchronous counseling [10-12]. However, synchronous counseling may not be convenient for all smokers as they need to coordinate their schedules with the counselors. Asynchronous counseling that allows the smokers the convenience of communicating with the counselor at any time can be used to augment synchronous counseling. Asynchronous counseling can be achieved using a secure Health Insurance Portability and Accountability Act-compliant form of email or secure messaging. Previous studies have provided functions to communicate asynchronously with experts such as a question and answer email service or a text messaging interface for communicating with a human coach [13-15]. These functions have mostly been provided within the context of a larger intervention. To our knowledge, none of these studies conducted secondary analysis to understand and explore the effect of these counseling efforts.

Objective

Our objective is to conduct a secondary evaluation of asynchronous web-based counseling provided by trained TTSs to smokers participating in the QUIT-PRIMO randomized trial testing a web-assisted tobacco intervention (WATI) [16,17]. As published before [16-18] and further described below, in the QUIT-PRIMO randomized trial, the recruited smokers were randomized to the following three groups: a WATI control (control), the WATI enhanced with proactive, pushed tailored email motivational messaging (messaging), and the WATI with messaging further enhanced with personal secure messaging with a TTS (personalized). Although we found significant differences between those who received the motivational messages (personalized and messaging groups) versus the control (17%; odds ratio [OR] 1.69; 95% CI 1.03-2.8; $P=.04$), we found no differences between the personalized and the messaging group of the study.

Understanding TTS use by the smokers might provide us more insight into why the access to TTS did not result in an increase in smoking cessation in our study and may help in designing more patient-centered asynchronous counseling support in future web-based interventions. In this paper, we add to this literature by conducting a semiquantitative analysis of the TTS-smoker communication and evaluating its association with smoking cessation. We report the frequency of TTS-smoker communication, whether the TTSs were able to incorporate evidence-based practices, such as motivational interviewing, into their counseling. Our analysis also included the communication initiated by the smokers.

Methods

Study Design, Setting, and Sample

We conducted a longitudinal, observational cohort study of secure asynchronous communication between 3 trained TTSs and 245 smokers. The communication occurred during a 6-month period [16]. For the purpose of this study, we only included smokers from the personalized group. These smokers were able to select from among 3 TTSs and initiate the communication by messaging their TTS using a secure web-based form. The TTSs were, then, able to message back using an administrative portal. We implemented the *secure* communication similar to how it is implemented on web-based patient portals. When someone initiated a contact through the website, an email alert was sent to the smoker and the TTS. This email alert did not contain any personal identifiers and the

smoker and the TTS had to log in to the website to view their messages. Our website was available only through an encrypted channel using the hypertext transfer protocol secure.

The approach for selecting and messaging the TTS and coding messages is described in detail below (Choosing a Tobacco Treatment Specialist in the Personalized Group). Our study was approved by the University of Alabama at Birmingham and the University of Massachusetts Medical School Institutional Review Boards.

The QUIT-PRIMO Randomized Trial

Recruitment

Smokers were recruited during June 2010–March 2012. Smoker recruitment was conducted through medical and dental practice patient referrals as well as through Google advertisements [16–19]. Our recruitment process has been described in detail in our previous publications. Briefly, primary care medical practices were recruited from a registered database of internal medicine and family or general practitioners. Dental practices were recruited from state lists of registered dentists and through the Dental Practice–Based Research Network. At these practices, we implemented an electronic referral program that allowed providers (doctors, nurses, or other clinical staff) to recruit smokers to the Decide2Quit.org WATI at the point of care by entering their email addresses into a web-based form. When smokers were electronically referred, they were sent up to 10 email messages encouraging registration over an 8-week period or until the patient registered. To participate on Decide2Quit.org, all smokers completed a web-based consent and registered on the website. To recruit smokers through Google advertisements, 3 advertisements were posted on Google AdWords. Advertisements were linked to searches for keywords related to smoking (eg, smoking, quit smoking, stop smoking, quit, quit smoking tips, and quit smoking programs) and included a link that took participants directly to the Decide2Quit.org home page where they could choose to register as new participants. Each smoker had access to Decide2quit.org for 6 months.

Design of the Randomized Trial

The Decide2Quit.org WATI was designed for all smokers, supporting cessation induction for those not ready to quit and acting as an aid to cessation for those preparing to quit [16–19]. Decide2Quit.org was implemented as an adaptable service with modules that could be engaged based on assigned group. Thus, smokers were allocated to one of three increasingly enhanced versions of Decide2Quit.org. The interventions received by the three randomized groups of smokers are described below.

The Control Group

Smokers randomized to the active control received an interactive, tailored quit smoking website. This module included motivational information tailored to readiness to quit (not thinking of quitting, thinking of quitting, and preparing to quit), and interactive risk, decisional balance, and cessation barrier calculators, and games linking the chemicals in smoking with their other uses (eg, formaldehyde is used in both cigarettes and in embalming). The control also included a library of informational resources about smoking, and sections on seeking

social support and talking to your doctor about quitting. These tools are variations of those in many previously studied evidence-based WATI.

The Messaging Group

Pushed Motivational Email Messages Module Plus Control Module

For this group, we enhanced the control with a pushed motivational email messaging system. Brief motivational email messages were further tailored to an individual smoker's readiness to quit (not ready to quit, thinking about quitting, preparing to quit, and actively quitting). In the first week of registration, 4 email messages were sent to the smoker, followed thereafter by 2 email messages per week. To enhance the personal relevance of messages, our motivational email messaging system included messages written by smokers for other smokers [20].

Personalized Group: Personal Support Module, Messages Module, and Control Module

In addition to the above functions, the personal support module further included an innovative secure messaging portal allowing asynchronous electronic communication between smokers and trained TTSs hired to participate in the intervention team. Further details on how the patient chose and communicated with the TTS are provided below.

Choosing a Tobacco Treatment Specialist in the Personalized Group

During registration over the web, smokers were asked to select between 3 trained TTSs. Smokers were presented the counseling philosophies of the 3 trained TTSs to facilitate this selection. The 3 TTSs created their philosophies from personal experiences and accredited TTS training [21]. Of the 3 TTSs, 2 were trained at the University of Mississippi, Medical Center ACT, while 1 was trained at the University of Massachusetts, Center for Tobacco Treatment Research and Training. Both training programs were accredited by the Council for Tobacco Treatment Training Programs [22,23]. Content of this training consisted of bio, psycho, and social determinants of nicotine dependence, pharmacotherapy, and counseling theory and practice, which included motivational interviewing techniques, treatment strategies, and program and system issues.

Counseling philosophies of the 3 TTSs were as follows:

It is not about the quantity of your life, it is about the quality. Smoking harms nearly every organ in the body and one out of every two smokers will die prematurely from a smoking-related disease. When you think about it, those are not very good odds. Many of my loved ones struggle with tobacco use so I know first-hand how difficult quitting can be. This is why I take a holistic approach to helping you quit tobacco. It is important for you to understand ALL the reasons why you smoke and for you to think about strategies that will work best for you—I want to help you do that. Quitting tobacco may be one of the most difficult things you will do for yourself but is also one of the

most important. Time is ticking...what are you waiting for? [TTS 1]

Being healthy and happy is about learning what's important to you and what you are willing to do to make that happen. Quitting smoking is no different. For some it's as simple as wanting more time with their children or grandchildren, for others perhaps it's about saving money to go on a trip. Either way, small steps now can make a big difference in years to come. For me, I like to help people look at why they want to quit and the reasons they smoke to figure out the best plan for them. Each person is unique and will have their own reasons for quitting smoking. By learning what motivates you and why you do certain things, you are better prepared to make healthy lifestyle changes. This way the skills learned from quitting smoking can be used in other areas of life. [TTS 2]

Quitting smoking can truly be one of the biggest challenges you may face...As a prior smoker, I know first-hand the hold nicotine, a highly addictive drug, can have on one's life. Quitting enabled me to see how much life I was missing. I would like to help you quit. Whether you are ready to quit today or just barely considering quitting, there are many things we can talk about! It is important for each smoker to know their own motivations. One very important motivation is to quit for those around you. Lead by example; kids of parents who quit while they are young are 5 times less likely to smoke themselves, and kids with parents who successfully quit are three times as likely to successfully quit themselves. I can help you to stop being "the smoker"; get rid of the shadow and join the ranks of those proud to say they are "smoke-free." Choosing to quit today can mean a new life tomorrow. [TTS 3]

Smokers were given the option to send a message at the time of registration or proceed to the website and send a message later. After login, smokers were presented with a web form (similar to an email form on a website) that included a textbox for writing the subject and another for writing the message to TTS ([Multimedia Appendix 1](#)). Please note that we did not require smokers to send messages. Once they submitted the message, an email was sent to the participant confirming their submission and indicating that the TTS will respond in 2 business days. The TTSs logged into the administrative portal daily to view any messages they had received. Each message on the administrative portal listed the assigned TTS. Messages were typically replied to by the TTSs within 2 business days. Once the TTS responded, the smokers were sent an email notifying that the TTS had responded to their message. Smokers had to login to the system to read the message. We designed the display of the message to resemble an email inbox. New messages were indicated by the subject in bold font and the smoker had to click on the message subject to read the response.

Coding Process

To code messages and collect codes by categories, we used MAXQDA 10 (VERBI Software) [24-26], a qualitative data analysis software. This software facilitates coding by creating a hierarchical coding dictionary that can be directly mapped to the messages within the software. Authors AK, KL, BP, JW, HC, and SC were all involved in coding the messages. The principal investigator TH guided the process and trained the initial set of coders (BP and AK). BP participated in a training offered by the MAXQDA company on the tool and trained the rest of the coders on the tool. They also created a documentation of codes and coded an initial set of 10 messages under the guidance of TH. Each additional coder was integrated using the following process: They were provided the documentation of codes and asked to code the same 10 messages that had originally been coded by the first two coders (BP and AK). We, then, compared the new and original codes for interrater reliability (IRR). We reviewed the IRR results and compared them with the coding completed by the original coders. On the basis of this comparison and discussion, changes were made to code definitions. Each message was coded by at least 2 coders, and we calculated IRR for each set of 100 messages. After discussion, coder pairs reached agreement on codes used and resolved any discrepancies during the coding process. In the end, 100% agreement was achieved by consensus.

Development of Coding Schema for Analyzing the Tobacco Treatment Specialist Messages

Our coding schema was developed using the Motivational Interviewing Self-Evaluation Checklist and the Smoking Cessation Counseling (SCC) Scale. Motivational interviewing is a directive, patient-centered counseling style for eliciting behavior change by helping people to explore, clarify, and resolve ambivalence [27]. The Motivational Interviewing Self-Evaluation Checklist is a 7-item checklist, used as a self-evaluation tool to improve motivational interviewing skills [28-32]. This checklist is focused on engagement, assessing motivation, addressing ambivalence, promoting internal motivation, eliciting change talk, rolling with resistance, and supporting client strength. The SCC Scale is a 24-item questionnaire based on a 4-level response format. It was originally developed for nurses by the US Department of Health and Human Services to help in assessing, improving, or testing evidence-based methodologies for SCC [33]. The scale has both clinical relevance and research applications and can be used clinically to assess the quality of smoking cessation services [33,34]. We revised the SCC by converting its 4-level response to a 2-level response format, and by using the 17 items that focused on advanced and basic counseling after revising the scale.

We also assessed the content of messages initiated by the smokers. For this purpose, we iteratively developed a topical content coding schema ([Multimedia Appendix 2](#)). To develop this coding system, we first reviewed 20 messages for both TTSs and smokers and developed a preliminary coding schema based on these. We used this schema to code 20 additional messages, and then refined and added new codes if needed. We proceeded in blocks of 20 messages, and code definitions were

revised several times before we finalized our content code category. The content themes included website content; treatment questions: general advice; treatment questions: advice behavioral; treatment questions: advice on treatments and over the counter medications (Rx and OTC); motivations; sociocultural; health; and current or past treatment and feedback from patients.

Data Collection

Data were collected online on the WATI. During registration, we collected demographic and smoking characteristics, and readiness to quit. The TTS and smoker messages were also recorded in the database. At follow-up, the 30-day point prevalence smoking cessation rate was assessed as self-report using the question *Did you smoke any cigarettes during the past 30 days?* [35]. We also assessed on a Likert scale whether communicating with the TTS was helpful to the smoker.

Data Analysis

Our analysis was conducted using STATA version 12 (StataCorp). All analyses were two-sided and alpha error was set at .05. We compared demographic characteristics and smoking behavior between those participants who had sent at least one message to the TTS and those who did not using the chi-square test or the Mantel-Haenszel method (MH odds) for trend. We used descriptive statistics (means or medians) to assess the codes in the TTS messages (Motivational Interviewing Self-Evaluation Checklist and SCC Scale codes). We compared the topical content codes in the TTS messages to the codes in the smoker messages. At the smoker level, we assessed the association between 6-month cessation and the frequency of each code in the TTS messages received by the smoker using separate logistic regression models for each code. For the independent variable for this analysis, we used the number of times the code appeared in messages to best represent the variations in TTS-smoker communication between the smokers. We assessed the association between smoking cessation and

the selected TTS by adjusting for demographic characteristics and smoking characteristics at baseline. Furthermore, we assessed the association between 6-month cessation and the amount of TTS use and created a multivariable logistic model; adjusting for demographic characteristics, smoking characteristics at baseline, and the selected TTS. For this analysis, our dependent variable was the 30-day point prevalence cessation rate as described above. All those with missing follow-up data were coded as continued smoking. The amount of TTS use was the total number of messages sent to the TTS by each smoker. For our analysis, we categorized the amount of TTS use as follows: did not message the TTS, messaged the TTS 1 or 2 times, and messaged the TTS more than 2 times. We implemented a selection model using inverse probability weighting to determine the potential effect of the missing data [36,37]. First, based on covariates available within the dataset, we developed a logistic regression model to predict the amount of missing data. Then, we calculated the inverse probability of not being missing and weighted the main analysis by this probability.

Results

Comparing Characteristics of Smokers Using a Tobacco Treatment Specialist and Those Who Did Not Use a Tobacco Treatment Specialist

Approximately, a third of smokers (245/725, 33.8%) who were offered asynchronous counseling support sent messages at least once to the TTS (see the example of communication between smokers and TTSs in [Multimedia Appendix 3](#)). Those smokers who allowed smoking at home and who had visited a WATI before this study were more likely to use the TTS ($P=.04$ for both; [Table 1](#)). Among those who used the TTS and completed follow-up, 71% (32/45) reported that the communication was very helpful or somewhat helpful. In all, 29% (13/45) indicated that the communication was not very helpful or not at all helpful.

Table 1. Demographics and smoking behavior of participants, comparing those who messaged the tobacco treatment specialist (TTS) at least once and those who did not.

| Patient characteristics | Total smokers included in the study (N=725), n | Messaged the TTS at least once during the 6-month period (n=245), n (%) | Did not message the TTS even once during the 6-month period, (n=480), n (%) | P value |
|--|--|---|---|------------------------|
| Overall | 725 | 245 (33.8) | 480 (66.2) | N/A ^a |
| Patient sex | | | | .28 |
| Female | 469 | 165 (35.2) | 304 (64.8) | N/A |
| Male | 256 | 80 (31.2) | 176 (68.8) | N/A |
| Patient age (years) | | | | .05^b |
| 19-34 | 151 | 37 (24.5) | 114 (75.5) | N/A |
| 35-55 | 361 | 127 (35.2) | 234 (64.8) | N/A |
| 55-64 | 167 | 63 (37.7) | 104 (62.3) | N/A |
| ≥65 | 46 | 18 (39.1) | 28 (60.9) | N/A |
| Patient race | | | | .06 |
| White | 605 | 195 (32.2) | 410 (67.8) | N/A |
| Black or African American | 60 | 28 (46.7) | 32 (53.3) | N/A |
| Other | 22 | 9 (40.9) | 13 (59.1) | N/A |
| Patient education | | | | .09^b |
| Some high school and high school graduate | 245 | 71 (29.0) | 174 (71.0) | N/A |
| Some college | 306 | 109 (35.6) | 197 (64.4) | N/A |
| College graduate or more | 166 | 64 (38.5) | 102 (61.5) | N/A |
| Readiness to quit | | | | .96^b |
| Not thinking of quitting and thinking of quitting | 518 | 176 (34.0) | 342 (66.0) | N/A |
| Set a quit date and already quit | 202 | 69 (34.2) | 133 (65.8) | N/A |
| Allow smoking at home | | | | .04 |
| No | 408 | 125 (30.6) | 283 (69.4) | N/A |
| Yes | 317 | 120 (37.8) | 197 (62.2) | N/A |
| Number of cigarettes smoked per day | | | | .67^b |
| 0-10 | 218 | 72 (33.0) | 146 (67.0) | N/A |
| 11-20 | 355 | 117 (32.9) | 238 (67.1) | N/A |
| >20 | 152 | 56 (36.8) | 96 (63.2) | N/A |
| Visited other smoking cessation websites before the study | | | | .04 |
| No | 565 | 180 (31.8) | 385 (68.2) | N/A |
| Yes | 160 | 65 (40.6) | 95 (59.4) | N/A |
| Quit attempt (1 day or more) in the past 12 months | | | | .70 |
| No | 321 | 106 (33.0) | 215 (67.0) | N/A |
| Yes | 404 | 139 (34.4) | 265 (65.6) | N/A |

^aN/A: not applicable.^bTest for trend analysis.

Message Use Frequency and Volume

A total of 1082 messages (TTSs: 565/1082, 52.2%; smokers: 517/1082, 47.8%) were exchanged between the smokers and TTSs. Among those smokers who messaged at least once, the mean number of messages sent was 2.6 (SD 4.5, median 1, IQR 1). The frequency of messages sent by TTS 1 was 341 (mean 6.1, SD 6.4; median 4, IQR 6), by TTS 2 was 370 (mean 9.7, SD 12.2; median 3, IQR 11), and by TTS 3 was 371 (mean 6.7, SD 8.9; median 2, IQR 7).

Comparison of Tobacco Treatment Specialist Motivational Messaging Codes

Motivational Interviewing Self-Evaluation Checklist

The majority of motivational interviewing codes were those that supported client strengths (280/517, 54.1%) and promoted engagement (280/517, 54.1%; [Table 2](#)). For example, a TTS message highlighting the client strength was “well it sounds like you certainly have good experience with nicotine replacement, which is good because you have an idea of what works and what doesn’t.”

Table 2. Frequency of codes in messages (N=517).

| Scale, category of codes, and codes | Messages, n (%) |
|--|-------------------|
| Motivational Interviewing Scale | |
| Supporting client strengths | 280 (54.1) |
| Explored previous successes | N/A ^a |
| Explored positive qualities | N/A |
| Accentuated <i>any</i> motivation for change | N/A |
| Highlighted any efforts towards change | N/A |
| Used affirmations to highlight strengths, motivation | N/A |
| Engagement | 280 (54.1) |
| Worked to fully understand the problem and the client's perspective before moving toward change | N/A |
| Focused on engagement before change | N/A |
| Used reflective listening to convey empathy and understanding | N/A |
| Used affirmations to build a positive relationship | N/A |
| Eliciting change talk | 110 (21.3) |
| Asked about concerns using open-ended questions or reflective listening | N/A |
| Asked for elaboration about concerns | N/A |
| Explored client values as they relate to change | N/A |
| Selectively responded to change talk with curiosity, interest | N/A |
| Explored pros and cons or decisional balance | N/A |
| Used <i>low-threshold</i> questions | N/A |
| Promoting internal motivation | 106 (20.5) |
| Elicited and respected the client's goals for treatment | N/A |
| Explored values underlying the motivation for change | N/A |
| Supported autonomy in decision making | N/A |
| Assessing motivation | 78 (15.1) |
| Identified a target behavior | N/A |
| Identified stage of change | N/A |
| Used importance, confidence, and readiness ruler | N/A |
| Differentiated between different areas of motivation (eg, substance use vs mental health; treatment vs change) | N/A |
| Addressing ambivalence | 72 (13.9) |
| Normalized ambivalence | N/A |
| Explored ambivalence | N/A |
| Reframed ambivalence | N/A |
| Used a decisional balance | N/A |
| Avoided direct persuasion | N/A |
| Explored pros and cons of change | N/A |
| Rolling with resistance | 48 (9.3) |
| Affirmation (external reframe) | N/A |
| Empathic response | N/A |
| Reflective listening | N/A |
| Providing choice | N/A |
| Nondefensive response | N/A |
| Smoking Cessation Counseling Scale | |

| Scale, category of codes, and codes | Messages, n (%) |
|--|-----------------|
| Basic counseling | |
| If willing to quit, provide assistance | 247 (47.8) |
| Ask if willing to quit | 26 (5.0) |
| If not quitting, help identify barriers | 23 (4.4) |
| Advanced counseling | |
| Helping smokers prepare to quit ^b | 206 (39.8) |
| Helping smokers anticipate barriers ^c | 197 (38.1) |
| Managing relapses ^d | 73 (14.1) |
| Advising smokers to set a quit date | 47 (9.2) |
| Getting social support ^e | 46 (8.9) |

^aN/A: not applicable.

^bHelping smokers prepare to quit codes include identify reasons and benefits for quitting; provide information for follow-up visits with doctor; and recommend over-the-counter nicotine patch, other medications.

^cHelp smokers anticipate barriers codes include anticipate challenges in the beginning; anticipate nicotine withdrawal; alcohol is associated with relapse; and review past quit attempts, what helped, and what led to relapse.

^dManage relapses codes include if relapse occurs, review and learn from experience; if relapse occurs, repeat quit attempts; if relapse occurs, reassess problems; and total abstinence is essential.

^eGetting social support codes include get support from family, friends, and coworkers; and other smokers in house are not helpful.

Smoking Cessation Counseling Scale


SCC Scale code analysis showed that the TTS messages provided assistance to smokers if they were willing to quit (247/517, 47.8%; Table 2). For example, a TTS message that included this code was *When you feel you are ready, I am here to help you*. The TTS messages also included the use of advanced counseling strategies to help the smokers prepare to quit (206/517, 39.8%) and anticipate barriers (197/517, 38.1%), such as *Have you thought about ways other than smoking that can help you deal with stress after you quit? Perhaps this is an*

important first step. It helps to plan as much as possible before you quit to avoid slips.

Topical Content Codes: Comparison of Tobacco Treatment Specialist and Smoker Messages

TTS messages and smoker messages differed in their content (Figure 1). The majority of smoker messages discussed motivations to quit (234/565, 41.4%) and current and past treatments, including their previous use of nicotine replacement therapy and medications (201/565, 35.6%). An example smoker message is as follows:

Figure 1. Proportion of topical content in messages: Smoker and tobacco treatment specialist messages. TTS: tobacco treatment specialist.



I just lost my mom and step father, both from smoking related disease's. I don't want to die like that ,so i will use all help, and support i can get.

The majority of TTS messages used behavioral strategies (233/517, 45.1%), offered advice on treatments (189/517, 36.5%), and highlighted motivations to quit (171/517, 33.1%).

In the following example, a TTS response established the importance of general behavioral strategy:

Most importantly, you should ask your husband to not smoke around you. Your friends and family can be a big help during your quit attempt if you let them know what you need and they support you in quitting. Talk to them about what you will be experiencing during your quit attempt. For example, you may find you have a shorter temper during your quit process. Let your friends or family know this and not to take it personal if you are grumpy.

Association Between the Point Prevalence Cessation Rate Assessed at 6 Months Using a 30-Day Window and the Selected Tobacco Treatment Specialist, the Content of Tobacco Treatment Specialist Motivational Messages, and Amount of Tobacco Treatment Specialist Use

After adjusting for gender, age, race, education, readiness at baseline, and number of cigarettes smoked per day at baseline, compared with TTS 1, the odds of quitting were lower for TTS 2 (OR 0.4; 95% CI 0.1-1.3) and TTS 3 (OR 0.9 95% CI 0.4-2.4),

but not significant. There was no association between the codes in the TTS messages and smoking cessation (Table 3). There was also no association between the total SCC score and smoking cessation (OR 1.02; 95% CI 0.96-1.08).

The amount of TTS use (ie, the total number of messages sent to the TTS by each smoker) ranged from 0 to 42. In all, 66.2% (480/725) of smokers had never used the TTS, 26.1% (189/725) of smokers had used the TTS one or two times, and 7.7% (56/725) of smokers had used it more than two times. There was no association between the amount of TTS use and cessation. In the multivariable model after adjusting for gender, age, race, education, readiness at baseline, number of cigarettes smoked per day at baseline, and the selected TTS, compared with smokers who did not message the TTS, smokers messaging the TTS one or two times had a smoking cessation OR of 0.8 (95% CI 0.4-1.4), and those that messaged the TTS more than two times had a smoking cessation OR of 1.0 (95% CI 0.4-2.3; $P=.50$ for trend across categories). In the inverse probability weighted model, the smokers messaging the TTS one or two times had a smoking cessation OR of 0.7 (95% CI 0.4-1.3), and those that messaged the TTS more than two times had a smoking cessation OR of 0.7 (95% CI 0.3-1.7; $P=.20$ for trend across categories).

Table 3. Association between smoking cessation and the frequency of codes.

| Scale, category of codes, and codes | Odds ratio (95% CI) using separate logistic regression models for each code |
|--|---|
| Motivational Interviewing Scale | |
| Supporting client strengths | 1.12 (0.98-1.32) |
| Explored previous successes | N/A ^a |
| Explored positive qualities | N/A |
| Accentuated <i>any</i> motivation for change | N/A |
| Highlighted any efforts towards change | N/A |
| Used affirmations to highlight strengths, motivation | N/A |
| Engagement | 1.06 (0.88-1.27) |
| Worked to fully understand the problem and the client's perspective before moving toward change | N/A |
| Focused on engagement before change | N/A |
| Used reflective listening to convey empathy and understanding | N/A |
| Used affirmations to build a positive relationship | N/A |
| Eliciting change talk | 1.10 (0.74-1.62) |
| Asked about concerns using open-ended questions or reflective listening | N/A |
| Asked for elaboration about concerns | N/A |
| Explored client values as they relate to change | N/A |
| Selectively responded to change talk with curiosity, interest | N/A |
| Explored pros and cons or decisional balance | N/A |
| Used <i>low-threshold</i> questions | N/A |
| Promoting internal motivation | 1.20 (0.74-1.82) |
| Elicited and respected the client's goals for treatment | N/A |
| Explored values underlying the motivation for change | N/A |
| Supported autonomy in decision making | N/A |
| Assessing motivation | 1.05 (0.52-2.09) |
| Identified a target behavior | N/A |
| Identified stage of change | N/A |
| Used importance, confidence, and readiness ruler | N/A |
| Differentiated between different areas of motivation (eg, substance use vs mental health; treatment vs change) | N/A |
| Addressing ambivalence | 0.94 (0.55-1.63) |
| Normalized ambivalence | N/A |
| Explored ambivalence | N/A |
| Reframed ambivalence | N/A |
| Used a decisional balance | N/A |
| Avoided direct persuasion | N/A |
| Explored pros and cons of change | N/A |
| Rolling with resistance | 0.82 (0.35-1.92) |
| Affirmation (external reframe) | N/A |
| Empathic response | N/A |
| Reflective listening | N/A |
| Providing choice | N/A |

| Scale, category of codes, and codes | Odds ratio (95% CI) using separate logistic regression models for each code |
|--|---|
| Nondefensive response | N/A |
| Smoking Cessation Counseling Scale | |
| Basic counseling | |
| If willing to quit, provide assistance | 1.0 (0.69-1.42) |
| Ask if willing to quit | 0.72 (0.18-2.97) |
| If not quitting, help identify barriers | 1.44 (0.60-3.41) |
| Advanced counseling | |
| Helping smokers prepare to quit ^b | 1.0 (0.82-1.21) |
| Helping smokers anticipate barriers ^c | 1.07 (0.88-1.31) |
| Managing relapses ^d | 0.88 (0.54-1.45) |
| Advising smokers to set a quit date | 1.24 (0.66-2.35) |
| Getting social support ^e | 1.04 (0.6-1.80) |

^aN/A: not applicable.

^bHelping smokers prepare to quit codes include identify reasons and benefits for quitting; provide information for follow-up visits with doctor; and recommend over-the-counter nicotine patch, other medications.

^cHelp smokers anticipate barriers codes include anticipate challenges in the beginning; anticipate nicotine withdrawal; alcohol is associated with relapse; and review past quit attempts, what helped, and what led to relapse.

^dManage relapses codes include if relapse occurs, review and learn from experience; if relapse occurs, repeat quit attempts; if relapse occurs, reassess problems; and total abstinence is essential.

^eGetting social support codes include get support from family, friends, and coworkers; and other smokers in house are not helpful.

Discussion

Principal Findings

Our discussion focuses on three topics: (1) approximately, one-third of the smokers who were offered an opportunity to asynchronously communicate with the TTS over the web chose to use the service, suggesting some level of interest in the service; (2) the coding analysis demonstrated that the TTSs were able to deliver best practice-based counseling on this asynchronous medium; and (3) the finding of no association between the amount of TTS use and smoking cessation has important implications.

Previous interventions with web-based counseling for a variety of behavioral and mental health conditions have had inconsistent results, with negative studies being a common finding. However, as these studies did not explore the message content, we cannot say whether these interventions delivered counseling that included evidence-based communication patterns. Thus, one goal was to understand the content of the messages. In our previous publication, we have qualitatively analyzed the communication between the smoker and the TTS and identified seven basic themes in the communication—talk about the process of quitting, barriers to quitting, reasons to quit, quit history, support and strategies for quitting, quitting with medication, and quit progress [15]. This paper adds to this evaluation by conducting a semiquantitative coding analysis and evaluating the association between the codes and the number of messages and smoking cessation. We also evaluated whether

the content and volume of messages were related to the cessation outcomes at 6 months.

As noted, a third of our participants chose to message the TTS without actively prompting them. Please note that we did not require the smoker to message the TTS. In a study with smokers from Arkansas Children's Hospital, 21% of smokers participated following a fax referral [38]. Fax referral is a process in which smokers are referred to a quit line by simply faxing a referral form to the quit line. Quitline TTS, then, proactively calls the smoker to setup the counseling sessions. In an observational study of faxed referrals to the Ohio Tobacco Quit Line, 23% (n=1616) of smokers were able to be enrolled out of a total of 6951 faxed referrals [39]. These results may suggest that there is some merit to our assumption that asynchronous counseling offers a lower barrier to access, acknowledging that the participants in our study may be more motivated to quit than the general population of smokers, as indicated by their registration on WATI. Furthermore, among those who communicated with the TTS, the majority (32/45, 71%) indicated the communication was helpful.

We used the Motivational Interviewing Self-Evaluation Checklist and the SCC Scale for developing our codes to evaluate the TTS messaging. As noted above, the Motivational Interviewing Self-Evaluation Checklist was originally developed as a self-evaluation tool to improve motivational interviewing skills [28-32], while the SCC Scale was originally developed to help in assessing, improving, or testing evidence-based methodologies for SCC [33]. A contribution of our work is adapting these to code messages between TTSs and smokers

on the web. Some of the messages included motivational interviewing and other behavioral strategies. The techniques used included supporting client strengths and promoting engagement, among others. TTSs were able to include messaging to help smokers prepare to quit and anticipate barriers.

In our content analysis of the smokers' messages, we found that smokers' messages most often discussed motivations to quit and current and past treatments used for smoking cessation, including their previous use of nicotine replacement therapy and medications. This suggested that smokers were able to express their motivations to quit, as well as the challenges they face while quitting. In addition, smokers were also able to express what worked or did not work in their past treatment. This ability of the smokers to include these concepts in their messages is important as it allows the TTSs to tailor their messaging to the smokers. This was evident in our analysis of the TTS messaging, during which TTSs were able to incorporate behavioral strategies, offer advice on treatments, and highlight motivations to quit.

We did not find any association between smoking cessation and the codes in the TTS messages, the selected TTS, and the amount of TTS use. As noted above, only a third of our participants chose to message the TTS and even among these participants, the mean number of messages sent was 2.6. This may have contributed to our inability to detect an effect. The tobacco treatment guidelines recommend four or more sessions to increase the effectiveness of counseling [1]. A previous meta-analysis has also indicated that the session length is an important factor in the effectiveness of counseling [1]. For the light-touch, low-intensity approach of asynchronous counseling, the number of interactions may need to be higher. In future work, we plan to explore approaches to increase engagement, both in terms of overall use of asynchronous counseling and the number of interactions between TTSs and smokers. First, accessing the TTS by logging into the web interface may be a barrier. Future studies could test whether text messaging and mobile phone-based access further reduce barriers to access. Second, users may need tailored and proactive messaging to continually motivate and encourage them to participate in asynchronous counseling. As noted, our study identified certain characteristics that were significantly associated with participation; compared with those who did not message the TTS, smokers who messaged the TTS were those who allowed smoking at home and who had visited a WATI before this study.

These characteristics could potentially be used to tailor messages to motivate the smokers to engage in asynchronous counseling with the TTS.

Future trials should test asynchronous counseling in combination with synchronous counseling, as asynchronous counseling may be most beneficial when used as an augmentation to synchronous counseling. For example, the asynchronous counseling could be used in between synchronous counseling sessions to continually support the smoker. The timing (whether the synchronous counseling session precedes or follows the start of the asynchronous counseling) and the number of synchronous counseling sessions need to be tested. Furthermore, it is difficult to find an overall impact of TTS use because the content of the messages varied considerably.

Limitations

This was an observational study. Furthermore, not all smokers who participated in the larger randomized clinical trials participated in this study. Although this circumstance helped identify those who are likely to use the service, this may have caused some selection bias. The number of interactions between the TTSs and the smokers were also low. As this was a secondary, exploratory analysis, our study had limited power. Comparing TTS use=0 with TTS use=top quartile, we found a difference in cessation of 3% (top quartile: 9/56, 16%; 0 use: 62/480, 12.9%). With the sample available, we were powered to detect a difference of 11%.

Conclusions

Asynchronous counseling may be able to augment synchronous counseling, providing smokers the ability to connect with counselors at their convenience. We did not find an association between asynchronous counseling and smoking cessation. There are several potential explanations for this finding, including a lack of intervention engagement by participants, a lack of intervention fidelity (the counselors did not deliver evidence-based support), and a lack of power. We have now demonstrated that evidence-based cessation counseling did occur. Our discussion outlines several approaches to potentially increasing the effectiveness of asynchronous counseling, including a discussion of approaches to increase participant engagement, and testing the asynchronous counseling in combination with synchronous counseling. Future studies are needed to address the best ways to use asynchronous counseling for helping smokers.

Acknowledgments

Funding for this study was received from the National Institutes of Health grant nos. R01-CA-12909, U01-DE-16746, U01-DE-16747, U19-DE-22516, and U19-DE-28717. TH is also supported by the VA eHealth Quality Enhancement Research Initiative that he directs. RS is funded by a National Cancer Institute Career Development Award (K07-CA-172677). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the US government.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Decide2Quit website.

[DOC File , 1413 KB-Multimedia Appendix 1]

Multimedia Appendix 2

Topical content coding schema.

[DOC File , 150 KB-Multimedia Appendix 2]

Multimedia Appendix 3

Email communications between smokers and TTS.

[DOC File , 78 KB-Multimedia Appendix 3]

References

1. Fiore MC, Jaén CR, Baker TB, Bailey WC, Benowitz NL, Curry SJ, et al. The Tobacco Dependence Program (TDP). Rockville, MD: US Department of Health and Human Services. Public Health Service; 2008 May. Clinical Practice Guideline. Treating Tobacco Use and Dependence: 2008 Update URL: <http://www.tobaccocontrol.org/clientuploads/documents/Consumer%20Materials/Clinicians%20Systems%20Mat/2008-Guidelines.pdf> [accessed 2020-02-12]
2. Centers for Disease Control and Prevention. 2014 Nov 28. Current Cigarette Smoking Among Adults — United States, 2005–2013 URL: <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6347a4.htm> [accessed 2017-12-05]
3. Centers for Disease Control and Prevention. Quitting Smoking URL: http://www.cdc.gov/tobacco/data_statistics/fact_sheets/cessation/quitting/index.htm [accessed 2017-12-05]
4. Fiore MC. Treating tobacco use and dependence: an introduction to the US Public Health Service Clinical Practice Guideline. *Respir Care* 2000 Oct;45(10):1196-1199. [Medline: [11203101](#)]
5. Sheffer C, Brackman S, Lercara C, Cottoms N, Olson M, Panissidi L, et al. When Free Is Not for Me: confronting the barriers to use of free quitline telephone counseling for tobacco dependence. *Int J Environ Res Public Health* 2015 Dec 22;13(1):ijerph13010015 [FREE Full text] [doi: [10.3390/ijerph13010015](#)] [Medline: [26703662](#)]
6. Johnston LD, Miech RA, O'Malley PM, Bachman JG, Schulenberg JE, Patrick ME. ERIC - Education Resources Information Center. Monitoring the Future National Survey Results on Drug Use, 1975-2017: Overview, Key Findings on Adolescent Drug Use URL: <https://eric.ed.gov/?id=ED589762> [accessed 2019-07-10]
7. Pew Research Center. 2019 Jun 12. Internet/Broadband Fact Sheet: Internet Use Over Time URL: <http://www.pewinternet.org/data-trend/internet-use/internet-use-over-time/> [accessed 2019-08-10]
8. Pew Research Center.: Pew Research Center; 2015 Jun 10. Broadband vs Dial-up Adoption Over Time URL: <https://www.pewresearch.org/internet/chart/broadband-vs-dial-up-adoption-over-time/> [accessed 2019-08-12]
9. Smith A. Pew Research Center. 2015 Apr 1. US Smartphone Use in 2015 URL: <http://www.pewinternet.org/2015/04/01/us-smartphone-use-in-2015/> [accessed 2019-06-12]
10. Friederichs SA, Oenema A, Bolman C, Guyaux J, van Keulen HM, Lechner L. Motivational interviewing in a web-based physical activity intervention: questions and reflections. *Health Promot Int* 2015 Sep;30(3):803-815. [doi: [10.1093/heapro/dat069](#)] [Medline: [24101160](#)]
11. Ramo DE, Thrul J, Chavez K, Delucchi KL, Prochaska JJ. Feasibility and quit rates of the tobacco status project: a Facebook smoking cessation intervention for young adults. *J Med Internet Res* 2015 Dec 31;17(12):e291 [FREE Full text] [doi: [10.2196/jmir.5209](#)] [Medline: [26721211](#)]
12. Schaub MP, Wenger A, Berg O, Beck T, Stark L, Buehler E, et al. A web-based self-help intervention with and without chat counseling to reduce cannabis use in problematic cannabis users: three-arm randomized controlled trial. *J Med Internet Res* 2015 Oct 13;17(10):e232 [FREE Full text] [doi: [10.2196/jmir.4860](#)] [Medline: [26462848](#)]
13. McKay HG, Danaher BG, Seeley JR, Lichtenstein E, Gau JM. Comparing two web-based smoking cessation programs: randomized controlled trial. *J Med Internet Res* 2008 Nov 18;10(5):e40 [FREE Full text] [doi: [10.2196/jmir.993](#)] [Medline: [19017582](#)]
14. Patrick H, Fujii CA, Glaser DB, Utley DS, Marler JD. A comprehensive digital program for smoking cessation: assessing feasibility in a single-group cohort study. *JMIR Mhealth Uhealth* 2018 Dec 18;6(12):e11708 [FREE Full text] [doi: [10.2196/11708](#)] [Medline: [30563807](#)]
15. Williams JH, DeLaughter K, Volkman JE, Sadasivam RS, Ray MN, Gilbert GH, QUIT-PRIMONational Dental PBRN Collaborative Groups. Exploring Online Asynchronous Counseling With Tobacco Treatment Specialists in the QUIT-PRIMO and National Dental PBRN HI-QUIT Studies: Who Uses It and What Do They Say? *Am J Health Promot* 2018 Jun;32(5):1170-1177 [FREE Full text] [doi: [10.1177/0890117116670972](#)] [Medline: [29848011](#)]
16. Houston TK, Sadasivam RS, Ford DE, Richman J, Ray MN, Allison JJ. The QUIT-PRIMO provider-patient Internet-delivered smoking cessation referral intervention: a cluster-randomized comparative effectiveness trial: study protocol. *Implement Sci* 2010 Nov 17;5:87 [FREE Full text] [doi: [10.1186/1748-5908-5-87](#)] [Medline: [21080972](#)]

17. Houston TK, Sadasivam RS, Allison JJ, Ash AS, Ray MN, English TM, et al. Evaluating the QUIT-PRIMO clinical practice ePortal to increase smoker engagement with online cessation interventions: a national hybrid type 2 implementation study. *Implement Sci* 2015 Nov 2;10:154 [FREE Full text] [doi: [10.1186/s13012-015-0336-8](https://doi.org/10.1186/s13012-015-0336-8)] [Medline: [26525410](https://pubmed.ncbi.nlm.nih.gov/26525410/)]
18. Ray MN, Funkhouser E, Williams JH, Sadasivam RS, Gilbert GH, Coley HL, National Dental PBRN Collaborative Group. Smoking-cessation e-referrals: a national dental practice-based research network randomized controlled trial. *Am J Prev Med* 2014 Feb;46(2):158-165 [FREE Full text] [doi: [10.1016/j.amepre.2013.10.018](https://doi.org/10.1016/j.amepre.2013.10.018)] [Medline: [24439349](https://pubmed.ncbi.nlm.nih.gov/24439349/)]
19. Sadasivam RS, Kinney RL, Delaughter K, Rao SR, Williams JH, Coley HL, National Dental PBRN Group, QUIT-PRIMO Collaborative Group. Who participates in web-assisted tobacco interventions? The QUIT-PRIMO and National Dental Practice-Based Research Network Hi-Quit studies. *J Med Internet Res* 2013 May 1;15(5):e77 [FREE Full text] [doi: [10.2196/jmir.2385](https://doi.org/10.2196/jmir.2385)] [Medline: [23635417](https://pubmed.ncbi.nlm.nih.gov/23635417/)]
20. Coley HL, Sadasivam RS, Williams JH, Volkman JE, Schoenberger Y, Kohler CL, National Dental PBRNQUITPRIMO Collaborative Group. Crowdsourced peer- versus expert-written smoking-cessation messages. *Am J Prev Med* 2013 Nov;45(5):543-550 [FREE Full text] [doi: [10.1016/j.amepre.2013.07.004](https://doi.org/10.1016/j.amepre.2013.07.004)] [Medline: [24139766](https://pubmed.ncbi.nlm.nih.gov/24139766/)]
21. Center for Tobacco Treatment Research & Training. Tobacco Training Programs at Our Center URL: <http://www.umassmed.edu/tobacco/training/> [accessed 2020-02-12]
22. National Center for Education Statistics (NCES). Percentage of the population 3 to 34 years old enrolled in school, by age group: Selected years, 1940 through 2008 URL: http://nces.ed.gov/programs/digest/d09/tables/dt09_007.asp [accessed 2019-08-15]
23. DiFranza JR, Rigotti NA, McNeill AD, Ockene JK, Savageau JA, St Cyr D, et al. Initial symptoms of nicotine dependence in adolescents. *Tob Control* 2000 Sep;9(3):313-319 [FREE Full text] [doi: [10.1136/tc.9.3.313](https://doi.org/10.1136/tc.9.3.313)] [Medline: [10982576](https://pubmed.ncbi.nlm.nih.gov/10982576/)]
24. MAXQDA. URL: <https://www.maxqda.com/> [accessed 2019-09-12]
25. Karim F, Johansson E, Diwan VK, Kulane A. Community perceptions of tuberculosis: A qualitative exploration from a gender perspective. *Public Health* 2011 Feb;125(2):84-89. [doi: [10.1016/j.puhe.2010.10.005](https://doi.org/10.1016/j.puhe.2010.10.005)] [Medline: [21288542](https://pubmed.ncbi.nlm.nih.gov/21288542/)]
26. Schmidt-Weitmann S, Bösch S, Weidt S, Brockes C. Psychological online consultation service attracts men: a new approach to close the gap between being ill and being treated. *Telemed J E Health* 2015 Jan;21(1):48-54. [doi: [10.1089/tmj.2013.0370](https://doi.org/10.1089/tmj.2013.0370)] [Medline: [25459521](https://pubmed.ncbi.nlm.nih.gov/25459521/)]
27. Rollnick S, Miller WR. What is motivational interviewing? *Behav Cogn Psychother* 1995 Oct;23(4):325-334. [doi: [10.1017/s135246580001643x](https://doi.org/10.1017/s135246580001643x)]
28. Possidente CJ, Bucci KK, McClain WJ. Motivational interviewing: a tool to improve medication adherence? *Am J Health Syst Pharm* 2005 Jun 15;62(12):1311-1314. [doi: [10.1093/ajhp/62.12.1311](https://doi.org/10.1093/ajhp/62.12.1311)] [Medline: [15947131](https://pubmed.ncbi.nlm.nih.gov/15947131/)]
29. Brodie DA, Inoue A, Shaw DG. Motivational interviewing to change quality of life for people with chronic heart failure: a randomised controlled trial. *Int J Nurs Stud* 2008 Apr;45(4):489-500. [doi: [10.1016/j.ijnurstu.2006.11.009](https://doi.org/10.1016/j.ijnurstu.2006.11.009)] [Medline: [17258218](https://pubmed.ncbi.nlm.nih.gov/17258218/)]
30. Erol S, Erdogan S. Application of a stage based motivational interviewing approach to adolescent smoking cessation: the Transtheoretical Model-based study. *Patient Educ Couns* 2008 Jul;72(1):42-48. [doi: [10.1016/j.pec.2008.01.011](https://doi.org/10.1016/j.pec.2008.01.011)] [Medline: [18304775](https://pubmed.ncbi.nlm.nih.gov/18304775/)]
31. van Eijk-Hustings YJL, Daemen L, Schaper NC, Vrijhoef HJ. Implementation of Motivational Interviewing in a diabetes care management initiative in The Netherlands. *Patient Educ Couns* 2011 Jul;84(1):10-15. [doi: [10.1016/j.pec.2010.06.016](https://doi.org/10.1016/j.pec.2010.06.016)] [Medline: [20638812](https://pubmed.ncbi.nlm.nih.gov/20638812/)]
32. TrainingTeams. Motivational Interviewing Self-Evaluation Checklist URL: http://www.trainingteams.org/Document/MI_SelfEvaluationChecklist.pdf [accessed 2020-02-10]
33. Newhouse RP, Himmelfarb CD, Liang Y. Psychometric testing of the smoking cessation counseling scale. *J Nurs Scholarsh* 2011 Dec;43(4):405-411. [doi: [10.1111/j.1547-5069.2011.01420.x](https://doi.org/10.1111/j.1547-5069.2011.01420.x)] [Medline: [22018103](https://pubmed.ncbi.nlm.nih.gov/22018103/)]
34. Chan SS, Sarna L, Wong DC, Lam T. Nurses' tobacco-related knowledge, attitudes, and practice in four major cities in China. *J Nurs Scholarsh* 2007;39(1):46-53. [doi: [10.1111/j.1547-5069.2007.00142.x](https://doi.org/10.1111/j.1547-5069.2007.00142.x)] [Medline: [17393965](https://pubmed.ncbi.nlm.nih.gov/17393965/)]
35. Hughes JR, Keely JP, Niaura RS, Ossip-Klein DJ, Richmond RL, Swan GE. Measures of abstinence in clinical trials: issues and recommendations. *Nicotine Tob Res* 2003 Feb;5(1):13-25. [Medline: [12745503](https://pubmed.ncbi.nlm.nih.gov/12745503/)]
36. Smolkowski K, Danaher BG, Seeley JR, Kosty DB, Severson HH. Modeling missing binary outcome data in a successful web-based smokeless tobacco cessation program. *Addiction* 2010 Jun;105(6):1005-1015 [FREE Full text] [doi: [10.1111/j.1360-0443.2009.02896.x](https://doi.org/10.1111/j.1360-0443.2009.02896.x)] [Medline: [20148782](https://pubmed.ncbi.nlm.nih.gov/20148782/)]
37. Seaman SR, White IR. Review of inverse probability weighting for dealing with missing data. *Stat Methods Med Res* 2013 Jun;22(3):278-295. [doi: [10.1177/0962280210395740](https://doi.org/10.1177/0962280210395740)] [Medline: [21220355](https://pubmed.ncbi.nlm.nih.gov/21220355/)]
38. Sisterhen LL, Sheffer CE, Bursac Z, Fischer EP. Rates of enrollment in smoking cessation services following fax referrals from a children's hospital. *Acad Pediatr* 2010;10(3):200-204. [doi: [10.1016/j.acap.2010.03.001](https://doi.org/10.1016/j.acap.2010.03.001)] [Medline: [20452567](https://pubmed.ncbi.nlm.nih.gov/20452567/)]
39. Willett JG, Hood NE, Burns EK, Swetlick JL, Wilson SM, Lang DA, et al. Clinical faxed referrals to a tobacco quitline: reach, enrollment, and participant characteristics. *Am J Prev Med* 2009 Apr;36(4):337-340. [doi: [10.1016/j.amepre.2008.12.004](https://doi.org/10.1016/j.amepre.2008.12.004)] [Medline: [19201150](https://pubmed.ncbi.nlm.nih.gov/19201150/)]

Abbreviations

IRR: interrater reliability
OR: odds ratio
SCC: smoking cessation counseling
TTS: tobacco treatment specialist
WATI: web-assisted tobacco intervention

Edited by G Eysenbach; submitted 04.01.19; peer-reviewed by R Paz Castro, J McClure, MS Aslam; comments to author 01.03.19; revised version received 24.05.19; accepted 28.01.20; published 06.05.20

Please cite as:

Sadasivam RS, Kamberi A, DeLaughter K, Phillips B, Williams JH, Cutrona SL, Ray MN, Gilbert GH, Houston TK, QUITPRIMO, National Dental PBRN Collaborative Group

Secure Asynchronous Communication Between Smokers and Tobacco Treatment Specialists: Secondary Analysis of a Web-Assisted Tobacco Intervention in the QUIT-PRIMO and National Dental PBRN Networks

J Med Internet Res 2020;22(5):e13289

URL: <https://www.jmir.org/2020/5/e13289>

doi: [10.2196/13289](https://doi.org/10.2196/13289)

PMID: [32374266](https://pubmed.ncbi.nlm.nih.gov/32374266/)

©Rajani Shankar Sadasivam, Ariana Kamberi, Kathryn DeLaughter, Barrett Phillips, Jessica H Williams, Sarah L Cutrona, Midge N Ray, Gregg H Gilbert, Thomas K Houston, QUITPRIMO, National Dental PBRN Collaborative Group. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 06.05.2020. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.