

Corrigenda and Addenda

Correction: Culturally Adapted Guided Internet-Based Cognitive Behavioral Therapy for Hong Kong People With Depressive Symptoms: Randomized Controlled Trial

Jia-Yan Pan¹, PhD; Jonas Rafi², PhD

¹Department of Social Work, Hong Kong Baptist University, Hong Kong, China (Hong Kong)

²Department of Psychology, Stockholm University, Stockholm, Sweden

Corresponding Author:

Jia-Yan Pan, PhD

Department of Social Work

Hong Kong Baptist University

Room AAB1026, 10/F, Academic and Administration Building

15 Baptist University Road, Baptist University Road Campus, Kowloon Tong, KLN

Hong Kong

China (Hong Kong)

Phone: 852 34116415

Email: jiayan@hkbu.edu.hk

Related Article:

Correction of: <https://www.jmir.org/2025/1/e64303>

(*J Med Internet Res* 2026;28:e88495) doi: [10.2196/88495](https://doi.org/10.2196/88495)

In “Culturally Adapted Guided Internet-Based Cognitive Behavioral Therapy for Hong Kong People With Depressive Symptoms: Randomized Controlled Trial”[1], the authors noted some errors.

One sentence has been changed. In the “Measurements” section, the third sentence in the third paragraph originally appeared as follows:

Item scores were summed for the positive and negative subscales, with higher total scores indicating more positive and negative automatic thoughts, respectively.

This sentence now reads:

Item scores were averaged for the positive and negative subscales, with higher total scores indicating more positive and negative automatic thoughts, respectively.

The originally published Tables 2 and 4 have been revised due to typos. The new **Table 2** and **Table 4** that are replacing the original published versions are shown below (with changes marked in italics).

Table 2. Comparison of treatment effects between active and WLC^a groups at post-treatment (intent-to-treat analysis).

Outcome	Unadjusted mean (SD)			Adjusted difference (95% CI)				Between group effect size <i>d</i> (95% CI)		Within group effect size (95% CI)		
	Web	App	WL	Web vs WL	<i>P</i> value	App vs WL	<i>P</i> value	Web vs WL	App vs WL	Web	App	WLC
BDI-II ^b	9.40 (8.90)	9.43 (8.45)	18.02 (9.02)	-8.24 (-10.52 to -5.96)	<.001	-9.38 (-11.65 to -7.10)	<.001	1.07 (0.81 to 1.34)	1.15 (0.88 to 1.43)	1.48 (1.18 to 1.78)	1.64 (1.33 to 1.94)	0.45 (0.2 to 0.71)
PHQ-9 ^c	5.59 (4.43)	5.24 (3.65)	8.99 (4.58)	-3.07 (-4.46 to -1.67)	<.001	-4.50 (-5.89 to -3.12)	<.001	0.78 (0.52 to 1.04)	0.95 (0.63 to 1.27)	1.11 (0.81 to 1.41)	1.69 (1.36 to 2.02)	0.46 (0.19 to 0.73)
GHQ-12 ^d	1.60 (3.06)	1.53 (2.77)	5.35 (3.85)	-3.65 (-4.75 to -2.54)	<.001	-4.49 (-5.6 to -3.38)	<.001	1.05 (0.77 to 1.32)	1.11 (0.83 to 1.39)	1.64 (1.34 to 1.95)	2.21 (1.87 to 2.55)	0.64 (0.38 to 0.9)
BAI ^e	8.89 (7.89)	8.49 (6.69)	11.09 (7.49)	-2.47 (-4.33 to -0.62)	.009	-3.32 (-5.17 to -1.47)	<.001	0.37 (0.11 to 0.63)	0.54 (0.29 to 0.8)	0.64 (0.37 to 0.91)	0.88 (0.59 to 1.15)	0.37 (0.11 to 0.62)
Positive automatic thoughts	2.40 (0.71)	2.33 (0.67)	1.98 (0.58)	0.38 (0.22 to 0.54)	<.001	0.40 (0.24 to 0.55)	<.001	-0.63 (-0.9 to -0.37)	-0.61 (-0.9 to -0.32)	-0.88 (-1.16 to -0.6)	-0.93 (-1.22 to -0.65)	-0.30 (-0.55 to 0.04)
Negative automatic thoughts	1.85 (0.78)	1.89 (0.63)	2.29 (0.80)	-0.43 (-0.63 to -0.22)	<.001	-0.48 (-0.68 to -0.27)	<.001	0.63 (0.37 to 0.90)	0.66 (0.35 to 0.97)	0.83 (0.55 to 1.11)	1.00 (0.72 to 1.28)	0.28 (0.02 to 0.53)
Positive emotions	3.26 (0.69)	3.27 (0.62)	2.89 (0.68)	0.35 (0.18 to 0.52)	<.001	0.43 (0.26 to 0.60)	<.001	-0.56 (-0.84 to -0.27)	-0.60 (-0.9 to -0.3)	-0.89 (-1.17 to -0.61)	-1.10 (-1.38 to -0.81)	-0.30 (-0.55 to -0.04)
Negative emotions	2.91 (0.82)	2.92 (0.69)	3.33 (0.74)	-0.45 (-0.64 to -0.25)	<.001	-0.47 (-0.66 to -0.27)	<.001	0.64 (0.37 to 0.91)	0.65 (0.38 to 0.91)	1.06 (0.78 to 1.34)	1.21 (0.91 to 1.5)	0.50 (0.24 to 0.75)

^aWLC: waitlist control.^bBDI-II: Beck Depression Inventory-II.^cPHQ-9: 9-item Patient Health Questionnaire.^dGHQ-12: 12-item General Health Questionnaire.^eBAI: Beck Anxiety Inventory.

Table 4. Comparisons between web-based and app-based iCBT^a groups on primary and secondary outcome measures at post-treatment (intention-to-treat analysis).

Measure and time	Unadjusted mean (SD)		Adjusted difference (95% CI)	P value	Between-group effect size <i>d</i> (95% CI)	Within-group effect size (95% CI)	
	Web	App				Web	App
BDI-II^b							
Baseline	22.24 (8.52)	23.66 (8.77)	N/A ^c	N/A	N/A	N/A	N/A
Post	9.40 (8.90)	9.43 (8.45)	0.21 (-2.28 to 2.70)	.87	-0.01 (-0.3 to 0.28)	1.48 (1.18 to 1.78)	1.64 (1.33 to 1.94)
3 months	9.43 (9.58)	9.15 (8.51)	-0.14 (-3.11 to 2.83)	.93	0.03 (-0.34 to 0.4)	1.45 (1.1 to 1.8)	1.67 (1.32 to 2.02)
6 months	10.15 (9.03)	13.47 (12.24)	-1.37 (-4.46 to 1.72)	.38	-0.37 (-0.76 to 0.02)	1.40 (1.05 to 1.74)	1.04 (0.69 to 1.38)
PHQ-9^d							
Baseline	10.77 (4.9)	12.07 (4.27)	N/A	N/A	N/A	N/A	N/A
Post	5.59 (4.43)	5.24 (3.65)	0.35 (-1.08 to 1.77)	.63	0.07 (-0.22 to 0.36)	1.11 (0.8 to 1.41)	1.69 (1.36 to 2.02)
3 months	5.41 (4.01)	5.83 (3.57)	-0.27 (-1.94 to 1.41)	.75	-0.1 (-0.47 to 0.28)	1.16 (0.81 to 1.51)	1.55 (1.18 to 1.91)
6 months	5.94 (5.44)	7.65 (5.06)	-0.86 (-2.67 to 0.94)	.35	-0.39 (-0.79 to 0.01)	0.95 (0.59 to 1.31)	0.97 (0.62 to 1.33)
GHQ-12^e							
Baseline	7.46 (3.86)	8.4 (3.31)	N/A	N/A	N/A	N/A	N/A
Post	1.60 (3.06)	1.53 (2.77)	0.05 (-0.84 to 0.93)	.91	0.02 (-0.27 to 0.31)	1.64 (1.34 to 1.95)	2.21 (1.87 to 2.55)
3 months	1.85 (3.02)	1.69 (2.82)	-0.08 (-1.17 to 1)	.88	0.05 (-0.32 to 0.42)	1.54 (1.19 to 1.89)	2.12 (1.74 to 2.49)
6 months	1.77 (3.20)	3.49 (4.10)	-1.46 (-0.33 to -2.59)	.01	-0.52 (-0.91 to -0.12)	1.54 (1.19 to 1.89)	1.39 (1.03 to 1.74)
BAI^f							
Baseline	14.53 (9.29)	15.36 (8.8)	N/A	N/A	N/A	N/A	N/A
Post	8.89 (7.89)	8.49 (6.69)	0.53 (-1.25 to 2.31)	.56	0.06 (-0.23 to 0.36)	0.64 (0.37 to 0.91)	0.88 (0.59 to 1.15)
3 months	7.67 (6.42)	8.03 (6.78)	0.59 (-1.56 to 2.73)	.59	-0.05 (-0.42 to 0.32)	0.80 (0.47 to 1.12)	0.89 (0.57 to 1.21)
6 months	7.89 (7.99)	9.88 (7.72)	0.32 (-1.91 to 2.56)	.78	-0.25 (-0.64 to 0.14)	0.74 (0.42 to 1.07)	0.64 (0.31 to 0.98)
Positive automatic thoughts							
Baseline	1.85 (0.55)	1.79 (0.52)	N/A	N/A	N/A	N/A	N/A
Post	2.40 (0.71)	2.33 (0.67)	0.01 (-0.18 to 0.2)	.93	0.1 (-0.19 to 0.39)	-0.88 (-1.16 to -0.6)	-0.93 (-1.21 to -0.65)
3 months	2.49 (0.67)	2.44 (0.84)	-0.1 (-0.32 to 0.12)	.38	0.07 (-0.3 to 0.44)	-1.08 (-1.41 to -0.75)	-1.03 (-1.36 to -0.71)
6 months	2.46 (0.85)	2.15 (0.71)	0.08 (-0.16 to 0.31)	.52	0.47 (0.08 to 0.86)	-0.93 (-1.26 to -0.6)	-0.63 (-0.96 to -0.29)
Negative automatic thoughts							
Baseline	2.55 (0.88)	2.65 (0.85)	N/A	N/A	N/A	N/A	N/A
Post	1.85 (0.78)	1.89 (0.63)	-0.02 (-0.22 to 0.18)	.86	-0.04 (-0.33 to 0.25)	0.83 (0.55 to 1.1)	1 (0.72 to 1.28)
3 months	1.74 (0.68)	1.82 (0.71)	-0.03 (-0.27 to 0.21)	.81	-0.09 (-0.46 to 0.28)	0.97 (0.64 to 1.29)	1.02 (0.7 to 1.34)
6 months	1.75 (0.78)	2.05 (0.93)	-0.15 (-0.4 to 0.1)	.24	-0.38 (-0.77 to 0.01)	0.93 (0.6 to 1.26)	0.68 (0.34 to 1.02)
Positive emotions							
Baseline	2.72 (0.54)	2.65 (0.54)	N/A	N/A	N/A	N/A	N/A
Post	3.26 (0.69)	3.27 (0.62)	-0.03 (-0.22 to 0.16)	.87	-0.04 (-0.33 to 0.25)	-0.89 (-1.17 to -0.61)	-1.1 (-1.38 to -0.81)
3 months	3.26 (0.75)	3.33 (0.81)	-0.1 (-0.33 to 0.13)	.93	-0.12 (-0.49 to 0.25)	-0.89 (-1.21 to -0.56)	-1.07 (-1.39 to -0.74)
6 months	3.24 (0.75)	2.97 (0.72)	0.12 (-0.11 to 0.36)	.38	0.4 (0.01 to 0.79)	-0.85 (-1.18 to -0.52)	-0.54 (-0.88 to -0.21)

Measure and time	Unadjusted mean (SD)		Adjusted difference (95% CI)	P value	Between-group effect size <i>d</i> (95% CI)	Within-group effect size (95% CI)	
	Web	App				Web	App
Negative emotions							
Baseline	3.73 (0.75)	3.78 (0.73)	N/A	N/A	N/A	N/A	N/A
Post	2.91 (0.82)	2.93 (0.69)	-0.02 (-0.23 to 0.2)	.63	-0.02 (-0.31 to 0.28)	1.06 (0.78 to 1.34)	1.21 (0.91 to 1.5)
3 months	2.78 (0.85)	2.93 (0.73)	-0.07 (-0.32 to 0.18)	.75	-0.19 (-0.56 to 0.18)	1.23 (0.89 to 1.57)	1.17 (0.84 to 1.5)
6 months	2.86 (0.91)	3.17 (0.88)	-0.1 (-0.36 to 0.16)	.35	-0.39 (-0.78 to 0)	1.09 (0.75 to 1.42)	0.8 (0.46 to 1.13)

^aiCBT: internet-based cognitive behavioral therapy

^bBDI-II: Beck Depression Inventory-II.

^cN/A: not applicable.

^dPHQ-9: 9-item Patient Health Questionnaire.

^eGHQ-12: 12-item General Health Questionnaire.

^fBAI: Beck Anxiety Inventory.

The correction will appear in the online version of the paper on the JMIR Publications website, together with the publication of this correction notice. Because this was made after submission

to PubMed, PubMed Central, and other full-text repositories, the corrected article has also been resubmitted to those repositories.

Reference

1. Pan J-Y, Rafi J. Culturally Adapted Guided Internet-Based Cognitive Behavioral Therapy for Hong Kong People With Depressive Symptoms: Randomized Controlled Trial. *J Med Internet Res*. Feb 25, 2025;27:e64303. [FREE Full text] [doi: [10.2196/64303](https://doi.org/10.2196/64303)] [Medline: [39998865](https://pubmed.ncbi.nlm.nih.gov/39998865/)]

This is a non-peer-reviewed article. Submitted 26.Nov.2025; accepted 27.Nov.2025; published 03.Feb.2026.

Please cite as:

Pan J-Y, Rafi J

Correction: Culturally Adapted Guided Internet-Based Cognitive Behavioral Therapy for Hong Kong People With Depressive Symptoms: Randomized Controlled Trial

J Med Internet Res 2026;28:e88495

URL: <https://www.jmir.org/2026/1/e88495>

doi: [10.2196/88495](https://doi.org/10.2196/88495)

PMID:

©Jia-Yan Pan, Jonas Rafi. Originally published in the Journal of Medical Internet Research (<https://www.jmir.org>), 03.Feb.2026. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research (ISSN 1438-8871), is properly cited. The complete bibliographic information, a link to the original publication on <https://www.jmir.org/>, as well as this copyright and license information must be included.