Original Paper

Assessing Patient Attitudes to Computerized Screening in Primary Care: Psychometric Properties of the Computerized Lifestyle Assessment Scale

Farah Ahmad¹, MBBS, MPH, PhD; Sheilah Hogg-Johnson², PhD; Harvey A Skinner³, PhD

¹Centre for Research on Inner City Health, St. Michael's Hospital, Toronto, ON, Canada

²Faculty of Medicine, University of Toronto and Institute of Work and Health, Toronto, ON, Canada
³Faculty of Health, York University, Toronto, ON, Canada

Corresponding Author: Farah Ahmad, MBBS, MPH, PhD Centre for Research on Inner City Health St. Michael's Hospital 30 Bond Street Toronto, ON M5B 1W8 Canada Phone: +1 416 864 6060 ext 3310 Fax: +1 416 864 5485 Email: <u>ahmadf@smh.toronto.on.ca</u>

Abstract

Background: Computer-based health-risk assessments are electronic surveys which can be completed by patients privately, for example during their waiting time in a clinic, generating a risk report for the clinician and a recommendation sheet for the patient at the point of care. Despite increasing popularity of such computer-based health-risk assessments, patient attitudes toward such tools are rarely evaluated by reliable and valid scales. The lack of psychometric appraisal of appropriate scales is an obstacle to advancing the field.

Objective: This study evaluated the psychometric properties of a 14-item Computerized Lifestyle Assessment Scale (CLAS).

Methods: Out of 212 female patients receiving the study information at a family practice clinic, 202 completed a paper questionnaire, for a response rate of 97.6%. After 2 weeks, 52 patients completed the scale a second time.

Results: Principal component analysis revealed that CLAS is a multidimensional scale consisting of four subscales (factors): (1) Benefits: patient-perceived benefits toward the quality of medical consultation and means of achieving them, (2) Privacy-Barrier: concerns about information privacy, (3) Interaction-Barrier: concerns about potential interference in their interaction with the physician, and (4) Interest: patient interest in computer-assisted health assessments. Each subscale had good internal consistency reliability ranging from .50 (2-item scale) to .85 (6-item scale). The study also provided evidence of scale stability over time with intraclass correlation coefficients of .91, .82, .86, and .67 for the four subscales, respectively. Construct validity was supported by concurrent hypotheses testing.

Conclusions: The CLAS is a promising approach for evaluating patients' attitudes toward computer-based health-risk assessments.

(J Med Internet Res 2008;10(2):e11) doi: 10.2196/jmir.955

KEYWORDS

Computers; scale; psychometric; screening; risk assessment; family practice

Introduction

The use of computer interactive technology in health care settings is on the rise. Many studies report using patient-administered computer programs for health-risk

```
http://www.jmir.org/2008/2/e11/
```

RenderX

assessments [1-4] as well as for preventative health education in clinical settings [5-7]. Interest is particularly growing in computer-based health-risk assessments for which patients complete a computer survey privately during their waiting time. The interactive program then prints a risk report for the clinician

Ahmad et al

and a recommendation sheet for the patient at the point of care. Such computer-based health-risk assessments can facilitate meaningful communication between clinicians and patients by saving the clinicians' screening time for thorough risk management and by allowing the patients to self-reflect on their risk profile before the medical consultation.

Many practical implications have also been recognized for computer-based health-risk assessments. At the organizational level, the advantages include speed and efficiency, accountability, quality improvement, and cost containment [8]. At the individual level, convenience to providers and patients includes tailored testing, accuracy of responses, unobtrusive means of branching or skipping questions, instant feedback on risks and referrals to clinicians and patients, and aids to diagnosis [9]. Technological advances, such as touch-screen, mobile, hand-held tablet computers, have amplified the utility of computer-based health-risk assessments. Such interactive computer technology has great potential in primary care settings where provider time is constrained due to a multitude of presenting health issues and preventive needs of patients [10].

However, user attitudes toward interactive computer technology are important when considering applications. In 1986, Nickell and Pinto developed a computer attitude scale for the general population [11]. Despite good psychometric properties [12], this scale has limited applicability in the physician-patient context due to specific communication patterns between health care providers and patients. While computer attitude scales have been developed and evaluated for physicians [13,14] and other health care providers [15], reliable and valid scales for general patients are lacking [16]. Further, little knowledge exists about scale reliability over time [16].

In our review of the literature on computer-based health-risk assessments, two scales were identified as potentially applicable to general patient populations. The first scale was developed by Lucas in 1977 and tested among patients visiting specialized clinics in hospital settings [17]. This 22-item scale tapped patient attitudes toward different types of clinical encounters, including computer-assisted visits, in-person visits, and ideal health care visits. The assessments used a semantic differential technique whereby participants rate each item on a bipolar scale with contrasting adjectives at the extremes, such as hot and cold [18]. The Lucas scale was subsequently used by others in a primary care setting [2]. Yet this scale is difficult to apply in today's world of globalization, given different interpretations of adjectives by respondents of diverse ethnocultural backgrounds. Further, scales based on a semantic differential technique are lengthy and, hence, of limited use in time-pressed clinical settings.

Addressing some of these concerns, Skinner developed a short 14-item Computerized Lifestyle Assessment Scale (CLAS) in 1993 [19]. First drawing on an initial study of family practice patients [2], a large pool of items was generated through group discussions with patients and providers about the pros and cons of computer-based health-risk assessments. This list was reviewed by the research team, and the final pool of items was established through consensus among the team members. An easy to comprehend Likert-type scale was used whereby

```
http://www.jmir.org/2008/2/e11/
```

XSL•FO

participants are asked to express their level of agreement or disagreement for each item. Given the centrality of decisional balance (ie, extent of pros compared to cons) in health behavior theories and research, the CLAS focuses on patient perceived benefits and barriers toward computer-based health-risk assessments. Several studies grounded in the Transtheoretical Model and Health Belief Model demonstrate that preventive behaviors, such as cancer screening visits, improve when perceived benefits exceed perceived barriers [20,21]. Thus, assessment of the decisional balance of patients in relation to computer-based health-risk assessments is meaningful in establishing their acceptance of future use. Although CLAS is a theoretically informed scale for primary care patients, its psychometric properties have not been previously reported.

Lack of psychometric appraisal of scales may impede research and innovation to advance the field. Recent studies have begun to report patients' general reactions to the use of computer interactive technology. In 2000, Dugaw et al reported patients' overall acceptance of computerized medical history taking in an emergency department, with limited description of the measurement [22]. Likewise, recent randomized trials on computer-based health-risk assessments by Rhodes et al in a US hospital emergency department reported general reactions of patients, their recall of advice after a 1-month follow-up, and satisfaction with the visit [3,4,23]. Although information on patient satisfaction is important, it does not generate knowledge specific to technologically mediated patient concerns or facilitators in medical encounters. In 2006, MacMillan and colleagues reported patients' higher preference for computer-based screening for the risk of partner abuse compared to face-to-face inquiry. However, the preference measurement relied on three de novo questions about "ease," "like answering," and "private enough" [24]. The use of nonstandardized methods of measurement could lead to difficulties in assessing and interpreting results over time and across sites. At the same time, clinical adoption of computer-based health-risk assessments is dependent on the availability of reliable and valid knowledge about patient attitudes.

Considering the potential of CLAS, this study evaluated its psychometric properties as part of a larger research program on computer-based screening for lifestyle risks, including partner abuse, among female patients. Using standard procedures [25,26], the aim of this study was threefold: (1) to assess the dimensionality and/or latent constructs of CLAS, (2) to measure test-retest reliability and internal consistency of the instrument, and (3) to provide initial evidence on its construct validity.

Methods

The study was conducted at a multidisciplinary family practice clinic affiliated with a teaching hospital in Toronto, ON, Canada. The study was approved by the hospital research ethics board as part of a research project on prevalence of partner abuse; details are provided elsewhere [27].

Participant Recruitment

All adult female patients with an appointment were eligible to participate if they were at least 18 years of age, could speak and

read English, and could provide informed consent. The study participants were recruited in 15 days over a period of three consecutive weeks in February of 2004. On recruitment days, all adult female patients with appointments were given a brief letter of invitation by the clinic receptionist at the time of arrival. These potential participants were then approached in the waiting area by a recruiter to confirm their eligibility and inquire about their interest in the study. Willing participants were taken to a separate room in the clinic, unaccompanied by family or friends, where they completed the survey after giving informed consent. At this time (T1), participants were also asked to consent to a subsequent contact after 2 weeks (T2) to administer the CLAS a second time. Participants sealed the survey in the provided envelope before returning it to the recruiter. Then, participants received health brochures (domestic violence, cancer, and heart health) with telephone numbers for domestic violence counsellors and the assaulted women's helpline.

Measurement

The survey included the CLAS, which is a 14-item scale that covers patients' positive and negative perceptions about computer-based health-risk assessments [19]. Participants read a vignette about such a "computer survey" (Textbox 1) and rated each CLAS statement on a Likert-type scale of 1 to 5 (strongly agree, agree, not sure, disagree, strongly disagree). Other sections in the survey gathered information on sociodemographics (age, marital status, country of birth, years lived in Canada, highest education completed, employment status, and income), health (self-rated health, partner violence, and number of visits to family practice), exposure to computers (access and use), comfort in completing the survey, and English language abilities (see Table 1 for details).

Textbox 1. Vignette

We would like to know your opinion about a computer survey of patients. This survey is completed by patients on a computer before seeing their family doctor. The computer survey asks questions about lifestyle and health risks such as smoking, stress, conflict in relationships, and safety. The questions appear on the computer screen one by one. The patient answers by touching one of the options on the computer screen using a non-ink pen. Patients do not type or use any computer parts but only touch the screen to give answers. This computer survey uses simple day-to-day language of 5th grade reading level. The computer system prints (1) a summary of patient health risks for the doctor to review, and (2) an information sheet for the patients about their reported health risks. *What is your opinion about such computer-based health-risk assessment of patients before seeing a family doctor*?

Sample Size

The aim was to recruit a sample of 200 participants. As CLAS included 14 items, a sample of 200 was expected to generate an adequate subject-to-variable ratio of 14:1 to derive latent constructs. For factor analytical approaches, Gorsuch (1983) and others recommend a subject-to-variable ratio of five when the communalities are high and there are many variables for each factor [28,29]. If these conditions are not met, then a subject-to-variable ratio of 10 is recommended [28]. Others suggest that a sample of 150 should be considered sufficient when the factor analyzed solutions have several high loading markers (> 0.80) [30]. Our sample size is fair in meeting both

of the established requirements (minimum sample size and sample size per item) for psychometric studies.

Data Analyses

The CLAS items [19] were reverse coded prior to analyses so that 1 referred to "strongly disagree" and 5 to "strongly agree." All analyses were conducted using Statistical Package for Social Sciences (SPSS) version 12(SPSS Inc, Chicago, IL, USA). Preliminary statistical procedures examined distributions of the individual items (eg, means, standard deviations, skewness, and kurtosis) and evidence of ceiling or floor effects. The quality of data was evaluated by percentage of missing responses, which were low; hence, we used the case deletion in subsequent analyses. The sampling adequacy was assessed by the Kaiser-Meyer-Olkin test.

Prior to reliability and validity analysis, we examined the latent structure of the scale. The latent constructs of the CLAS were examined by employing principal component analysis (PCA) [31,32]. The number of latent constructs or factors was determined using scree plots and the criterion of eigenvalues greater than 1.0. We considered three-factor, four-factor, and five-factor solutions with varimax rotation. Salient loadings were defined using a critical value of 0.38 [28].

The scale reliability was estimated by both internal consistency and test-retest reliability of the subscales. To examine homogeneity of items or internal consistency, item-total correlation [33] and change in Cronbach alpha coefficient upon item deletion were used [34]. For test-retest reliability, intraclass correlation (ICC) from a two-way random effects model was executed [35]. We also compared T2 participants to the remaining participants at T1 with respect to characteristics measured at T1 to assess the potential to generalize the reliability findings.

After factors were derived and reliability established, construct validitywas investigated. For this analysis, we tested hypotheses that were based on existing literature. Further details on the hypotheses are presented in the Results section under construct validity. The hypotheses were tested by using Pearson product moment (r_p) , point biserial (r_{pb}) , or Spearman rank (r_s) correlation analyses, as appropriate.

Results

Participants

Among 361 women approached, 212 eligible women received the study details in privacy, 207 provided written consent (response rate 97.6%), and 202 returned the completed surveys. Participants had a mean age of 45.3 years (range 19 to 86) and 36% were immigrants, with the top two groups from Europe and Asia (Table 1). Almost 75% of the participants were in a current intimate relationship, and 77% had at least university education. Nearly 64% were currently employed and reported annual household income of at least Can \$40,000; 87% of the participants had access to computers, and 66% used one every day. Self-perceived health was rated as "good" on a scale of one to five with a mean of 3.2 (SD 1). The mean number of

visits to the family practice during the last year was 4.6 (median 3.5; mode 1; range 0 to 30).

| Table 1. Sociodemographic characteristics (N = | = 202) |
|--|--------|
|--|--------|

| Variable | No. | % |
|--|-----|-------------|
| Age (years), mean (SD) | 201 | 45.3 (15.4) |
| Current marital status | 202 | |
| Married or common law or intimate | | 74.9 |
| Separated or divorced or widowed | | 13.9 |
| Single, not in relationship | | 11.4 |
| Country of birth: Canada | 129 | 63.9 |
| If immigrant: years lived in Canada | 71 | |
| Less than 10 years | | 23.9 |
| 11 to 20 years | | 16.9 |
| More than 20 years | | 59.2 |
| If immigrant: country of birth | 72 | |
| Europe | | 36.1 |
| East or South East Asia or South Asia | | 29.1 |
| West Indies, Latin America, or Caribbean | | 20.8 |
| Middle East or West Asia | | 6.9 |
| Africa | | 5.6 |
| Highest education | 201 | |
| Less than high school | | 3.0 |
| High school, some or complete | | 19.9 |
| University or higher, some or complete | | 77.1 |
| Current employment | 201 | |
| Full-time or part-time | | 64.2 |
| Unemployed | | 13.9 |
| Retired or on disability | | 21.9 |
| Household annual income (Can \$) | 181 | |
| Less than 20,000 | | 15.5 |
| 20,001 to 40,000 | | 19.9 |
| 40,001 to 60,000 | | 20.5 |
| More than 60,000 | | 44.2 |
| Access to computer at home or work | 200 | 87.0 |
| Use of computer in the last month | 200 | |
| Every day or two to three times a week | | 81.5 |
| Once a week or once a month | | 6.5 |
| Not at all | | 12.0 |
| English ability, [*] mean (SD) | 201 | 4.5 (0.87) |
| Survey comfort level, [†] mean (SD) | 199 | 4.0 (1.2) |

*Scale of 1 to 5: 1 = poor, 2 = fair, 3 = good, 4 = very good, 5 = excellent.

[†]Scale of 1 to 5: 1 = very uncomfortable, 2 = uncomfortable, 3 = not sure, 4 = comfortable, 5 = very comfortable.

XSL•FO RenderX

Table 2. Item summary statistics and Pearson correlations

| | | - | | | | | | | | | | | | | | | | | |
|-------------------------------------|------------------------|-------------------|------|---------------|---------------|------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|-----|----|
| Item [†] | % Miss [*] | Mean [‡] | SD | Skew- ness | Kur- tosis | Item Correlation | | | | | | | | | | | | | |
| | | | | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| 1. Routine | 1.0 | 3.93 | 0.86 | -0.74 | 0.62 | 1 | | | | | | | | | | | | | |
| 2. Lifestyle | 0.5 | 3.67 | 0.94 | -0.73 | 0.39 | .58 | 1 | | | | | | | | | | | | |
| 3. Save time | 2.0 | 3.73 | 0.92 | -0.64 | 0.55 | .62 | .47 | 1 | | | | | | | | | | | |
| 4. Better as- sess | 1.0 | 3.28 | 0.84 | -0.07 | 0.17 | .54 | .43 | .50 | 1 | | | | | | | | | | |
| 5. Comfort- able | 1.0 | 3.92 | 0.85 | -0.86 | 0.93 | .60 | .52 | .41 | .34 | 1 | | | | | | | | | |
| 6. Trusted | 2.0 | 3.36 | 0.94 | -0.33 | 0.02 | .47 | .33 | .37 | .42 | .59 | 1 | | | | | | | | |
| 7. Confiden- tiality | 0.5 | 3.30 | 1.14 | -0.17 | -1.06 | 08 | 05 | 80 | 06 | 24 | 29 | 1 | | | | | | | |
| 8. Certain in- formation | 0.5 | 3.39 | 1.12 | -0.24 | -0.78 | 14 | 04 | 15 | 04 | 22 | 26 | .47 | 1 | | | | | | |
| 9. Mistakes | 1.0 | 2.63 | 0.84 | 0.51 | 0.27 | 18 | 16 | 14 | 27 | 39 | 41 | .41 | .38 | 1 | | | | | |
| 10. Less time | 2.0 | 3.35 | 0.99 | -0.34 | -0.41 | 06 | .02 | .16 | 02 | 19 | 13 | .25 | .17 | .26 | 1 | | | | |
| 11. Personal touch | 1.0 | 3.38 | 1.19 | -0.14 | -1.12 | 32 | 24 | 20 | 30 | 34 | 32 | .28 | .40 | .43 | .40 | 1 | | | |
| 12. Another doctor [§] | 0.5 | 2.13 | 0.98 | 0.98 | 0.92 | 35 | 28 | 30 | 33 | 47 | 27 | .29 | .26 | .42 | .32 | .50 | 1 | | |
| 13. Answer honestly [§] | 1.5 | 4.42 | 0.68 | -1.36 | 3.36 | .40 | .19 | .32 | .23 | .50 | .28 | 07 | 19 | 20 | 12 | 20 | 35 | 1 | |
| 14. No pat info [§] | 4.5 | 4.26 | 0.70 | -0.97 | 2.15 | .35 | .19 | .19 | .20 | .25 | .23 | 04 | 03 | 06 | 06 | 06 | 12 | .34 | 1 |

^{*}% Miss, % missing response.

[†]Full item statements are provided in Table 3 and Table 4.

[‡]Scale of 1 (strongly disagree) to 5 (strongly agree).

[§]Skewed items.

Item Descriptive Statistics

The item means and standard deviations were acceptable, while three items were skewed (Table 2). These items were transformed and PCA was executed with and without transformations. As the two PCAs were similar in factor structure and factor loadings, we report PCA without transformed items in this paper. Sampling adequacy was indicated by a Kaiser-Meyer-Olkin test value of .82 and the absence of ceiling or floor effects. The factorability was indicated by correlation and partial correlation matrices.

Factor Structure

On conducting the PCA, the first 10 eigenvalues were 4.7, 2.1, 1.1, 1.0, .85, .76, .69, .53, .50, and .44. Four factors emerged

with eigenvalues greater than or equal to one, accounting for 63.7% of the total variance. Based on the scree plot, either a three-factor or four-factor solution was indicated. We considered three-, four-, and five-factor solutions, and the four-factor solution yielded the most interpretable results. A summary of the PCA with varimax rotation is presented in Table 3. The named Benefits, Privacy-Barrier, factors were Interaction-Barrier, and Interest. Three- and four-factor solutions were also compared for the internal consistency of the derived subscales. Although the Privacy-Barrier and Interaction-Barrier factors merged into one factor upon forcing a three-factor solution, the internal consistency of the subscales was higher in the four-factor solution than in the three-factor solution. This internal consistency comparison was based on the reliability coefficients adjusted for the length of the subscales [33].



Ahmad et al

Table 3. Summary of principal component analysis with varimax rotation

| Item | Factor Load | h ^{2*} | | | |
|--|-------------|--------------------|--------------------|----------|-----|
| | Benefits | Privacy | Interaction | Interest | |
| | | -Barrier | -Barrier | | |
| 1. Computers will help doctors with routine lifestyle questions | .79 | | | | .74 |
| 2. The computer is a good way to ask lifestyle questions | .79 | | | | .65 |
| 3. It would save doctors time. | .78 | | | | .65 |
| 4. Doctors will make better assessments with such computer systems | .74 | | | | .60 |
| 5. I would feel comfortable answering questions on a computer | .58 | | | | .61 |
| 6. Computers can be trusted | .54 | (.41) [†] | | | .51 |
| 7. I would worry about confidentiality | | .82 | | | .69 |
| 8. I do not want certain information about me on the computer | | .81 | | | .69 |
| 9. Too many mistakes will be made with computer | | .60 | (.39) [†] | | .55 |
| 10. Doctors would spend less time with patients | | | .81 | | .71 |
| 11. There will be loss of personal touch of a doctor | | | .69 | | .67 |
| 12. I would find another doctor | | | .63 | | .57 |
| 13. I would want to read patient information sheet | | | | .80 | .67 |
| 14. I would answer honestly | | | | .74 | .62 |

 $*h^2$ refers to communalities.

[†]Item shared loading between factors above the critical value.

Variances accounted for by the four identified factors (Benefits, Privacy-Barrier, Interaction-Barrier, and Interest) after the rotation were 33.6%, 15.0%, 8.0%, and 7.2%, respectively. The item "Computers can be trusted" in the first factor (Benefits) shared loading (.41) with the second factor (Privacy-Barrier) above the critical value of .38. Also, the item "Too many mistakes will be made with computer" in the second factor (Privacy-Barrier) shared loading (.39) with the third factor (Interaction-Barrier) above the critical value.

The Benefits factor consisted of six items with factor loadings ranging from .79 to .54. The items loading on this factor cover perceived benefits toward the quality of medical consultation and means of achieving the benefits. The Privacy-Barrier factor consisted of three items dealing with patient concerns about privacy, with loadings ranging from .82 to .60. The Interaction-Barrier factor consisted of three items covering

patient concerns about interference in the interaction with the physician, with loadings ranging from .81 to .63. Although the Interest factor consisted of only two items, both items had strong factor weightings (ie, .80 and .79). The stability of this factor was also apparent during execution of the five-factor solution. Both items of this factor continued to load together while the fifth factor consisted of one item pulled from the Interaction-Barrier factor.

Reliability

To estimate internal consistency reliability, we considered the following criteria for each subscale: (1) an item-total correlation of at least .3 for all items, (2) no increase in the Cronbach alpha coefficient if an item was deleted, and (3) general acceptability of the item means and standard deviations. All three criteria were met for the subscales (Table 4).



Table 4. Internal consistency of the subscales

| Item | Mean (SD) [*] | Corrected | Cronbach Alpha |
|--|------------------------|-----------------------------|-----------------|
| | | Item-Total Corre- lation | if Item Deleted |
| Benefits: Cronbach alpha .85 | | | |
| 1. Computers will help doctors with routine lifestyle questions | 3.9 (0.87) | .77 | .80 |
| 2. The computer is a good way to ask lifestyle questions | 3.7 (0.94) | .62 | .83 |
| 3. It would save doctors time. | 3.7 (0.91) | .61 | .83 |
| 4. Doctors will make better assessments with such computer systems | 3.3 (0.85) | .59 | .83 |
| 5. I would feel comfortable answering questions on a computer | 3.9 (0.85) | .66 | .82 |
| 6. Computers can be trusted | 3.4 (0.95) | .57 | .84 |
| Privacy-Barrier: Cronbach alpha .70 (alpha $.81^{\dagger}$) | | | |
| 7. I would worry about confidentiality | 3.3 (1.1) | .53 | .54 |
| 8. I do not want certain information about me on the computer | 3.4 (1.1) | .52 | .55 |
| 9. Too many mistakes will be made with computer | 2.6 (0.84) | .46 | .64 |
| Interaction-Barrier: Cronbach alpha .67 (alpha .80 †) | | | |
| 10. Doctors would spend less time with patients | 3.4 (0.99) | .42 | .66 |
| 11. There will be loss of personal touch of a doctor | 3.4 (1.2) | .56 | .47 |
| 12. I would find another doctor | 2.1 (0.97) | .49 | .57 |
| Interest: Cronbach alpha .50 (alpha .75 †) | | | |
| 13. I would want to read patient information sheet | 4.3 (0.70) | .34 | - |
| 14. I would answer honestly | 4.4 (0.63) | .34 | _ |

*Scale 1 to 5: strongly disagree, agree, not sure, agree, strongly agree.

[†]Adjusted reliability coefficient, adjusted to compare to scales with six items.

The Cronbach alpha coefficients for the four subscales Benefits, Privacy-Barrier, Interaction-Barrier, and Interest were .85, .70, .67, and .50, respectively. There was no increase in Cronbach alpha if items were deleted from the first three subscales. This analysis did not apply to the Interest subscale as it had two items only. The item-total correlation for the subscales Benefits, Privacy-Barrier, and Interaction-Barrier ranged from .77 to .57, .53 to .46, and .52 to .44, respectively. We also calculated the reliability coefficients adjusted for the length of subscale [33], given that the number of items loading on the subscales varied and that Cronbach alpha is sensitive to number of items. The adjusted reliability coefficients were .81, .80, and .75 for the Privacy-Barrier, Interaction-Barrier, and Interest subscales, respectively, where adjustment was made to assume six items as for the Benefits subscale. This analysis assumes that the new items would be similar to the old items with respect to content and reliability.

Scale reliability over time was assessed with the test-retest data (n = 52). At T2, 52 patients were successfully reached out of 145 T1 participants who consented to the second contact. The reduced participation at T2 was due to (1) the study requirement that the second administration of the CLAS occur within 2 weeks of the first administration, and (2) the fact that many patients were difficult to reach because they had provided telephone numbers at work. The T2 participants were similar to the other T1 participants (n = 150) on sociodemographic

```
http://www.jmir.org/2008/2/e11/
```

RenderX

characteristics, including age, country of birth, number of years lived in Canada, education level, employment status, income, English language abilities, access to computers, computer use in the last month, relationship status, experiences of intimate partner violence, number of visits to family practice, and perceived health. However, the T2 participants were less likely to be employed than participants who consented but could not be reached for second contact ($\chi^2_2 = 7.0$, P < .05). The time between T1 and T2 contacts averaged 16 days (SD 2.6, median 15, mode 15). The ICC analysis based on a two-way random effect model gave coefficients of .76 for the overall scale and .91, .82, .86, and .67 for the subscales of Benefits, Privacy-Barrier, Interaction-Barrier, and Interest, respectively. As the CLAS is a multidimensional scale, the test-retest reliability of the subscales was higher than the test-retest reliability of the overall scale.

Construct Validity

To evaluate validity of the derived constructs, several hypotheses were formulated based on a literature review. We hypothesized that the Benefits factor would be positively associated with participants' frequent use of computers as greater familiarity with computers is likely to increase peoples' comfort and perceptions of the benefits [2]. Also, we hypothesized that patients with poorer health would perceive the benefits of computer-based screening as high due to the limited time

available for lifestyle inquiries during their routine health care visits. As computer-based screening has been found specifically beneficial for socially sensitive issues [2-4,36-39], it was hypothesized that participants reporting victimization by intimate partners would perceive the benefits as high. Existing studies report that patients are likely to perceive barriers in using preventive health services if they have low socioeconomic status or are immigrants [40,41]. Accordingly, it was hypothesized that the Privacy-Barrier and Interaction-Barrier factors would be positively associated with participants' non-Canadian-born status, low household income, unemployment, and lesser years of education. We also hypothesized that the Interest factor would be significantly associated with less use of computers and older age.

The hypotheses were tested by correlation analyses. The Benefits factor was positively associated with poorer self-perceived health and intimate partner victimization ($r_p = .15$, P = .03; $r_{pb} = .19$, P = .02) as hypothesized. However, it was not significantly associated with the use of computers, in contrast to our hypothesis. To explore further, we examined the mean scores of the Benefits subscale by participants' frequency of computer use in the last month. Participants who used computers every day or two to three times a week somewhat agreed with the Benefits (mean 3.7, SD 0.67), while participants who used computers once a week or once a month (mean 3.5, SD 0.50) or not at all (mean 3.6, SD 0.67) seemed to neither agree nor disagree with the Benefits.

As hypothesized, the Privacy-Barrier and Interaction-Barrier factors had positive significant associations with participants' non-Canadian-born status ($r_{\rm pb}$ = .19, P = .006; $r_{\rm pb}$ = .22, P = .001), low household income ($r_{\rm p}$ = .23, P = .002; $r_{\rm p}$ = .21, P = .004), and lower use of computers ($r_{\rm s}$ = .16, P = .03; $r_{\rm s}$ = .18, P = .01). Furthermore, older age at the time of immigration had a positive association with both the Privacy-Barrier and Interaction-Barrier factors ($r_{\rm p}$ = .27, P = .02; $r_{\rm p}$ = .28, P = .02). The Interaction-Barrier factor also had significant associations with participants' unemployment status and lesser years of education ($r_{\rm pb}$ = .16, P = .03; $r_{\rm s}$ = .18, P < .01). The Interest factor had significant positive associations with older age ($r_{\rm p}$ = .16, P = .03) and less use of computers ($r_{\rm s}$ = .14, P = .04).

Discussion

The CLAS has demonstrated good preliminary psychometric properties and shows promise as a tool for assessing patient attitudes toward computer-based health-risk assessments. Each of the four latent constructs or derived subscales of the CLAS had good internal consistency that exceeded the recommended threshold of 0.7 [42] after adjusting for the number of items. Furthermore, the multidimensionality of the CLAS highlights different clusters of barriers perceived by patients in the use of interactive technology, namely privacy and interaction with physicians. This study also provides much needed initial evidence of the scale stability over time through test-retest analysis. This is important as some researchers and health care interventionists aim to assess patient attitudes toward computer-based health-risk assessments before and after new initiatives.

Implications

The use of a psychometrically validated scale is an essential element in facilitating clinical and policy decisions about the application of computer-based health-risk assessments. This is of particular importance for sensitive health risks and conditions where superiority of computer-based risk assessments over personal interviews is already well documented with respect to patient disclosure of socially sensitive information. These health risks and conditions include behaviors related to sex, alcohol, drugs, HIV, and violence [2-4,36-39]. A similar link is demonstrated in our study as a positive association between women's victimization at the hands of their intimate partner and the Benefits subscale. Literature shows that women experiencing partner abuse seldom spontaneously disclose it to health care providers [43,44], who frequently fail to detect victims of abuse due to time pressure, priority of acute problems, and discomfort [45,46]. At the same time, clinicians' questioning about abuse is the most significant predictor of women's disclosure [47]. Computer-based screening matches abused women's preferences for "direct questioning," and it has limited dependency on physician time. Above all, it is a nonjudgmental and anonymous way of asking about socially sensitive health risks. Perhaps it explains why abused women in our study perceived higher benefits of the computer-based screening. Our future work will test the computer-based screening intervention in a family practice setting for the detection and disclosure of partner abuse.

The findings also highlight the complex nature of human behavior. Study participants perceived barriers in two distinct ways: barriers regarding privacy and barriers regarding interaction with physicians. At the implementation level, this underscores the need to measure both domains to understand and thereafter address effective use of computer-based health-risk assessments. At the theoretical level, this distinction is novel to the original conception of the scale. Possibly, patient attitudes have taken specific forms with the increasing use of computers. Recent studies reveal that use of the Internet for health information influences the way people relate to physicians, make medical decisions, and access health services [48,49]. In 2007, a telephone survey with 2479 Canadians examined their attitudes toward electronic health information and their privacy [50]. The survey found that 9 out of 10 people perceived the use of electronic health information as integral to the provision of high-quality care but had mixed confidence about the protection of health information. Future research should further examine the domains of privacy and interaction barriers in the use of computer-based health-risk assessments. Other studies report that patients' perceptions toward computer-based lifestyle assessments are positively increased after they are provided the actual experience [2,19].

Our post hoc analyses indicate that study participants who were immigrants or had lower socioeconomic status perceived more barriers. This raises two critical questions: (1) Is this an extension of the "digital divide?", and (2) What does it mean for implementation? The term "digital divide" stems from

research and refers to "decreased access to information technologies, particularly the Internet, for racial and ethnic minorities, person with disabilities, rural populations, and those with low socioeconomic status" [51] (p 449). The digital divide requires vigilance when using certain health information technologies, such as the Internet [52,53]. In contrast, computer-based health-risk assessments in health care settings bridge the digital divide because these programs provide tailored health information to the patients at the point of care. They may play a positive role in addressing patients' unequal access to health information and care—an anticipated impact similar to telemedicine [54].

Limitations

Several limitations of this study should be noted. The CLAS predominantly measures the decision-making aspect of human behavior, though it has relevance for research on explaining and changing behavior regarding computerized assessments. Future studies should explore other aspects such as patient self-efficacy and cue-to-action. The construct of Interest would also benefit from further conceptual development. Further, our analysis of the construct validity is post hoc in nature. Many of the correlations were not strong even when significant. This is possibly due to our convenience-based use of a larger survey to select variables which in turn had a more distal than proximal relationship with the CLAS constructs. Although we found support for most of our hypothesized relationships, the Benefits subscale was not associated with the participants' use of computers, contrary to our hypothesis. The study sample was relatively more educated than the average general population, and 87% of the participants had access to computers at home or work; almost a similar proportion reported using the computer every day or at least two to three times a week. Perhaps frequent use of computers makes people think critically about their advantages and disadvantages, leading to a cautious assessment of their benefits. On other side, it is also possible that computers have now become part of our everyday life and their benefits are taken for granted, reducing the level of perceived benefits seen a few years ago. Future research with larger samples should examine this further and establish the construct validity with a priori selection of variables. Also, it will be important to conduct

a classic multitrait-multimethod study in which the four constructs on the CLAS are assessed via different methods (eg, peer ratings, behavior observations). This type of study will provide evidence for both convergent and discriminate aspects of the CLAS construct validity.

Caution is warranted regarding the generalizability of our study findings. We evaluated psychometric properties of the CLAS with female patients only. A future study involving both men and women is needed to ensure its applicability to all patients visiting primary health care settings. Further, patients were recruited from a single site. However, the collaborating clinic had several physicians and served a large number of diverse patients with estimated annual visits of 50,000. The study obtained a high response rate and, reassuringly, the participants were similar to females residing in Toronto in terms of immigration and marital status [55,56]. At the same time, study participants had relatively higher levels of income and education than the general population. The test-retest results of our study may have limited generalizability as participants in the second administration of the CLAS were more likely to be unemployed than the rest of participants. Nevertheless, the two groups were similar for all other sociodemographic and health-related variables that were measured. Further research is needed with a heterogeneous sample as an important next step to advance the generalizability of the scale.

Conclusion

This study of patients in a family practice setting advances our understanding of the properties, applicability, and generalizability of the CLAS. This is an important improvement over previous assessments of other scales that relied on samples of convenience or were not specific to patient populations. Furthermore, the phrasing of items in the CLAS is expected to allow people from different ethnocultural backgrounds to reply in a meaningful way, unlike some other existing scales. At the same time, future research with a heterogeneous sample is needed to enhance its generalizability by gender and socioeconomic status while examining the utility for low and high users of computers. In conclusion, this study is a step toward facilitating research and interventions for promoting patient acceptance of computer interactive technology.

Acknowledgments

The study contributed to doctoral and fellowship training of Farah Ahmad, funded by the Canadian Institutes of Health Research (No. 17744), the Institute of Gender and Health, and Ontario Women's Health Council. The authors wish to thank Wendy Levinson, thesis/fellowship supervisor, and Donna E. Stewart, thesis advisor, for their conceptual contributions. All in-kind support provided by the Centre for Research on Inner City Health, The Keenan Research Centre in the Li Ka Shing Knowledge Institute of St. Michael's Hospital, is much appreciated.

Conflicts of Interest

None declared.

References

- 1. Lucas RW, Mullin PJ, Luna CB, McInroy DC. Psychiatrists and a computer as interrogators of patients with alcohol-related illnesses: a comparison. Br J Psychiatry. Aug 1977;131:160-167. [Medline: <u>334310</u>]
- Skinner HA, Allen BA, McIntosh MC, Palmer WH. Lifestyle assessment: applying microcomputers in family practice. Br Med J (Clin Res Ed). Jan 19, 1985;290(6463):212-214. [FREE Full text] [Medline: <u>3917757</u>]

- 3. Rhodes KV, Lauderdale DS, Stocking CB, Howes DS, Roizen MF, Levinson W. Better health while you wait: a controlled trial of a computer-based intervention for screening and health promotion in the emergency department. Ann Emerg Med. Mar 2001;37(3):284-291. [Medline: <u>11223765</u>] [doi: <u>10.1067/mem.2001.110818</u>]
- 4. Rhodes KV, Lauderdale DS, He T, Howes DS, Levinson W. "Between me and the computer": increased detection of intimate partner violence using a computer questionnaire. Ann Emerg Med. Nov 2002;40(5):476-484. [FREE Full text] [Medline: 12399790] [doi: 10.1067/mem.2002.127181]
- 5. Jenny NYY, Sing T. Evaluating the effectiveness of an interactive multimedia computer-based patient education program in cardiac rehabilitation. Occup Ther J Res. 2001;21(4):260-275.
- 6. Schinke SP, Di Noia J, Glassman JR. Computer-mediated intervention to prevent drug abuse and violence among high-risk youth. Addict Behav. Jan 2004;29(1):225-229. [Medline: <u>14667434</u>] [doi: <u>10.1016/j.addbeh.2003.07.002</u>]
- Prochaska JJ, Zabinski MF, Calfas KJ, Sallis JF, Patrick K. PACE+: interactive communication technology for behavior change in clinical settings. Am J Prev Med. Aug 2000;19(2):127-131. [Medline: <u>10913904</u>] [doi: <u>10.1016/S0749-3797(00)00187-2</u>]
- Flower J. Transformations of 21st century health care, Part 1. Beyond the digital divide. Health Forum J. 2003;46(1):8-13, 1. [Medline: <u>12690711</u>]
- 9. Gavin DR, Skinner HA, George MS. Computerized approaches to alcohol assessment. In: Litten R, Allen J, editors. Measuring Alcohol Consumption. Totowa, NJ. The Humana Press Inc; 1992.
- 10. Glasgow RE, McKay HG, Boles SM, Vogt TM. Interactive computer technology, behavioral science, and family practice. J Fam Pract. Jun 1999;48(6):464-470. [Medline: 10386491]
- 11. Nickell GS, Pinto JN. The computer attitude scale. Comput Human Behav. 1986;2(4):301-306. [doi: 10.1016/0747-5632(86)90010-5]
- 12. Rainer J, Kelly R, Miller MD. An assessment of the psychometric properties of the computer attitude scale. Comput Human Behav. 1996;12(1):93-105. [doi: 10.1016/0747-5632(95)00021-6]
- Cork RD, Detmer WM, Friedman CP. Development and initial validation of an instrument to measure physicians' use of, knowledge about, and attitudes toward computers. J Am Med Inform Assoc. 1998;5(2):164-176. [FREE Full text] [Medline: 9524349]
- 14. Brown SH, Coney RD. Changes in physicians' computer anxiety and attitudes related to clinical information system use. J Am Med Inform Assoc. 1994;1(5):381-394. [FREE Full text] [Medline: 7850562]
- 15. Shumway JM, Jacknowitz AI, Abate MA. Analysis of physicians', pharmacists', and nurses' attitudes toward the use of computers to access drug information. Methods Inf Med. Mar 1990;29(2):99-103. [Medline: <u>2342434</u>]
- 16. Shaft TM, Sharfman MP, Wu WW. Reliability assessment of the attitude towards computers instrument. Comput Human Behav. 2004;20(5):661-689. [doi: <u>10.1016/j.chb.2003.10.021</u>]
- 17. Lucas RW. A study of patients' attitudes to computer interrogation. J Man-Machine Studies. 1977;9:69-86.
- 18. Osgoode CE, Suci GJ, Tannenbaum PH. The measurement of meaning. Urbana, IL. University of Illinois Press; 1957.
- Skinner HA. Early identification of addictive behaviors using computerized life-style assessment. In: Baer JS, Marlatt GA, McMahon RJ, editors. Addictive Behaviors across the Lifespan: Prevention, Treatment and Policy Issues. Newbury Park, CA. Sage; 1993;89-110.
- 20. Prochaska JO, Redding CA, Evers KE. The Transtheoretical Model and stages of change. In: Glanz K, Lewis FM, Rimer BK, editors. Health Behavior and Health Education. 2nd edition. San Francisco, CA. Jossey-Bass Inc; 1997;60-84.
- 21. Strecher VJ, Rosenstock IM. The Health Belief Model. In: Glanz K, Lewis FM, Rimer BK, editors. Health Behavior and Health Education. 2nd edition. San-Francisco, CA. Jossey-Bass Inc; 1997;41-59.
- 22. Dugaw JE, Civello K, Chuinard C, Jones GN. Will patients use a computer to give a medical history? J Fam Pract. Oct 2000;49(10):921-923. [Medline: <u>11052165</u>]
- 23. Rhodes KV, Drum M, Anliker E, Frankel RM, Howes DS, Levinson W. Lowering the threshold for discussions of domestic violence: a randomized controlled trial of computer screening. Arch Intern Med. May 22, 2006;166(10):1107-1114. [FREE Full text] [Medline: 16717173] [doi: 10.1001/archinte.166.10.1107]
- 24. MacMillan HL, Wathen CN, Jamieson E, Boyle M, McNutt LA, Worster A, et al; McMaster Violence Against Women Research Group. Approaches to screening for intimate partner violence in health care settings: a randomized trial. JAMA. Aug 2, 2006;296(5):530-536. [FREE Full text] [Medline: <u>16882959</u>] [doi: <u>10.1001/jama.296.5.530</u>]
- 25. DeVellis RF. Scale Development: Theory and Applications. 2nd edition. Newbury Park, CA. Sage; 1991.
- 26. Specter PE. Summated Rating Scale Construction. Newbury Park, CA. Sage; 1992.
- 27. Ahmad F, Hogg-Johnson S, Stewart DE, Levinson W. Violence involving intimate partners: prevalence in Canadian family practice. Can Fam Physician. Mar 2007;53(3):461-8, 460. [FREE Full text] [Medline: <u>17872682</u>]
- 28. Norman GR, Streiner DL. Factor analysis. In: Norman GR, Streiner DL, editors. Biostatistics: The Bare Essentials. Hamilton, ON. BC Decker Inc; 2000;163-176.
- 29. Gorsuch RL. Factor Analysis. Hilldale, NJ. Lawrence Erlbaum Associates; 1983.
- 30. Guadagnoli E, Velicer WF. Relation of sample size to the stability of component patterns. Psychol Bull. Mar 1988;103(2):265-275. [Medline: <u>3363047</u>] [doi: <u>10.1037/0033-2909.103.2.265</u>]

RenderX

- 31. Tabachnick BG, Fidell LS. Principal Components and Factor Analysis. Using Multivariate Statistics. 4th edition. Needham Heights, MA. Allyn and Bacon; 2001;582-652.
- 32. Kline P. An Easy Guide to Factor Analysis. New York. Routledge; 1994.
- 33. Nunnally JCJ. Introduction to Psychological Measurement. New York. McGraw-Hill; 1970.
- 34. Cronbach LJ. Coefficient Alpha and the Internal Structure of Tests. Psychometrika. 1951;16(3):297-334. [doi: 10.1007/BF02310555]
- 35. Shrout PE, Fleiss JL. Intraclass Correlations: Uses in Assessing Rater Reliability. Psychol Bull. 1979;86(2):420-428. [doi: 10.1037/0033-2909.86.2.420]
- Turner CF, Ku L, Rogers SM, Lindberg LD, Pleck JH, Sonenstein FL. Adolescent sexual behavior, drug use, and violence: increased reporting with computer survey technology. Science. May 8, 1998;280(5365):867-873. [FREE Full text] [Medline: 9572724] [doi: 10.1126/science.280.5365.867]
- Locke SE, Kowaloff HB, Hoff RG, Safran C, Popovsky MA, Cotton DJ, et al. Computer-based interview for screening blood donors for risk of HIV transmission. JAMA. Sep 9, 1992;268(10):1301-1305. [FREE Full text] [Medline: <u>1507376</u>] [doi: <u>10.1001/jama.268.10.1301</u>]
- 38. Lucas RW, Card WI, Knill-Jones RP, Watkinson G, Crean GP. Computer interrogation of patients. Br Med J. Sep 11, 1976;2(6036):623-625. [FREE Full text] [Medline: 786434]
- 39. Tourangeau R, Smith TW. Asking sensitive questions. The impact of data collection mode, question format, and question context. Public Opin Q. 1996;60(2):274-304. [doi: 10.1086/297751]
- 40. Maxwell CJ, Kozak JF, Desjardins-Denault SD, Parboosingh J. Factors important in promoting mammography screening among Canadian women. Can J Public Health. Sep 1997;88(5):346-350. [FREE Full text] [Medline: 9401172]
- 41. Maxwell CJ, Bancej CM, Snider J. Predictors of mammography use among Canadian women aged 50-69: findings from the 1996/97 National Population Health Survey. CMAJ. Feb 6, 2001;164(3):329-334. [FREE Full text] [Medline: <u>11232132</u>]
- 42. Nunnally JC, Berstein IH. Psychometric Theory. New York. McGraw-Hill; 1994.
- 43. Reidy R, Von Korff M. Is battered women's help seeking connected to the level of their abuse? Public Health Rep. 1991;106(4):360-364. [FREE Full text] [Medline: 1908586]
- 44. Rodriguez MA, Quiroga SS, Bauer HM. Breaking the silence. Battered women's perspectives on medical care. Arch Fam Med. Mar 1996;5(3):153-158. [Medline: <u>8620256</u>] [doi: <u>10.1001/archfami.5.3.153</u>]
- 45. Rodriguez MA, Bauer HM, McLoughlin E, Grumbach K. Screening and intervention for intimate partner abuse: practices and attitudes of primary care physicians. JAMA. Aug 4, 1999;282(5):468-474. [FREE Full text] [Medline: 10442663] [doi: 10.1001/jama.282.5.468]
- 46. Sugg NK, Inui T. Primary care physicians' response to domestic violence. Opening Pandora's box. JAMA. Jun 17, 1992;267(23):3157-3160. [Medline: <u>1593735</u>] [doi: <u>10.1001/jama.267.23.3157</u>]
- 47. Rodríguez MA, Sheldon WR, Bauer HM, Pérez-Stable EJ. The factors associated with disclosure of intimate partner abuse to clinicians. J Fam Pract. Apr 2001;50(4):338-344. [Medline: <u>11309220</u>]
- 48. Murray E, Lo B, Pollack L, Donelan K, Catania J, Lee K, et al. The impact of health information on the Internet on health care and the physician-patient relationship: national U.S. survey among 1.050 U.S. physicians. J Med Internet Res. Aug 29, 2003;5(3):e17. [FREE Full text] [Medline: <u>14517108</u>] [doi: <u>10.2196/jmir.5.3.e17</u>]
- 49. Ahmad F, Hudak PL, Bercovitz K, Hollenberg E, Levinson W. Are physicians ready for patients with Internet-based health information? J Med Internet Res. 2006;8(3):e22. [FREE Full text] [Medline: <u>17032638</u>] [doi: <u>10.2196/jmir.8.3.e22</u>]
- 50. ; EKOS Research Associates. Electronic Health Information and Privacy Survey: What Canadians Think 2007. Executive Summary. Ottawa, ON. EKOS Research Associates; 2007. URL: <u>http://www.infoway-inforoute.ca/Admin/Upload/Dev/ Document/EKOS Final%20report Executive%20Summary EN.pdf%20</u> [accessed 2008 Apr 14] [WebCite Cache ID <u>5WenlFzSy</u>]
- Chang BL, Bakken S, Brown SS, Houston TK, Kreps GL, Kukafka R, et al. Bridging the digital divide: reaching vulnerable populations. J Am Med Inform Assoc. 2004;11(6):448-457. [FREE Full text] [Medline: 15299002] [doi: 10.1197/jamia.M1535]
- 52. Bernhardt JM. Health education and the digital divide: building bridges and filling chasms. Health Educ Res. Oct 2000;15(5):527-531. [FREE Full text] [Medline: <u>11184212</u>] [doi: <u>10.1093/her/15.5.527</u>]
- Smith-Barbaro PA, Licciardone JC, Clarke HF, Coleridge ST. Factors associated with intended use of a Web site among family practice patients. J Med Internet Res. May 17, 2001;3(2):E17. [FREE Full text] [Medline: <u>11720959</u>] [doi: <u>10.2196/jmir.3.2.e17</u>]
- 54. ; President's Advisory Commission on Consumer Protection and Quality in the Health Care Industry. Quality First. Better Health Care for All Americans. Advisory Commission on Consumer Protection and Quality in the Health Care Industry; 1998. URL: <u>http://www.hcqualitycommission.gov/final</u> [accessed 2008 Apr 14] [WebCite Cache ID 5PcolvCUr]
- 55. ; Statistics Canada. Immigrant population by place of birth, by census metropolitan areas (2006 Census). In: Statistics Canada, Census of Population. Ottawa, ON. Statistics Canada; 2007. URL: <u>http://www40.statcan.ca/l01/cst01/demo35c.</u> <u>htm</u> [accessed 2008 Apr 14] [WebCite Cache ID 5X6070QPM]



RenderX

56. ; Statistics Canada. Population in private households, showing living arrangements, by census metropolitan areas (2001 Census). In: Statistics Canada, Census of Population. Ottawa, ON. Statistics Canada; 2005. URL: <u>http://www40.statcan.ca/101/cst01/famil52g.htm</u> [accessed 2008 Apr 14] [WebCite Cache ID 5XAgeLEXj]

Abbreviations

CLAS: Computerized Lifestyle Assessment Scale **ICC:** intraclass correlation **PCA:** principal component analysis

Edited by G Eysenbach; submitted 12.07.07; peer-reviewed by MA Vallejo Pareja, L Nichols, P Bendtsen; comments to author 28.01.08; revised version received 28.03.08; accepted 08.04.08; published 18.04.08

<u>Please cite as:</u> Ahmad F, Hogg-Johnson S, Skinner HA Assessing Patient Attitudes to Computerized Screening in Primary Care: Psychometric Properties of the Computerized Lifestyle Assessment Scale J Med Internet Res 2008;10(2):e11 URL: <u>http://www.jmir.org/2008/2/e11/</u> doi: <u>10.2196/jmir.955</u> PMID: <u>18440918</u>

© Farah Ahmad, Sheilah Hogg-Johnson, Harvey A Skinner. Originally published in the Journal of Medical Internet Research (http://www.jmir.org), 18.04.2008. Except where otherwise noted, articles published in the Journal of Medical Internet Research are distributed under the terms of the Creative Commons Attribution License (http://www.creativecommons.org/licenses/by/2.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided 1) the original work is properly cited, including full bibliographic details and the original article URL on www.jmir.org, and 2) this statement is included.

