

Original Paper

Ability to Generate Patient Registries Among Practices With and Without Electronic Health Records

Adam Wright^{1,2}, PhD; Elizabeth A McGlinchey³, BA; Eric G Poon², MD, MPH; Chelsea A Jenter³, MPH; David W Bates^{2,3}, MD, MSc; Steven R Simon⁴, MD, MPH

¹Clinical Informatics Research & Development, Partners Healthcare System, Boston, MA, USA

²Division of General Medicine, Brigham & Women's Hospital, Boston, MA, USA

³Clinical and Quality Analysis, Partners Healthcare System, Boston, MA, USA

⁴Department of Ambulatory Care and Prevention, Harvard Medical School and Harvard Pilgrim Health Care, Boston, MA, USA

Corresponding Author:

Adam Wright, PhD

Division of General Medicine

Brigham and Women's Hospital

75 Francis St

Boston, MA 02115

USA

Phone: +1 781 416 8764

Fax: +1 781 416 8912

Email: awright5@partners.org

Abstract

Background: The ability to generate registries of patients with particular clinical attributes, such as diagnoses or medications taken, is central to measuring and improving the quality of health care. However, it is not known how many providers have the ability to generate such registries.

Objectives: To assess the proportion of physician practices that can construct registries of patients with specific diagnoses, laboratory results, or medications, and to determine the relationship between electronic health record (EHR) usage and the ability to perform registry functions.

Methods: We conducted a mail survey of a stratified random sample of physician practices in Massachusetts in the northeastern United States (N = 1884). The survey included questions about the physicians' ability to generate diagnosis, laboratory result, and medication registries; the presence of EHR; and usage of specific EHR features.

Results: The response rate was 71% (1345/1884). Overall, 79.8% of physician practices reported being able to generate registries of patients by diagnosis; 56.1% by laboratory result; and 55.8% by medication usage. In logistic regression analyses, adjusting for urban/rural location, practice size and ownership, teaching status, hospital affiliation, and specialty, physician practices with an EHR were more likely to be able to construct diagnosis registries (adjusted odds ratio [OR] 1.53, 95% confidence interval [CI] 1.25 - 1.86), laboratory registries (OR 1.42, 95% CI 1.22 - 1.66), and medication registries (OR 2.30, 95% CI 1.96 - 2.70).

Conclusions: Many physician practices were able to generate registries, but this capability is far from universal. Adoption of EHRs appears to be a useful step toward this end, and practices with EHRs are considerably more likely to be able to carry out registry functions. Because practices need registries to perform broad-based quality improvement, they should consider adopting EHRs that have built-in registry functionality.

(*J Med Internet Res* 2009;11(3):e31) doi: [10.2196/jmir.1166](https://doi.org/10.2196/jmir.1166)

KEYWORDS

Registries; chronic disease; quality assurance, health care; primary health care; family practice; hospital information systems; medical records systems, computerized

Introduction

With the publication of the Institute of Medicine's *Crossing the Quality Chasm* report [1], physicians and health care delivery systems have sharpened their focus on the care of populations and panels of patients. Managing the health of populations demands the ability to identify individuals within the population with specific clinical or sociodemographic characteristics. The core of population management is the patient registry—a list of multiple patients who share some common clinical trait, such as being overdue for a laboratory test, receiving a medication, or having a particular diagnosis.

Registries play a key part in the Chronic Care Model [2,3], a broadly promulgated framework for managing patients with chronic illnesses. In the traditional paradigm of care, a physician reviews the record of, and evaluates, one patient at a time. In the paradigm of the Chronic Care Model and population health management, the physician looks simultaneously at all of his or her patients with a particular diagnosis or combination of diagnoses. This activity is commonly, but not universally, carried out by utilizing an electronic health record (EHR) [4,5].

Laboratory result registries have been used for several purposes, but the most common is detection of patients overdue for screening tests [6-8]. Laboratory result registries can also serve as surrogates for, or adjuncts to, diagnosis-based registries [9,10]. In comparison, medication registries can be used to identify and communicate with patients who are receiving a medication for which some change is recommended or required, as in the case of a safety recall or the new availability of a more effective or less expensive alternative [11-13].

Taken together, this evidence suggests that diagnosis, medication and laboratory registries are essential and effective tools for improving the quality and safety of health care at the population level. A variety of studies at different sites, using different registries and with different disease foci have shown positive results [2-3,6-13]. It is not known, however, how easy or difficult it is for practices in the community to generate such registries. Furthermore, it is apparent that many studies demonstrating the potential for registries to improve care arise in settings with robust electronic health records and computerized provider order entry systems [4,6,8,11,12].

In order to understand better the registry generation capabilities of community ambulatory practices, as well as the relationship between EHR usage and registry capabilities, we undertook the present study. Our goal was to measure physicians' general abilities to perform registry functions in office practice and to explore further the hypothesis that use of electronic health records is associated with the ability to perform registry functions. This study is one aim of a larger study which used a variety of methods, including surveys, focus groups, direct observation, and quality assessment. The goal of the larger study was to measure adoption and use of EHRs in Massachusetts and to compare state-wide adoption to three specific communities in the state that were in the process of implementing community-wide electronic health records with information exchange. Other results of the larger study have been published previously [14,15].

Methods

Sampling

We carried out a statewide survey of physician practices in Massachusetts between June 2005 and November 2005. We began with a commercial database of physicians in Massachusetts (Folio Associates, Hyannis, MA) which contained contact information for 20,704 physicians practicing at 6308 distinct practice sites in the state. We drew a stratified random sample of practice sites from this database and selected one physician at random from each practice. Our sample was stratified by geography (urban vs nonurban based on county designation, except in the case where there were rural ZIP codes in urban counties, where ZIP codes were used instead), practice size (1 physician, 2 - 3 physicians, 4 - 6 physicians, and 7 or more physicians, exclusive of residents), and practice type (hospital-based primary care, hospital-based specialty/mixed, non-hospital-based primary care, or non-hospital-based specialty/mixed). These sampling characteristics were based on values in the commercial database.

Practices in rural parts of Massachusetts, primary care practices within hospitals, and large practices were oversampled by 100% in our sampling plan to ensure we had adequate representation of these particular practice types. Further details of the sampling plan have been reported previously [14,15].

Survey

Ultimately, we identified a sample of 1884 physician practices across the state of Massachusetts. We mailed a randomly chosen physician at each practice the survey and a US \$20 cash incentive to encourage participation. We contacted non-respondents by phone several times and also sent the survey to them two more times (without further cash incentive). The survey contained demographic questions (relating to practice size, teaching, and practice ownership), as well as a variety of questions about quality improvement, practice satisfaction, use of technology, and finances. Some of these questions have been analyzed as part of other aims of this study [14,15]. Two questions (practice size and type) were asked in the survey and were also present in the commercial database used for sampling. In the case that a practice's response differed from the commercial database (eg, because the practice had changed in size), the survey responses were used in analysis as they were more current.

The survey also asked questions about the use of registry functions and availability of an EHR (defined as "an integrated clinical information system that tracks patient health data, and may include such functions as visit notes, prescriptions, lab orders, etc") in the physician's practice. Specifically, physicians were asked to rate the ease of creating lists of patients by diagnosis or health risk (eg, diabetes), by laboratory results (eg, patients with abnormal hematocrit levels), and by medications they currently take (eg, patients on warfarin), using a five-point scale: very easy, somewhat easy, somewhat difficult, very difficult, and cannot generate. Furthermore, physicians were asked if their practice had components of an EHR, specifically defined as "an integrated clinical information system that tracks patient health data, and may include such functions as visit

notes, prescriptions, laboratory orders, etc" and were also surveyed on the availability and use of specific EHR components, such as structured problem or medication lists and electronic reporting and review of laboratory results. The survey instrument and study protocol were reviewed and approved by the Partners Healthcare Institutional Review Board (IRB). The instrument is available as an appendix to this article (See [Multimedia Appendix 1](#)).

Analysis

We used SAS 9.1.3 (SAS Institute, Cary, NC), applying weights throughout our analysis to control for both our stratified sampling plan (which included over-sampling of specific groups) and for variable response rates in different strata (specialty, category of practice size, hospital affiliation, and urban/rural location). We used frequency weights (fweights) which are the inverse of the response proportion for each stratum or, equivalently, the weights were determined by taking the population size for each stratum divided by the number of responses. Conceptually, the fweight for a particular response corresponds to the number of physician practices in Massachusetts that this response represents. The ultimate purpose of this weighting strategy was to make our results representative of the population of ambulatory care physician practices in Massachusetts.

We used logistic regression to assess the relationship between the presence of electronic health records in the practice and the ability to create diagnosis registries, laboratory test registries, and medication registries, adjusting each model for the following potential confounding factors:

- urban/rural location
- practice size
- practice ownership (owner, part-owner, non-owner)
- teaching status (whether any students or residents were present in the past year)
- hospital affiliation

- practice type (chosen from solo primary care practice, solo specialty care practice, primary care group/partnership, single specialty group/partnership, multi-specialty group/partnership)

In a secondary analysis limited to practices that had EHRs, we used chi-square tests to examine the relationship between use of key EHR features and the ability to generate each type of registry (diagnosis, laboratory test, medication). For the feature-specific analysis we looked at the effect of problem list use on the ability to generate diagnosis registries, electronic laboratory result review on laboratory test registries, and electronic medication list use on medication registries. In each one of the three cases, the associated feature was chosen for analysis because it was the most directly related feature to the registry type.

Results

A total of 1345 physicians (71%, 1345 of 1884) completed the survey, 1328 by mail and 17 by phone. There were no significant differences between respondents and non-respondents on the sampling characteristics (specialty, practice size, hospital affiliation, and rural practice). The practices reported using a wide variety of commercially available and self-developed EHR systems. [Table 1](#) shows the practice characteristics. Note that some respondents omitted certain questions on the survey so the number of practices does not always add up to 1345. The rural/urban classification was applied based on practice location, so it was available for all practices.

Among the 356 practices which had an EHR and reported its name, a total of 187 (52.5%) used one of the 4 most prevalent systems while the remaining 169 (47.5%) reported using one of 78 other systems that were named. There were also 31 practices that reported having an EHR but did not provide its name—they were still counted as having an EHR for purposes of the analysis.

Table 1. Practice characteristics

	n of practices	% of practices
Practice Location		
Rural	331	24.6
Urban	1014	75.4
Practice Size		
≥ 6 physicians	504	37.5
3 - 5 physicians	280	20.8
1 - 2 physicians	383	28.5
Practice Ownership		
Full owner	460	34.2
Part owner	172	12.8
Non owner	542	40.3
Practice Involvement in Teaching		
Involved in teaching	552	41.0
Not involved in teaching	628	46.7
Hospital Affiliation		
Hospital based practice	360	26.8
Non-hospital based practice	974	72.4
Practice Type		
Solo primary care practice	154	11.4
Solo specialty care practice	192	14.3
Primary care group/partnership	309	23.0
Single specialty group/partnership	338	25.1
Multi-specialty group/partnership	177	13.2
EHR Usage		
Yes	387	28.8
No	794	59.0

Overall, 79.8% of physicians reported that their practices could generate registries of patients with a particular diagnosis; 56.1% could generate registries of patients with a specific laboratory result; and 55.8% could generate registries of patients taking a particular medication. Among physicians who reported that their practices were able to generate registries, the reported ease

with which such registries could be generated varied greatly, as shown in [Table 2](#). While 38.9% of physician practices that could generate diagnosis registries said their practice could do it easily or very easily, considerably fewer said that it was easy or very easy to generate registries based on laboratory test results (14.5%) or medications (17.8%).

Table 2. Ease or difficulty of generating registries of patients based on diagnosis, laboratory result and medication use^a

Ease or Difficulty	Diagnosis registry	Laboratory result registry	Medication registry
Very Easy	15.0%	6.5%	7.4%
Somewhat Easy	23.9%	8.0%	10.4%
Somewhat Difficult	21.7%	16.0%	14.1%
Very Difficult	19.2%	25.6%	23.9%
Cannot Generate	20.2%	43.9%	44.2%

^aWeighted proportion of physicians reporting that their practice can generate each kind of registry with a particular ease or difficulty.

[Table 3](#) shows the proportion of physician practices that were capable of carrying out registry functions, stratified by pre-specified subgroups of interest. For all three registry types, providers with EHRs were significantly more likely to be able

to perform registry functions than providers using non-electronic record systems ($P < .001$ for all three registry types). Also, larger practices, practices involved in teaching, hospital-based practices, and primary care practices were more likely to be able to generate registries.

In logistic regression analyses controlling for urban/rural location, practice size, practice ownership, teaching status, hospital affiliation, and practice type, the relationship between

the presence of EHR and the ability to carry out each registry function remained robust. EHR adopters were more likely than non-adopters to be able to develop registries based on diagnosis (adjusted odds ratio [OR] 1.53, 95% confidence interval [CI] 1.25 - 1.86), laboratory results (OR 1.42, 95% CI, 1.22 - 1.66), and medications (OR 2.30, 95% CI, 1.96 - 2.70). Rural location, practice size, practice ownership, hospital affiliation, and practice type also remained significant correlates of one or more registry capability in the multivariate analyses (Table 4).

Table 3. Ability to perform registry functions according to practice characteristics (these data are weighted but not adjusted for confounding factors)

	Percentage of physicians able to perform function		
	Diagnosis registry	Laboratory result registry	Medication registry
Practice Location ^a			
Rural	78.9%	58.9%	62.6%
Urban	79.9%	55.9%	55.3%
Practice Size ^b			
≥ 6 physicians	82.4%	60.1%	59.2%
3 - 5 physicians	84.8%	63.4%	53.9%
1 - 2 physicians	75.8%	50.6%	54.8%
Practice Ownership ^c			
Full owner	77.8%	51.2%	54.1%
Part owner	81.6%	49.9%	48.3%
Non owner	81.6%	66.6%	61.7%
Practice Involvement in Teaching ^d			
Involved in teaching	85.0%	63.2%	61.8%
Not involved in teaching	76.9%	52.3%	52.5%
Hospital Affiliation ^e			
Hospital based practice	81.9%	69.9%	65.2%
Non-hospital based practice	79.4%	54.5%	54.8%
Practice Type ^d			
Solo primary care practice	74.2%	57.2%	61.7%
Solo specialty care practice	77.1%	47.0%	55.8%
Primary care group/partnership	88.6%	71.4%	67.2%
Single specialty group/partnership	78.5%	56.0%	44.5%
Multi-specialty group/partnership	84.5%	58.7%	59.7%
EHR Usage ^d			
Yes	85.9%	66.7%	71.6%
No	78.0%	52.9%	51.1%

^a $P = .64$ using a chi-square test for diagnosis registry functions, $P = .25$ for laboratory registry functions, $P = .005$ for medication registry functions.

^b $P < .001$ for diagnosis and laboratory registry functions, $P = .009$ for medication registry functions.

^c $P = .004$ for diagnosis registry functions and $P < .001$ for laboratory and medication registry functions.

^d $P < .001$ for all three registry types.

^e $P = .153$ for diagnosis registry functions, $P < .001$ for medication and laboratory registry functions.

Table 4. Multivariate correlates of registry function capability

	Diagnosis registry OR (95% CI)	Laboratory result registry OR (95% CI)	Medication registry OR (95% CI)
Practice Location			
Rural	0.95 (0.73 - 1.23)	1.08 (0.86 - 1.34)	1.32 (1.05 - 1.65)
Urban	1 (Ref)	1 (Ref)	1 (Ref)
Practice Size			
≥ 6 physicians	1.09 (0.85 - 1.41)	0.92 (0.74 - 1.14)	1.10 (0.89 - 1.37)
3 - 5 physicians	1.62 (1.29 - 2.03)	1.39 (1.16 - 1.68)	1.20 (1.00 - 1.44)
1 - 2 physicians	1 (Ref)	1 (Ref)	1 (Ref)
Practice Ownership			
Full owner	1.28 (1.04 - 1.58)	0.71 (0.60 - 0.85)	0.65 (0.54 - 0.78)
Part owner	1.19 (0.92 - 1.53)	0.53 (0.43 - 0.65)	0.71 (0.58 - 0.87)
Non owner	1 (Ref)	1 (Ref)	1 (Ref)
Practice Involvement in Teaching			
Involved in teaching	1.42 (1.19 - 1.70)	0.98 (0.85 - 1.13)	1.08 (0.94 - 1.25)
Not involved in teaching	1 (Ref)	1 (Ref)	1 (Ref)
Hospital Affiliation			
Hospital based practice	1.09 (0.85 - 1.39)	1.87 (1.52 - 2.30)	1.35 (1.10 - 1.66)
Non-hospital-based practice	1 (Ref)	1 (Ref)	1 (Ref)
Practice Type			
Solo primary care practice	0.67 (0.47 - 0.96)	1.23 (0.92 - 1.65)	1.97 (1.46 - 2.66)
Solo specialty care practice	0.84 (0.60 - 1.17)	0.90 (0.68 - 1.18)	1.78 (1.35 - 2.34)
Primary care group/partnership	1.47 (1.07 - 2.02)	1.90 (1.50 - 2.42)	1.61 (1.26 - 2.04)
Single specialty group/partnership	0.70 (0.54 - 0.91)	0.91 (0.74 - 1.13)	0.62 (0.50 - 0.77)
Multi-specialty group/partnership	1 (Ref)	1 (Ref)	1 (Ref)
EHR Usage			
Yes	1.53 (1.25 - 1.86)	1.42 (1.22 - 1.66)	2.30 (1.96 - 2.70)
No	1 (Ref)	1 (Ref)	1 (Ref)

OR = adjusted odds ratio; CI = confidence interval

Ref: Reference Category

We also observed a relationship between use of key EHR features and ability to perform related registry functions. Specifically, within the group of physicians who had access to an EHR in their practice, 90.4% of physicians who reported using an electronic problem list at least some of the time had the ability to perform diagnosis registry functions, while only 67.7% of physicians using an EHR without access to an electronic problem list could perform these functions ($P < .001$). Similarly, while 75.2% of physicians who used an electronic medication list could perform medication registry functions, only 53.0% of physicians who used an EHR without a medication list reported they could perform them ($P < .001$). Finally, while 71.5% of physicians who used their EHR to view laboratory results could perform laboratory registry functions, only 33.9% of physicians whose EHR could not be used to view laboratory results reported they could perform these functions ($P < .001$).

Discussion

Principal Results

While many studies have demonstrated the value of being able to perform registry functions for improving the quality and safety of health care [2-3,6-13], few data are available regarding the capability of carrying out registry functions in community-based practices. In this study, about 80% of physicians reported being able to generate registries of patients according to diagnoses, but nearly half of all physicians in ambulatory care practice in Massachusetts could not create registries by medication or laboratory result.

Having EHRs was strongly associated with the reported ability to generate registries based on diagnosis, laboratory test result, or medication, but even among EHR users, 14% could not generate lists of diagnoses, 33% could not do so for laboratory

tests, and 28% could not do so for medications. Furthermore, we found that physicians who reported active use of key EHR functions were considerably more likely to report being able to generate registries. Thus, these data suggest that EHRs appear important for delivering care using registries, and that most but not all EHR users could generate registries using their electronic records.

We are uncertain about why some physician practices with EHRs were unable to create registries. Many of these practices reported using EHRs which we knew to have this capability. It is likely that at least some of the EHR users who reported an inability to generate registries actually have the ability to generate them using their EHR, but are unaware of the feature. This suggests that improvements in documentation, training, and ease of use to help more physicians take advantage of the existing registry capabilities of their EHRs may be useful.

We were also a bit surprised by the relatively high proportion of EHR non-users who reported being able to generate registries. We are uncertain as to the mechanism employed by these users, since our survey did not ask them to explain how they were generating registries. These users may have been using retrospective chart review, prospective tracking, or analysis based on administrative data (such as billing and claims data in a practice management system). Each of these methods has a significant downside. Retrospective chart review is extremely time-consuming and error prone; prospective tracking requires criteria to be developed in advance; and non-clinical data are often less sensitive and specific than clinical data.

Taken together, our findings raise concerns about the ability of many ambulatory care practices, particularly the majority of practices without EHRs, to provide effective care for their patients on a population level. Physicians and practices need to consider population-level care management, not only as an essential component to effective practice within the Chronic Care Model [2,3], but also as a necessary tool for responding to the exigencies of forces driving quality improvement, such as pay-for-performance. Without EHRs, and without active and effective use of key features in robust EHRs [16], physicians and practices will have much greater difficulty in efficiently delivering safe and effective care for patients with acute and chronic problems.

EHRs can and should either include the inbuilt ability to query across patient records by a variety of criteria, or support extracting patient data which can be fed into other applications which do this. The ability to generate registries by diagnosis is common in many commercial EHR systems. In fact, it is a requirement of the 2007 and 2008 Certification Commission for Health Information Technology (CCHIT) criteria for ambulatory EHR certification [17]. However, such certification is voluntary, and many commercial products are not certified, or are certified under the older 2006 criteria, which did not require the availability of registry functions. CCHIT certification is valid for only three years, however, so as products certified under the 2006 criteria are re-certified under the current criteria, this gap will close. Also, although certification is voluntary, there are increasing incentives (or, in some limited cases, mandates) encouraging use of CCHIT-certified products, so the

use of non-certified products which may not have registry capabilities is likely to be lessened over time [18]. Furthermore, just because an EHR can be used to generate a registry, the software does not always make it easy or user-friendly for a provider to do so. However, even when EHRs do not come "pre-loaded" with the ability to generate registries, the organization of patient information in structured fields in an electronic format facilitates the creation of registries more easily than in paper-based systems.

It is also worth noting that the ability to create registries in an EHR is generally predicated on the use of structured documentation features within the EHR. For example, if a clinician documents patient problems only in unstructured clinical notes, it is nearly impossible in most commercially available EHRs to build medication registries based on this unstructured information. However, if the clinician uses a structured medication list with a controlled medication vocabulary, generating such a registry becomes much easier.

Finally, our survey was conducted in 2005. Since then, adoption of EHRs in ambulatory practices has increased somewhat [19], though is still far from universal. Since our data suggest that physicians with EHRs are more able to generate registries than non-users, we expect that the current ability of physicians to generate such registries is likely now higher than it was in 2005. Moreover, the capabilities of EHR systems, as a rule, increase over time, so we likewise expect that EHR users are more likely, today, to be able to generate registries than they were in 2005.

Implications

Our findings have several important implications for physicians, for the health care system, and for developers of electronic health records. Because our findings suggest that the ability to generate registries is less than universal, and because generating registries is integral to quality and safety enhancing activities, it may be necessary to take steps to increase these capabilities in office practice. Providing physicians and practices with training and activities to increase awareness of the role of generating registries may be beneficial, but these changes are unlikely to be sufficient. Incentives also likely play an important role; physicians are more likely to adopt and use registry functions if financial incentives are in place to do so [20]. As pay-for-performance initiatives become more prevalent, physicians are likely to embrace the use of registries as a foundation for building practice-level population management capabilities. Our findings also show that EHR users are more likely to be able to generate registries than non-users, so incentives aimed at EHR adoption alone are also likely to have a positive effect on registry capabilities.

Limitations

The study has several limitations. First, our survey was limited to physicians in ambulatory care in Massachusetts, and the results may not be generalizable to other states or regions. However, given that Massachusetts is a state in which more than 45% of physicians have EHRs [15], the large proportions of physicians and practices reporting inability to perform registry functions in this study are likely to be even larger in other states, where EHR adoption has lagged. As such, the need to consider

efforts to adopt EHRs and expand their use may have even greater imperative in other regions.

Another limitation is the self-reported nature of survey studies such as ours. We are, of course, not truly measuring physicians' abilities to generate registries, but instead their self-reports of the ability to generate three specific types of registries. This raises the possibility of social desirability bias influencing physicians' responses to survey questions. However, if this bias were present, then one might expect that physicians overestimated their abilities to perform registry functions, which would mean that even fewer physicians than reported have the ability to generate registries. Also, our survey asked providers how easily they could perform registry functions but did not ask how frequently they actually did perform such functions, or for what purposes and with what results. It is important to note that, among those practices that reported the ability to make these registries, we do not know the frequency with which they did so. It is possible that some practices, although able to create the registries, never actually do. Future qualitative and quantitative studies should explore how physicians and practices are using registries, as well as the barriers to, and facilitators of, effective use of these important tools. Intervention studies will then be able to test strategies for improving physicians' use of registries to improve quality of care and patient safety.

Acknowledgments

This study was funded in part by the United States Agency for Healthcare Research and Quality cooperative agreement #1UC1HS015397-01 and the Massachusetts e-Health Collaborative. The Agency for Healthcare Research and Quality and the Massachusetts e-Health Collaborative had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Agency for Healthcare Research and Quality or the Massachusetts e-Health Collaborative.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Massachusetts Survey of Physicians and Computer Technology

[[PDF file \(Adobe PDF, 134 KB\)](#)-Multimedia Appendix 1]

References

1. ; Committee on Quality of Health Care in America, Institute of Medicine. Crossing the Quality Chasm: A New Health System for the 21st Century. Washington, DC. The National Academies Press; 2001.
2. Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness: the chronic care model, Part 2. *JAMA*. Oct 16, 2002;288(15):1909-1914. [Medline: [12377092](#)] [doi: [10.1001/jama.288.15.1909](#)]
3. Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness. *JAMA*. Oct 9, 2002;288(14):1775-1779. [Medline: [12365965](#)] [doi: [10.1001/jama.288.14.1775](#)]
4. Thomas KG, Thomas MR, Stroebel RJ, et al. Use of a registry-generated audit, feedback, and patient reminder intervention in an internal medicine resident clinic—a randomized trial. *J Gen Intern Med*. 2007;22(12):1740-1744. [doi: [10.1007/s11606-007-0431-x](#)]
5. Wagner EH. Population-based management of diabetes care. *Patient Educ Couns*. Sep 1995;26(1-3):225-230. [Medline: [7494727](#)] [doi: [10.1016/0738-3991\(95\)00761-N](#)]
6. Dorr DA, Wilcox A, Donnelly SM, Burns L, Clayton PD. Impact of generalist care managers on patients with diabetes. *Health Serv Res*. Oct 2005;40(5 Pt 1):1400-1421. [[FREE Full text](#)] [Medline: [16174140](#)] [doi: [10.1111/j.1475-6773.2005.00423.x](#)]

7. Jung E, Li Q, Mangalampalli A. Report Central: quality reporting tool in an electronic health record. In: AMIA Annu Symp Proc. 2006. Presented at: :971. [Medline: [17238590](#)]
8. Olsha-Yehiav M, Einbinder JS, Jung E, et al. Quality Dashboards: technical and architectural considerations of an actionable reporting tool for population management. AMIA Annu Symp Proc. 2006:1052. [[FREE Full text](#)] [Medline: [17238671](#)]
9. McCulloch DK, Price MJ, Hindmarsh M, Wagner EH. A population-based approach to diabetes management in a primary care setting: early results and lessons learned. Eff Clin Pract. 1998;1(1):12-22. [[FREE Full text](#)] [Medline: [10345254](#)]
10. Miller DR, Safford MM, Pogach LM. Who has diabetes? Best estimates of diabetes prevalence in the Department of Veterans Affairs based on computerized patient data. Diabetes Care. May 2004;27(Suppl 2):B10-B21. [[FREE Full text](#)] [Medline: [15113777](#)] [doi: [10.2337/diacare.27.suppl_2.B10](#)]
11. Cheetham TC, Chan J, Benson V, Richmond C, Levin E, Campen D. Successful conversion of patients with hypercholesterolemia from a brand name to a generic cholesterol-lowering drug. Am J Manag Care. Sep 2005;11(9):546-552. [[FREE Full text](#)] [Medline: [16159044](#)]
12. Jain A, Atreja A, Harris CM, Lehmann M, Burns J, Young J. Responding to the rofecoxib withdrawal crisis: a new model for notifying patients at risk and their health care providers. Ann Intern Med. Feb 1, 2005;142(3):182-186. [[FREE Full text](#)] [Medline: [15684206](#)]
13. Usher-Smith JA, Ramsbottom T, Pearmain H, Kirby M. Evaluation of the cost savings and clinical outcomes of switching patients from atorvastatin to simvastatin and losartan to candesartan in a Primary Care setting. Int J Clin Pract. Jan 2007;61(1):15-23. [Medline: [17229176](#)] [doi: [10.1111/j.1742-1241.2006.01217.x](#)]
14. Simon SR, Soran CS, Kaushal R. Physicians' usage of key functions in electronic health records from 2005 to 2007: a statewide survey. J Am Med Inform Assoc. Apr 23, 2009. [Epub ahead of print. [Medline: [19390104](#)]
15. Simon SR, Kaushal R, Cleary PD, et al. Physicians and electronic health records: a statewide survey. Arch Intern Med. 2007;167(5):507-512. [Medline: [17353500](#)] [doi: [10.1001/archinte.167.5.507](#)]
16. Frieden TR, Mostashari F. Health care as if health mattered. JAMA. Feb 27, 2008;299(8):950-952. [Medline: [18314438](#)] [doi: [10.1001/jama.299.8.950](#)]
17. ; Certification Commission for Health Information Technology. Final Criteria for 2007 Certification of Ambulatory EHRs. Certification Commission for Health Information Technology. 2007. URL: http://www.cchit.org/files/Ambulatory_Domain/CCHIT_Ambulatory_FUNCTIONALITY_Criteria_2007_Final_16Mar07.pdf [accessed 2008 Feb 19] [[WebCite Cache ID 5cOCkQmJL](#)]
18. ; Certification Commission for Health Information Technology. The CCHIT Incentive Index. Certification Commission for Health Information Technology. 2008. URL: <http://ehrdecisions.com/wp-content/files/CCHITIncentiveIndex20080925.pdf> [accessed 2009 Feb 28] [[WebCite Cache ID 5ew7Yky6p](#)]
19. DesRoches CM, Campbell EG, Rao SR, et al. Electronic health records in ambulatory care—a national survey of physicians. N Engl J Med. 2008;359(1):50-60. [Medline: [18565855](#)] [doi: [10.1056/NEJMsa0802005](#)]
20. Campbell S, Reeves D, Kontopantelis E, Middleton E, Sibbald B, Roland M. Quality of primary care in England with the introduction of pay for performance. N Engl J Med. Jul 12, 2007;357(2):181-190. [Medline: [17625132](#)] [doi: [10.1056/NEJMsr065990](#)]

Abbreviations

CCHIT: Certification Commission for Health Information Technology

EHR: electronic health record

IRB: institutional review board

Edited by K El Emam; submitted 21.Nov.2008; peer-reviewed by V Harrop, S Woods; comments to author 22.Feb.2009; revised version received 04.May.2009; accepted 23.May.2009; published 10.Aug.2009

Please cite as:

Wright A, McGlinchey EA, Poon EG, Jenter CA, Bates DW, Simon SR

Ability to Generate Patient Registries Among Practices With and Without Electronic Health Records

J Med Internet Res 2009;11(3):e31

URL: <http://www.jmir.org/2009/3/e31/>

doi: [10.2196/jmir.1166](#)

PMID: [19674961](#)

© Adam Wright, Elizabeth A McGlinchey, Eric G Poon, Chelsea A Jenter, David W Bates, Steven R Simon. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 10.Aug.2009. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted

use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.